



Carbon Stock Assessment of Four Selected Agroforestry Systems in Owerri-West Local Government Area, Nigeria

Chris O. Nwoko ^{a*}, Samuel Umeohana ^a,
Jonathan Anyanwu ^a, Lucy Izunobi ^a
and Chidinma A. Peter-Onoh ^b

^a Department of Environmental Management, Federal University of Technology, Owerri, Nigeria.

^b Department of Crop Science and Technology, Federal University of Technology, Owerri, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2024/v14i34033

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/113905>

Original Research Article

Received: 23/12/2023

Accepted: 27/02/2024

Published: 04/03/2024

ABSTRACT

Aim: The aim of the study is to generate data on the carbon stock in four predominant agroforestry systems in the southeast Nigeria. Agroforestry (AF) system studied were oil palm, plantain, Bush mango and mixed agroforestry systems.

Methods: Methodologies adopted were measurement of diameter at breast height (dbh) of woody species, soil and litter sampling and the use of allometric equations in the determination of biomass carbon.

Results: The results obtained showed that total agroforestry carbon ranged from $160.2 \pm 102.2 \text{ Mg C ha}^{-1}$ to $130.7 \pm 93.2 \text{ Mg C ha}^{-1}$. Mixed AF system recorded the highest total biomass carbon of $8.39 \pm 1.4 \text{ Mg C ha}^{-1}$ and lowest in bush mango AF. There was significant positive relationship between

*Corresponding author: E-mail: chrisnwoko3@gmail.com;

total biomass carbon and species richness and total agroforestry carbon. Also, SOC positively with all the ecological parameters examined. Soil organic carbon contributed significantly to the total agroforestry carbon recorded in this study.

Conclusion: With the level of carbon stock observed in the biomass and soils of the agroforestry systems in the region, agroforestry practices should be encouraged to boost climate change mitigation.

Keywords: Carbon stock; agroforestry systems; farming practice.

1. INTRODUCTION

“Agroforestry is one of the land-use management systems whereby trees are planted with agricultural crops. This farming practice plays very important role in sustaining ecosystem goods and services and in mitigating climate change. Agroforestry systems absorb excess carbon dioxide from the atmosphere for the process of photosynthesis. Carbon is stored in tree biomass and in soil that helps protect natural carbon sinks through the improvement of land productivity and the provision of forest products on agricultural lands” [1].

“Agroforestry practice is seen as one of the greenhouse gas reduction strategies under the Kyoto Protocol, which was adopted in 1997. As a result, the sequestration potential of agroforestry systems has received attention by the global community and many countries are currently adopting the practice in their climate change mitigation strategy” [2,3].

“Globally, the land areas covered with various types of Agroforestry systems is estimated to be around 1.6×10^9 hectares, with an aboveground biomass sequestration potential of 1.1–2.2 billion t C in the coming 50 years” [4]. “The carbon sequestered in an agroforestry system mainly depends on the species diversity, arrangement, and role of components within the system, which in turn is determined by the ecological, socioeconomic aspects and previous land use practices” [5,1].

“The capacity of agroforestry systems to store C differs across different agroecological landscapes”. [6] Montagnini, and Nair, [7] reported that “storage potential for semiarid, sub-humid, humid, and temperate regions was estimated at 9, 21, 50, and 63 t C ha⁻¹, respectively”. Also extensive reviews were carried out by [8] for “West African countries (extending from Arid Sahara Desert to humid region Guinea) and they reported that biomass C stocks ranged from 22.2 to 70.8 t C ha⁻¹”.

“There are many similar studies conducted in various regions of the continents” [9]. “From global report, the total biomass C stock for agroforestry systems ranges between 12 and 228 t C ha⁻¹” [1,10] also reported that “agroforestry practices stored C ranging from 0.29 to 15.21 t C ha⁻¹ yr⁻¹ in their aboveground biomass and can have from 30 to 300 t C ha⁻¹ in their soil down to one-meter depth. Soil C stock for the 0–60 cm soil layer differs among different land uses and regions”. Nair et al. [11,12] noted that “C stock in the 0-60 cm soil layer is 121–123 t ha⁻¹ for tropical forests and 110–117 t ha⁻¹ for tropical savanna”. “The results obtained from various studies on agroforestry practices in diverse ecological conditions showed that tree-based agricultural systems, compared to treeless systems, stored more carbon in deeper soil layers near the tree than away from the tree; higher soil organic carbon content was associated with higher species richness and tree density” [11].

“Agroforestry systems promote efficient water utilization in smallholder farmers in rural communities, improve microclimate, enhance soil productivity and nutrient cycling, improve farm productivity, and increase farm income as well as sequestering carbon” [13]. “There is need for better understanding of above-ground tree biomass and soil organic carbon stocks for proper management of the carbon pools in agroforestry systems. However, extensive research is required for agroforestry systems to be used in global agendas of carbon sequestration” [11].

The objective of this research work was to assess the carbon stock in selected agroforestry systems in Owerri-West Local Government Area in Imo state, Nigeria.

2. MATERIALS AND METHODS

This research was carried out in Owerri-West Local Government Area (LGA) Imo state, Southeast Nigeria (Fig. 1). which is located

between 6°59'19.1" and 6°54'42.3"E longitude and between 5°31'43.8" and 5°16'49.2"N latitude. Fig. 2 shows a satellite imagery of the Owerri-west Local government area. The soil formation which is Pliocene to Miocene in age consists of coastal plain sands, that is about 0.05–2.0 mm in size, with minor clay beds of isolated gravels, conglomerates, and very coarse

sandstone in some places. Benin Formation is overlain by alluvium deposits and underlain by Ogwashi-Asaba formation which consists of lignite, sandstones, clays and shale. The area has humid tropical climate, with mean annual precipitation of 2200 mm and average annual temperature of 30° C. Rain fall pattern is bimodal with peaks in July and August.




Fig. 1. Map of Owerri west local government area showing the major towns




Fig. 2. Land use satellite imagery of Owerri west Local government area

Table 1. Land use of the study area

s/n	Study area	Area covered (sq.km)	%
1	Built Up	68.5	23.3
2	Vegetation	50.2	17
3	Bare Soil/Cultivated Land	80.5	27.3
4	Forest	95.4	32.4
	Total area	294.6	100

Table 2. Characteristics and location of the agroforestry system of the study area

Characteristics	Agroforestry systems			
	Oil palm tree AFS	Bush mango AFS	Plantain AFS	Mixed multistory AFS
Location	Ndigwu	Obinze	Eziobodo	Avu/ Umuguma
Topography	Gentle slope	Flat land	Gentle slope	Flat land
Dominant plant species	Oil palm tree, cassava, <i>Albizia amara</i> (Roxb.)	Bush mango, elephant grass	Plantain plant, <i>Pleiospermium alatum</i>	<i>Azadirachta indica</i> , <i>Magnifera indica</i> , <i>Trichilia prieuriana</i>
Handling activities	Tree trimming, ripping of unwanted plants,	Trimming, , pollarding, Farm house waste	Tree trimming, ripping of unwanted plants, Farm house waste	Tree trimming, pollarding

The entire area of the zone is 294.6 sq.km, which is divided into different land-use types, such as agricultural land and agroforestry together, accounts for 27.3%, along with natural forest (32.4%) built up area (23.3% and vegetation 17% (Table 1).

“The soil type is Nitisol and the texture is predominantly clay. The livelihood of the community under this district is mainly dependent on agroforestry practices and paid employment. They use mixed farming, including non-fruit trees, fruit trees, crops, vegetables, spices production, and very limited animal husbandry focused on, goats, piggery, and poultry”. [6] Although there are several types of agroforestry systems practiced by the farmers, the main and most common which were used for this study were (1) *Elaeis guineensis* (oil palm tree) based agroforestry system (2) *Irvingia gabonensis* (bush mango) based agroforestry system and (3) Plantain (*Musa paradisiaca*) agroforestry system and (4) mixed multistory agroforestry system (Table 2).

These agroforestry systems identified for sampling are located at Ndigwu, Obinze, Eziobodo and Avu / Umuguma. The topography ranged from gentle slope to flat and AF systems are mainly maintained by occasional trimming of over grown plants and cutting down dead trees.

2.1 Sample Collection

The agroforestry systems with similar gradient and altitudes within Owerri-west landscape were identified and mapped out for data collection. In each agroforestry site a randomly nested quadrat of 20 x 20 m size was mapped out for the inventory of trees/shrubs. All woody species including fruit trees and non- fruit trees with DBH ≥2.5cm diameter and height ≥1.5m was measured and recorded. The inventory was aimed to obtain data for the biomass estimation. The quadrats size in this study corroborate with previous work in the literature [2]. Within each quadrant, three 50 x 50 cm small plots for litter sampling were laid out. In addition, soils (0-20cm, 20-40cm) were collected using soil auger, from five different points and bulked to form composite soil sample. A total of 140 composite samples were collected from the three subplots. The soils were sieved through a 5-mm mesh screen and with 0.02 mm size and mixed to a uniform color and consistency then a subsample of 500g was taken for carbon analysis. Soil samples were taken to Environmental Management Central laboratory, Federal University of Technology, Owerri for Soil Organic Carbon (SOC) analysis using Walkley-Black method and an additional 80 soil samples were collected using soil core sampler for bulk density determination (g cm⁻³) [14] The soil C stocks (Mg

C ha⁻¹) were computed by multiplying the C content (%), bulk density (g cm⁻³), and layer thickness (cm).

2.2 Biomass and Carbon Stock Determination

Estimation of total above ground biomass (AGB) and below ground biomass (BGB) for tree and shrubs were done for each of the 4 selected agroforestry systems. The biomass estimation was done using nondestructive method which involves use of allometric equations. This is because destructive methods of tree harvesting is costly, time consuming and labour intensive.

To compute the AGB and BGB, and their respective carbon stock, allometric equation developed by [15] was used. Tree/shrub biomass was converted to C by multiplying the above-ground biomass by 0.5

$$TCS = (AGB + BGB) \times 0.5 \quad 1$$

TCS= total carbon stock, AGB= above ground biomass, BGB= below ground biomass and 0.5 conversion factor. Earlier researchers have come to the conclusion that carbon stock is 50% of the total biomass of a tree [16-19].

The AGB was estimated using the allometric equation thus.

$$AGB = 0.091 \times d^{2.472}; R^2 = 0.78, n=60 \quad 2$$

Where,

AGB in (dry mass per tree in kg, 'd' is diameter at breast height in cm. previous authors revealed that AGB is strongly correlated with tree diameter [20-22].

$$BGB = 0.49 \times AGB^{0.923}; R^2 = 0.90, n=60 \quad 3$$

Where,

BGB (dry matter per tree in kg)

Bulk density (BD) and soil organic carbon:

SOC stock (Mg C ha⁻¹) was calculated by multiplying the concentrations (%) of soil carbon, the Bulk density (g cm⁻³) and depth of the sampled soil [23].

$$SOC \text{ (Mg C ha}^{-1}\text{)} = BD \text{ (g cm}^{-3}\text{)} * \% \text{ C} * \text{soil depth (cm)} \quad 4$$

For carbon litter determination a small sample (2 grams) of each one of the herbaceous vegetation and litter layer was analyzed for carbon content determination using Walkley-Black method. Carbon storage in herbaceous vegetation and litter layer was computed using the formula [24].

$$C \text{ stored (Mg C ha}^{-1}\text{)} = \text{Total dry weight} * \text{C content}$$

2.3 Vegetation Analysis

In this study, species richness, evenness and species diversity were determined.

Species richness(Y) = S/√N, where Y = Species richness, S= Number of species, N = Total number of individuals.

$$\text{Species diversity Shannon-Wiener} = H' = - \sum_{i=1}^s p_i \log p_i$$

where s is the total number of species in the sample, i is the total number of individuals in one species, p_i (a decimal fraction) is the number of individuals of one species in relation to the number of individuals in the population, and the log is to base-2 or base-e. Equitability (Evenness) was calculated by using Simpson's index and expressing it as a proportion of the maximum value.

2.4 Statistical Analysis

The vegetation composition of the different agroforestry systems were compared. Analysis of Variance (ANOVA) (α=0.05) was performed to assess the differences in vegetation composition, biomass and soil carbon stock in the agroforestry system under study. Regression analysis was used to test the relationship between tree carbon stock with tree species richness- and tree density.

3. RESULTS AND DISCUSSION

The highest number of species, genera and family were recorded in the mixed agroforestry system. These species were composed of trees, shrubs and crop components. Bush mango AFS has 13 different species and 10 genera and forms the second largest diverse agroforestry system (Table 3). Oil palm recorded the least number of species and genera in the AFS.

Table 3. The species, genera and their families identified in various agroforestry systems in the study area

AFS	Species	Genera	families
Oil palm	10	8	5
Bush mango	13	10	6
Plantain	11	9	7
Mixed AF	16	11	9

Table 4. Summary of the mean (\pm Std) vegetation characteristics of the agroforestry systems

AFS	Species richness	Species diversity (H')	Equitability
Mixed AF	3.5 \pm 1.3a	0.78 \pm 0.02a	0.84 \pm 0.2b
Bush mango	3.4 \pm 1.7a	0.68 \pm 0.01a	0.92 \pm 0.6a
Plantain	1.7 \pm 0.9b	0.34 \pm 0.02b	0.56 \pm 0.2c
Oil palm	1.2 \pm 0.3c	0.21 \pm 0.03c	0.43 \pm 0.3d
p-value	0.05	0.001	0.001

Same letters within column are not significantly different

Table 5. The Mean (\pm sd) tree species density, DBH and height of agroforestry practices

AFS	Tree stand (ha ⁻¹)	DBH(cm)	Height (m)	BA(m ² ha ⁻¹)
Oil palm	12.3 \pm 1.2c	11.2 \pm 3.4b	13.2 \pm 5.6b	2.5 \pm 0.22c
Bush mango	14.8 \pm 3.5b	9.2 \pm 4.5c	11.3 \pm 6.3c	2.3 \pm 0.42c
Plantain	8.2 \pm 2.1d	8.1 \pm 2.3d	9.4 \pm 4.2d	3.4 \pm 1.3b
Mixed AF	45.3 \pm 13.4a	12.8 \pm 5.3a	15.2 \pm 5.6a	16.2 \pm 2.3a
p-value	0.05	0.05	0.02	0.05

Same letters within column are not significantly different

Table 6. Mean (\pm standard deviation; n=3) biomass carbon, soil carbon (SOC), litter and agroforestry system total carbon stocks (Mg ha⁻¹) for each of the four studied agroforestry practices) and results of 1-way ANOVA (at α =0.05)

Biomass	Oil palm AF (Mg ha ⁻¹)	Bush mango AF (Mg ha ⁻¹)	Plantain AF (Mg ha ⁻¹)	Mixed AF (Mg ha ⁻¹)	P=0.05
Above ground biomass(AGB)	3.8 \pm 1.2	2.5 \pm 1.3	3.4 \pm 0.7	4.5 \pm 2.3	0.02
Below ground (BGB)	2.3 \pm 1.9	1.23 \pm 0.8	2.34 \pm 0.89	3.89 \pm 1.3	0.067
Biomass total	6.1 \pm 2.2 ^b	3.73 \pm 0.7 ^c	5.74 \pm 1.2 ^b	8.39 \pm 1.4 ^a	0.022
Litter	0.42 \pm 0.02 ^c	1.03 \pm 0.05 ^a	1.34 \pm 0.65 ^a	0.98 \pm 0.2 ^b	0.12
SOC(0-20cm)	76.4 \pm 12.5	77.4 \pm 34.2	78.3 \pm 23.1	89.5 \pm 23.4	0.065
SOC(20-40cm)	47.8 \pm 9.4	54.3 \pm 13.4	52.2 \pm 10.5	61.3 \pm 12.7	0.043
Agroforestry total	130.7 \pm 93.2 ^b	136.5 \pm 89.2 ^b	137.6 \pm 79.3 ^b	160.2 \pm 102.2 ^a	0.032

Same letters within rows are not significantly different at 5% probability level

Table 7. Relationship between ecological characteristics and carbon stock in the AFS

Ecological characteristics	DBH	SOC	TBC	TAFC
Richness	-0.57*	0.21	0.56*	0.66**
Shannon diversity (H')	-0.68**	0.32	0.19	-0.32
Equitability	-0.66*	0.23	0.21	0.41

DBH=diameter at breast height, SOC= soil organic carbon, TBC=total biomass carbon, TAFC=total agroforestry carbon. * Significant at 5% probability

The agroforestry systems studied had a total of 50 plant species, 38 genera and 27 families. All the agroforestry systems differed ($p < 0.05$) in terms of their species richness, diversity and equitability (Table 4). The Shannon diversity index of mixed agroforestry was significantly higher than other AF. However, Bush mango number of individual species and equitability differed among plantain and mixed AF systems. Species richness and Shannon diversity index didn't differ for the mixed and Bush mango AF systems.

The tree stand, diameter at breast height (DBH) and basal area of the mixed AF were higher than other AF systems and differed significantly with other AFS (Table 5) the mean diameter at breast height ranged from 9.2 cm in bush mango AF to 12.8cm in mixed AF. The least mean tree stand occurred in plantain AF.

3.1 Biomass Carbon Stock

The mean values of above and below ground carbon stock are presented in Table 6. The total biomass carbon stock ranged from $3.73 \pm 0.7 \text{ Mg C ha}^{-1}$ to $8.39 \pm 1.4 \text{ Mg C ha}^{-1}$. The bush mango AF biomass carbon significantly differed from oil palm, plantain and mixed AF systems. The total biomass carbon in the mixed AF was higher than all the other AFS. The litter biomass at the plantain AF was higher than other AF systems and differs significantly from oil palm and mixed AF.

Soil organic carbon of the top soil (0-20cm) ranged from $76.4 \text{ Mg C ha}^{-1}$ to $89.5 \text{ Mg C ha}^{-1}$. The highest was recorded at mixed AF and the least was at oil palm AF. The sub-soil organic carbon (20-40cm) also ranged from $47.8 \text{ Mg C ha}^{-1}$ to $61.3 \text{ Mg C ha}^{-1}$ and highest value recorded in mixed AF and the least in oil palm AF. The total agroforestry biomass carbon was highest in mixed AF ($160.2 \text{ Mg C ha}^{-1}$) and lowest in Oil palm AF. The mixed AF biomass carbon differed significantly from other agroforestry systems studied. However, total biomass carbon in other agroforestry systems didn't differ significantly ($P > 0.05$).

The spearman correlation between the ecological characteristics and carbon stock is presented in Table 7. There were significant negative relationship between the diameter at breast height and the species richness, Shannon diversity index and equitability. The soil organic carbon had positive relation with all the

ecological parameters but the relationship was not significant ($p > 0.05$). Total biomass carbon and total agroforestry carbon had positive and significantly correlated with species richness.

3.1 DISCUSSION

3.1.1 Vegetation characteristics

From the study, the highest species richness and species diversity recorded in mixed AF is similar to results obtained in previous studies in south east Nigeria and some parts of western Nigeria [25] in natural agroforestry systems but lower than results obtained in north central of Nigeria. This could be due to environmental variability such as altitude, soils, topography, species adaptability and management strategy [26]. Relatively high species richness recorded in Bush mango AF may be as a result of large number of exotic and indigenous tree species present. Although the species evenness was highest in bush mango, woody species diversity was highest in mixed AF due to higher species richness. This result corroborate earlier result from a humid tropical ecosystem in south east Nigeria [19]

3.1.2 Carbon stock

Generally, the amount of carbon stored varies between different agroforestry systems [27]. The mean above and below ground biomass values in this study were within values reported by [2] in "indigenous AF systems in Southeastern Rift-Valley Landscapes, Ethiopia". "The mean AGB ranged from 81.1 to 255.9 t ha^{-1} and for BGB from 26.9 to 72.2 t ha^{-1} . The highest C stock was found in Coffee-Fruit tree-Enset based ($233.3 \pm 81.0 \text{ t ha}^{-1}$). But however, our result is higher than the carbon stock obtained in West African Sahel region and in Kenya", respectively [16,28]. "The level of variation observed in biomass among different AF systems might be attributed to the types of trees and shrubs in AF, environmental conditions, soil chemical characteristics, extent of land degradation, and age of the AF system" [1]. Very low land degradation, good environmental conditions, and longer-aged AF systems probably accounted for high biomass production and storage. The AFS reported in this study represent stable AF with less degradation, thus the appreciable biomass recorded in this study.

"To combat rising CO_2 in the atmosphere, carbon sequestration becomes very beneficial when

stored in woody portion of plant species in AFS. Another advantage of having these perennial systems is that C sequestration does not have to end after harvesting the wood component because the stems or branches can also store carbon if processed in any form of long lasting products” [29].

Globally soils in AF systems help in sequestering atmospheric CO₂ [30]. The SOC stocks recorded in the top and sub-soils are considerably higher than the biomass C stocks of AF systems. These results corroborated with the findings of [31], who reported SOC (0–60 cm) from 77–135 Mg ha⁻¹. Swamy [32] also reported the SOC stocks from Agrisilviculture of Chhattisgarh, Central India is 27 Mg ha⁻¹ on average for of 0–60 cm soil depth.

The mixed forest and plantain AF had the highest SOC at both top and sub-soils when compared with Oil palm and bush mango AF. The mixed AF had more herbaceous species and appears to be less disturbed relative to other AFS. The Cultivation of land using tillage practice causes soil disturbance which might cause aggregate breakdown thus releasing previously held SOC [33].

Regarding the biomass C of individual AFS, mixed forestry recorded highest values when compared with other AFS. Luedeling, and Neufeldt [8] did an extensive reviews of biomass C stocks for West African Sahel countries. They discovered that extremely arid and humid regions of Guinea had lower C values from 22.2 to 70.8 t ha⁻¹ than values reported in this study. A global biomass C review on AFS by [1,34] showed that biomass C ranges from 12 to 228 t ha⁻¹. Therefore one can conclude that the biomass C stocks of the studied AF systems fall within the global value range for AF systems and fall within the range of tropical forests and savanna in Brazil and India [35,21].

Carbon sequestration and biodiversity conservation are the most fundamental global environmental challenges, particularly in drylands. Sustainable biodiversity conservation and carbon sequestration are proven means to mitigate current global warming. In the AF systems examined in this study, there was a negative but significant correlation between species richness and diameter at breast height, and positive correlation with total Biomass carbon and total agroforestry carbon on the other hand. These results are consistent with the positive correlation reported by [36]. “We report

that SOC stock increases with increasing species richness and Shannon diversity index in all four AF system. Naturally, ecosystems with high tree diversity sequester more carbon in the soil than those with lower tree diversity” [37].

4. CONCLUSIONS

The agroforestry systems studied include oil palm AF, bush mango AF, mixed AF and plantain AF systems. These agroforestry systems were randomly selected in the area as they were predominant among the indigenous people. Dominant plant species include woody trees, shrubs and herbaceous species.

Results obtained showed that all the agroforestry systems stored carbon. The highest amount of biomass carbon was recorded in mixed AF followed by oil palm AF. However, the highest carbon stored was in the soil component and the litter. This calls for proper soil management for sustainable carbon sequestration.

We noted that there is relationship between SOC stock and species diversity (i.e. species richness and Shannon diversity). The ecosystem C stock of these AF systems compared favorably well with those obtained from other tropical forests and other AF systems in West African countries and India.

ACKNOWLEDGEMENTS

The authors acknowledge the financial support provided by Nigeria Tertiary Education Fund (Tetfund) through Institution Based Research (IBR) grant. Many thanks are also extended to farmers in the study areas where field surveys were undertaken, who shared their precious experience, thoughts and concerns. We are also grateful to the staff of State Ministry of Forestry for their expert advice

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Albrecht A, Kandji ST. Carbon sequestration in tropical agroforestry systems. *Agric. Ecosyst. Environ.* 2003;99:15–27.
2. Tesfay HM, Negash M, Godbold DL, Hager H. Assessing carbon pools of three

- indigenous agroforestry systems in the southeastern rift-valley Landscapes, Ethiopia. Sustainability. 2022;14:4716.
3. Labata MM, Aranico EC, Tabaranza ACE, Patricio JHP, Amparado RFJr. Carbon stock assessment of three selected agroforestry systems in Bukidnon, Philippines. *Advances in Environmental Sciences*. 2012;4(1):5-11.
 4. Lorenz K, Lal R. Soil organic carbon sequestration in agroforestry systems. A review. *Agron. Sustain. Dev.* 2014;34:443–454.
 5. Nair PKR, Kumar BM, Nair VD. Agroforestry as a strategy for carbon sequestration. *J. Plant Nutr. Soil Sci.* 2009;172:10–23.
 6. Tesfay HM, Negash M, Godbold DL, Hager H. Assessing carbon pools of three indigenous agroforestry systems in the Southeastern Rift-Valley Landscapes, Ethiopia. *Sustainability*. 2022 Apr 14;14(8):4716.
 7. Montagnini F, Nair PKR. Carbon sequestration: An underexploited environmental benefit of agroforestry systems. *Agroforestry Systems*. 2004;61-62:281-295.
 8. Luedeling E, Neufeldt H. Carbon sequestration potential of parkland agroforestry in the Sahel. *Clim. Chang.* 2012;115:443–461.
 9. Siyum GE, Tassew T. The use of homegarden agroforestry systems for climate change mitigation in lowlands of southern tigray, Northern Ethiopia. *Asian Soil Research Journal*. 2019;4(1):1-10.
 10. Nair PKR, Nair VD. Carbon storage in North American agroforestry systems. pp. 333–346. In: Kimble J., Heath L.S., Birdsey R.A, Lal R. (eds) *The Potential of U.S. Forest Soils to Sequester Carbon and Mitigate the Greenhouse Effect*. CRC Press, Boca Raton, FL, USA; 2003.
 11. Nair PKR, Vimala D. Nair B, Mohan Kumar, Julia M. Showalter. Carbon sequestration in agroforestry systems, Editor(s): Donald L. Sparks. *Advances in Agronomy*, Academic Press. 2010;108: 237-307.
 12. Basuki TM, Leksono B, Baral H, Andini S, Wahyuni NS, Artati Y, Choi E, Shin S, Kim R, Yang A-R, et al. Allometric equations for the biomass estimation of *Calophyllum inophyllum* L. in Java, Indonesia. *Forests*. 2022;13:10571.
 13. Lasco RD, Group C, Agricult I, Jane R, Oscar D. Agroforestry systems: Helping smallholders adapt to climate risks while mitigating climate Agroforestry systems: Helping smallholders adapt to climate risks. *Wiley Interdiscip Rev Clim Chang.* 2014;5(6):825–33.
 14. Blake GR, Hartge KH. Bulk density. In: Klute, Ed., *Methods of soil analysis, Part I physical and mineralogical methods*, 2nd Edition, ASA-SSSA, Madison. 1986;363-375.
 15. MacDicken K. A guide to monitoring carbon storage in forestry and agroforestry projects. *Forest carbon Monitoring Program*. Winrock Int Inst Agric Dev. 1997;87.
 16. Takimoto A, Nair PKR, Nair VD. Carbon stock and sequestration potential of traditional and improved agroforestry systems in the West African Sahel. *Agric Ecosyst Environ.* 2008;125(1–4):159–66.
 17. Khan AB. Assessment of carbon storage in Pondicherry mangroves, Pondicherry, India. *Acta Biologica Malaysiana*. 2013; 2:95-99.
 18. Sundarapandian SM, Amritha S, Gowsalya L, Kayathri P, Thamizharasi M. Estimation of biomass and carbon stock of woody plants in different land-uses. *Forest Res.* 2013;3:115.
 19. Nwoko CO, Anuna SC, Anyanwu JC. Assessment of carbon sequestration potential of a disturbed humid tropical ecosystem, Southeast Nigeria. W. Leal Filho et al. (eds.), *Climate Change, Hazards and Adaptation Options, Climate Change Management*, Springer Nature Switzerland. 2020;213-224.
 20. Brown S (1997) Estimating biomass and biomass change of tropical forests: A carbon pools of three indigenous agroforestry systems in the comparisons. *Forest Sci.* 2020;45:573-593.
 21. Pragasan AL. Tree carbon stock assessment from the tropical forests of bodamalai hills located In India. *J Earth Sci Clim Change*. 2015;6:314.
 22. Nelson BW, Mequita R, Pereira JLG, de Souza SGA, Batista GT, et al. Assessing tree and stand biomass: A review with examples and critical comparisons. *Forest Sci.* 1999;45:573-593.
 23. Pearson T, Walker S, Brown S. *Sourcebook for land-use, landuse change and forestry projects*. Winrock International

- and the Biocarbon fund of the World Bank. Arlington, USA. 2005;19-35.
24. Lasco RD, MacDicken KG, Pulhin FB, Guillermo IQ, Sales RF, Cruz RVO. Carbon stock assessment of a selectively logged dipterocarp forest and wood processing mill in the Philippines. *Journal of Tropical Science*. 2006;18(4):166-172.
 25. Olayode OO, Saka O Bada, Labode Popoola. Carbon stock in teak stands of selected forest reserves in Southwestern Nigeria. *Environment and Natural Resources Research*. 2015;5(3): 109-115.
 26. Negash M, Yirdaw E, Luukkanen O. Potential of indigenous multistrata agroforests for maintaining native floristic diversity in the south-eastern Rift Valley escarpment, Ethiopia. *Agroforest Syst*. 2012;85(1):9–28.
 27. Luedeling E, Sileshi G, Beedy T, Dietz J. Carbon sequestration potential of agroforestry systems in Africa. 2011;(C):61–83.
 28. Glenday J. Carbon storage and emissions offset potential in an African dry forest, the arabuko-sokoke forest, Kenya. *Environ Monit Assess*. 2008;142(1–3):85–95.
 29. Roshetko JM, Lasco RD, Delos Angeles MS. Smallholder agroforestry systems for carbon storage. *Mitig Adapt Strateg Glob Chang*. 2007;12(2):219–42.
 30. Xie TT, Su PX, An LZ, Shi R, Zhou ZJ. Carbon stocks and biomass production of three different agroforestry systems in the temperate desert region of northwestern China. *Agroforestry Syst*; 2016.
 31. Manaye Ashenaf , Berihu Tesfamariam, Musse Tesfaye, Adefres Worku, Yirga Guf. Tree diversity and carbon stocks in agroforestry systems in northern Ethiopia. Manaye et al. *Carbon Balance Manage*. 2021;16:14.
 32. Swamy SL, Puri S. Biomass production and C-sequestration of *Gmelina arborea* in plantation and agroforestry system in India. *Agroforestry Syst*; 2005.
 33. Dixon RK. Agroforestry systems: Sources of sinks of greenhouse gases? *Agrofor. Syst*. 1995;31:99–116.
 34. Six J, Conant RT, Paul EA, Paustian K. Stabilization mechanisms of soil organic matter: implications for C-saturation of soils. *Plant Soil*. 2002;241:155–76.
 35. Silva LCR, Hoffmann WA, Rossatto DR, Haridasan M, Franco AC, Horwath WR. Can savannas become forests? A coupled analysis of nutrient stocks and fire thresholds in central Brazil. *Plant Soil*. 2013;373:829–842.
 36. Negash M. The indigenous agroforestry systems of the south-eastern Rift Valley escarpment, Ethiopia: Their biodiversity, carbon stocks, and litterfall. *Tropical Forestry Reports*. 2013;44: 75.
 37. Akinremi LRO. Physical properties of earthworm casts and surface soil as influenced by management. *Soil Sci*. 1983;135(2):114–22.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/113905>