



Revolutionizing Maize (*Zea mays* L.) Growth: Evaluating the Impact of Tokyo 8 Microorganism-based Biofertilizer in Southern Côte d'Ivoire Farmers' Fields

Kanga N'guessan Martial ^{a*},
Yao Kouadio Jacques Edouard ^b, Kouamé Konan Didier ^c,
Blafou Adjoua Grace Vanessa ^c, Motchié Fato Edouard ^d
and Kassi Koffi Fernand Jean-Martial ^c

^a University of Man, UFR Agronomic, Forestry and Environmental Engineering (IAFE), Côte d'Ivoire.

^b Agricultural Production Improvement Laboratory, Faculty of Agroforestry, Jean Lorougnon Guédé University, Daloa, Côte d'Ivoire.

^c Plant Physiology and Pathology Teaching and Research Unit of the Biotechnology, Agriculture and Biological Resources Development Laboratory of the Biosciences UFR of the Université Félix Houphouët-Bogny, Côte d'Ivoire.

^d Science and Technology Department, Life and Earth Sciences Section, Ecole Normale Supérieure, Abidjan, Côte d'Ivoire.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2024/v36i54528

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/113340>

Original Research Article

Received: 24/01/2024
Accepted: 29/02/2024
Published: 27/03/2024

*Corresponding author: E-mail: nguessmarti@yahoo.fr;

ABSTRACT

The exclusive use of mineral fertilisers for production contributes to the destruction of certain microflora and microfauna in the soil, as well as reducing the organic matter content, which in turn reduces soil fertility. With the aim of finding an alternative to crop fertilisation on the farm, a micro-organism-based biofertiliser (Tokyo 8) was introduced and tested on maize crops. Three types of Tokyo 8 doses were tested: T2: Tokyo 8 diluted 50 times, T3: Tokyo 8 diluted 75 times and T4: Tokyo 8 diluted 100 times. In addition, two controls were tested, namely T0: control without fertiliser and T1: control with NPK 12-22-22 fertiliser. The trials were set up in a Fisher block design with three replicates. The results show that, like NPK, biofertiliser improves maize growth parameters and yield components. It provides the mineral elements required for maize growth and development. However, to be used effectively, Tokyo 8 biofertiliser needs to be diluted sensibly. A dilution of 75% seems to be the ideal for optimum use in maize crops.

Keywords: Biofertilizer (Tokyo 8); micro-organisms; maize (*Zea mays*); Ivory Coast, maize crops; yield components; grain; pests.

1. INTRODUCTION

Maize, also known by its scientific name *Zea mays* L., is a tropical plant belonging to the Poaceae family, with the world's largest area of cultivation [1]. Maize plays an important role in maintaining food security. Maize grain is eaten in a variety of ways: cooked, grilled or in soup. It is also used to make biodegradable plastics, biofuels and even alcohol [2].

Maize is the world's most widely grown crop, 2023-2024 is estimated at 142.3 Mt (-6% on the previous estimate), compared with 151.5 Mt in 2022-2023. [3]. Among cereals, it ranks third and first respectively ahead of wheat and rice in terms of cultivated area and production. Maize and oilseed grains are the most important source of food for humans. In many developing countries, it accounts for the bulk of people's diets [4].

In Côte d'Ivoire, maize is the second most widely grown cereal after rice [5]. Annual national maize production is estimated at around 600,000 tonnes, 60% of which comes from the savannah region, with a yield of 1.9 tonnes per hectare from a total planted area of 350,000 ha [6]. In our country, maize is essential for food security [7], and is also used in animal feed for poultry, pigs and cattle, and as a raw material in certain industries (brewing, soap production and oil production) [8].

However, maize yields in Côte d'Ivoire remain low and local production covers only 30% of consumption needs [9]. This low yield is due to a combination of factors [10], including, along with disease and pests, a decline in soil fertility [11].

This is mainly due to intensive cultivation and soil erosion. To overcome this major constraint to maize cultivation, farmers use mineral and organic fertilisers [12,13,14,15], but above all chemical fertilisers, which are known for their ability to improve crop yields. Unfortunately, chemical fertilisers are expensive and harmful to the environment and human well-being [16]. Given the decline in soil fertility, the rise in the price of chemical fertilisers on the market and their harmful effects on the environment and human health, it is vital to look for alternatives that will enable sustainable agriculture. It is against this backdrop that the effect of a biofertiliser (Tokyo 8) has been assessed with a view to sustainable soil restoration.

The aim of this study is to contribute to sustainable maize production on the farm. More specifically, the aim is to assess the effect of the biofertiliser (Tokyo 8) on maize growth and yield.

1.1 Study Area

The field trial was carried out at the Université Félix HOUPHOUËT-BOIGNY (UFHB) in Cocody (Abidjan) on the experimental plot of the Unité Pédagogique et de Recherche de Physiologie et Pathologie Végétales (UPR PPV). The plot is located in Abidjan, a city in the south of Côte d'Ivoire with geographical coordinates between latitudes 4°10' and 5°30' North and longitudes 3°50' and 4°10' West.

The relief varies between 0 and 300 m from the Gulf of Guinea, between latitudes 4°17'N and 7°N and longitudes 2°40'W and 8°40'W. It is made up of Precambrian bedrock with a sedimentary cover of Upper Cretaceous-Quaternary age in




Fig. 1. Map of the study area

Table 1. Soil granulometric and chemical composition

Granulometric compositions									
Proportion of clay (%)			Proportion of silt (%)				Proportion of sand (%)		
11,59			6,51				80,98		
Proportions of minerals (%)									Ph
C	N	P	Mg	K	Ca	Fe	O		
12,50	8,3	5	0,11	4	0,06	1,46	62,11	4,73	

the south-eastern part of the geological level. The soils found there are mainly ferrallitic and ferruginous, with some brown eutrophic tropical soils. By virtue of its geographical location, the Université Félix HOUPHOUËT-BOIGNY site belongs to a humid tropical climate zone characterised by a highly contrasting rainfall and temperature regime (Fig. 1).

Granulometric and chemical composition of the soil is shown in Table 1.

2. MATERIALS AND METHODS

2.1 Materials

The experiment involved the LG501 maize hybrid. It has a development cycle of 90 days and a yield of 7 t/ha. It originates from Limagrain, a group of semi-processing cooperatives.

2.1.1 Fertilisers used

The synthetic fertiliser NKP (12-22-22) usually applied in the field for market gardening, vegetables and maize was used as a reference control.

Biofertilisant Tokyo 8 is a formulation of micro-organisms, enzymes and trace elements and nutrients (Table 2). It was tested at 3 doses obtained by dilution to 50, 75 and 100%. It is already being used in Ghana and Liberia, and is currently being tested in Côte d'Ivoire and Guinea. As Tokyo 8 is a biofertiliser, it has less harmful effects on human health. However, it must be applied in accordance with plant protection product safety measures.

2.1.2 Technical equipment

The technical equipment used in this study consisted mainly of tools and instruments

required for growing, maintaining and observing plants in the field and in the laboratory. Items such as machetes, a decameter, ropes, stakes and a backpack sprayer were used to set up and care for the plants. Observations and measurements were made using a variety of tools, including a tape measure, a calliper, a 30 kg weighing scale of a specific brand, transparent polyethylene plastic bags, permanent markers and a digital camera.

2.2 Methods

2.2.1 Experimental design

The experimental design was a Fisher randomised complete block design. The factor studied was the dose of biofertiliser with three (3) treatments and three (3) replications. The experimental plot was 14 m long and 11 m wide, i.e. an area of 154 m². The distance between the blocks was 2 m, and that between the individual plots 1 m (Fig. 2). Each individual plot had an area of 6 m².

2.2.2 Setting up and running the trial

The previous crop in the experimental plot was fallow. As a result, the plot was prepared before the trial was set up. This involved clearing the land and ploughing to a depth of 15-20cm. The experimental layout was then laid out. Each micro-plot consisted of 4 rows, each bearing 52 bunches (Should be minimum 6 rows, more rows are better in fertilizer experiment, because border rows effect a lot to experiment error. Another attention is how to maintain uniform plant density from emergence to harvest, otherwise effects a lot to error of experiment). Sowing was carried out after a useful watering of 45 l of water per micro plot. Sowing was carried out manually, with two seeds per plot. Fertiliser was applied 15 and 35 days after sowing. Weeding was carried out before each application. A total of three doses of Tokyo 8 biofertiliser were tested in comparison with NPK (12- 22- 22) at a dose of 150 kg/ha and an untreated control. The biofertiliser doses were obtained by diluting the recommended dose of 1 ml/m². The dilution factors were 50, 75 and 100%. Thus, 6 ml of biofertiliser were placed in a beaker and then water was added until a volume of 900, 1350 and 1800 ml was reached for the 50, 75 and 100% dilutions respectively. This quantity of spray mixture was then put into a sprayer and applied to the microplots concerned. The biofertiliser was applied by sprinkling at the foot of the plants, while the NPK fertiliser was

applied by spreading at the foot of the maize plants. The biofertiliser was applied to the plants by foliar spraying. Phytosanitary treatments based on cypercal (50ml/7.5l of water) were carried out to reduce attacks by the armyworm.

2.2.3 Data collection

Observations focused on measurements of agromorphological parameters and yield components of maize plants. The quantitative variables were measured on twenty (20) marked plants selected at random (Should be consecutively) from each microplot. Observations were made every week from 30 DAS until flowering began.

2.2.4 Agro-morphological parameters

Measurements were taken on 5 plants per row selected at random (consecutively is better) on each elementary plot and in each block, i.e. a total of 20 plants per microplot. Some characteristics were selected from among the maize descriptors. Plant height was measured with a tape from the crown to the stem apex 50 cm above the ground. This was done every week until flowering. The number of leaves was determined by counting them every week. Stem diameter was also measured using a caliper at the neck of the plants.

These data were used to calculate the vigour index (IV) according to the equation in [17]:

$$IV = \text{Log} [(H * C2) / 4\pi]$$

Where π is the constant 3.14; IV : Vigour index ; H : Plant height ; C : Plant circumference.

2.2.5 Yield components

Flowering: As soon as the first male flowers (panicle) and female flowers (ear) appeared, i.e. 52 days after sowing, the number of flowers that appeared was counted on all the plants. They were assessed by counting the flowers every two days, which made it possible to calculate the number of days at 50% male flowering and female flowering.

Fresh yield: At harvest, the fresh yield was assessed. To do this, the two central rows of each microplot, bearing a total of 15 plants, were harvested. The number of ears and the mass of each ear were determined. The number of ears per m² was calculated by dividing the number of ears by the area harvested. Similarly, the

Table 2. Tokyo 8's constituents

Micro-organismes	Enzymes	Mineral
proteobacteria	lipases	Magnesium
chlorophyceae	proteases	potassium
actinobacteria	cellulase	phosphorus
firmicutes		calcium
fungi		iron
		chlorine
		manganese
		Nickel copper
		zinc
		molybdene




Fig. 2. Experimental set-up

T0: control without fertiliser ; T1 : control with NPK 12-22-22 fertiliser ; T2 : Tokyo 8 diluted 50 times ; T3 : Tokyo 8 diluted 75 times ; T4 : Tokyo 8 diluted 100 times

average ear mass was calculated by dividing the total ear mass by the number of ears. The yield induced by each treatment was calculated by dividing the total mass of ears by the area harvested.

The method for measuring dry grain yields follows the same principle as for fresh grain yields. Except for the evaluation, the ears were dried for 4 weeks in the sun.

Data analysis and processing: All the data collected was entered using Excel. An analysis of variance (ANOVA) was performed using Statistica software version 7.1. The Newman Keul test was used for the comparison of means to establish the significance of differences between treatments at the 5% threshold.

3. RESULTS

3.1 Effect of Biostimulant Doses on Maize Agro-Morphological Parameters

3.1.1 Changes in plant height

Changes in average plant height of treatment after sowing are shown in Fig. 3, which shows that average plant height increased sharply over time, regardless of treatment. The average height ranged from 18.97 to 23.13 cm at week 1, and by week 4 had risen to values ranging from 86.99 to 111.17 cm. However, the average plant height in treatments T1 and T3 was higher throughout the evaluation period.

The average height of the maize plants obtained at the end of the observations varied significantly

from one treatment to another. Treatment T1 (NPK fertiliser) produced the highest heights, with an average height of 58.77 cm. On the other hand, treatments T2 and T4 (Tokyo 8 diluted 50 and 100%) showed low heights with mean heights of 45.23 and 46.19 cm respectively. Treatments T0 (no fertiliser) and T3 (biofertiliser diluted 75%) gave intermediate heights with mean heights of 52.66 and 54.64 cm respectively (Fig. 4).

3.1.2 Changes in plant diameter

At the end of the observations, a significant difference was recorded between the stem diameters obtained on the treatments. Treatment T1 (NPK fertiliser) produced the largest diameter with an average of 17.66 mm. The smallest diameters were observed in treatments T2, with

averages of 14.23 (Fig. 5). Treatments T0, T3 and T4 produced statistically identical intermediate diameters.

3.1.3 Changes in maize plant leaf

Fig. 6 shows the evolution of full maize plant leaves. This Fig. 6 shows that full leaves increased slightly with time, whatever the treatment. The greatest number of leaves was recorded in treatments T0, T1 and T3 over the entire observation period. T2 and T4 induced the lowest number of leaves in weeks 1 and 3. At the end of the observations, data analysis showed no significant difference ($p > 0.05$) between treatments. The number of leaves emitted was 11.25, 11.01, 10.98, 10.38 and 10.31 respectively for treatments T0, T1, T2, T3 and T4 (Fig. 7).




Fig. 3. Height versus time curve




Fig. 4. Plant height as a function of treatment

T0: control without fertiliser, T1: control with NPK 12-22-22 fertiliser, T2: Tokyo 8 diluted to 50%, T3: Tokyo 8 diluted to 75%. T4: Tokyo 8 diluted to 100 ; Error bars with the same letters are statistically identical according to the N-Keuls test with a threshold of $\alpha=0.05$




Fig. 5. Effect of treatments on plant diameter

T0: control without fertiliser, T1: control with NPK 12-22-22 fertiliser, T2: Tokyo 8 diluted to 50%, T3: Tokyo 8 diluted to 75%. T4: Tokyo 8 diluted to 100 ; Error bars with the same letters are statistically identical according to the N-Keuls test with a threshold of $\alpha=0.05$





Fig. 6. Evolution of the number of leaves as a function of time

Fig. 7. Effect of treatments on the number of leaves emitted

T0: control without fertiliser, T1: control with NPK 12-22-22 fertiliser, T2: Tokyo 8 diluted to 50%, T3: Tokyo 8 diluted to 75%. T4: Tokyo 8 diluted to 100 ; Error bars with the same letters are statistically identical according to the N-Keuls test with a threshold of $\alpha=0.05$




Fig. 8. Effect of treatments on vigour index





Fig. 9. Male flowering rate as a function of time

Fig. 10. Male flowering rate as a function of treatment

T0: control without fertiliser, T1: control with NPK 12-22-22 fertiliser, T2: Tokyo 8 diluted to 50%, T3: Tokyo 8 diluted to 75%. T4: Tokyo 8 diluted to 100 ; J1,2,3 ;4,5 : Day 1,2,3 ;4,5 ; Error bars with the same letters are statistically identical according to the N-Keuls test with a threshold of $\alpha=0.05$

3.1.4 Vigour index

Analysis of variance of the data revealed significant differences between treatments in terms of vigour index. The most vigorous plants were observed in treatment T1, with an average index of 3.05. On the other hand, the least vigorous plants were observed in treatments T2 and T4, with averages of 2.72 and 2.77 respectively. Treatments T3 and T0 recorded intermediate vigour indices of 2.91 and 2.90 respectively (Fig. 8). T0 should be least vigorous? Why in intermediate vigorous ?

3.2 Effect of Treatment on Yield Components

3.2.1 Evolution of flowering of plants according to treatments

The first male and female flowers were observed on days 52 and 56 respectively after sowing. The rate of male flowering increased slightly until day 54 and then sharply until the end of the observations. Treatment T1 induced the highest rates of male flowering throughout the observation period. It was followed by T3 and then T0. Treatments T2 and T4 induced the lowest flowering rates. At the end of the observations, the male flowering rate was 78, 62, 57, 43 and 41% for treatments T1, T3, T0, T4 and T2 respectively. However, analysis of variance showed no significant difference ($p > 0.05$) between treatments (Figs. 9 and 10).




Fig. 11. Female flowering rate as a function of time

T0: control without fertiliser, T1: control with NPK 12-22-22 fertiliser, T2: Tokyo 8 diluted to 50%, T3: Tokyo 8 diluted to 75%. T4: Tokyo 8 diluted to 100 ; J1,2,3 ;4,5 : Day 1,2,3 ;4,5 ; Error bars with the same letters are statistically identical according to the N-Keuls test with a threshold of $\alpha=0.05$

In the case of female flowers, the rate of flowering increased sharply over time. Treatment T1 recorded the highest flowering rate over the entire observation period. It was followed by treatment T3 and then the control T0. The female flowering rates observed in treatments T4 and T2 were below those of the control.

At the end of the observations, the female flowering rate was 23, 21, 17.14 and 9% respectively on treatments T1, T3, T0, T4 and T2. However, analysis of variance showed no significant difference ($p > 0.05$) between treatments (Figs. 11 and 12).

3.2.2 Number of fresh ears as a function of treatment

The number of ears per plant varied significantly from one treatment to another. Treatment with T1 induced the highest number of ears with an average of 24.67. The lowest number of ears was recorded in the untreated control with an average of 16.3. Treatments T3 and T4 induced ear numbers very close to those of T1. Treatment T2 produced an ear number very close to that of T0 (Fig. 13).

3.2.3 Average fresh ear weight by treatment

The average mass of the ear was 0.24, 0.25, 0.21, 0.20 and 0.20 kg respectively for treatments T1, T2, T3, T4 and T0. However, analysis of the data revealed no significant differences. (Fig. 14)




Fig. 12. Female flowering rate as a function of treatment




Fig. 13. Number of spikes as a function of treatment

T0: control without fertiliser, T1: control with NPK 12-22-22 fertiliser, T2: Tokyo 8 diluted to 50%, T3: Tokyo 8 diluted to 75%. T4: Tokyo 8 diluted to 100 ; Error bars with the same letters are statistically identical according to the N-Keuls test with a threshold of $\alpha=0.05$




Fig. 14. Average ear weight as a function of treatment

T0: control without fertiliser, T1: control with NPK 12-22-22 fertiliser, T2: Tokyo 8 diluted to 50%, T3: Tokyo 8 diluted to 75%. T4: Tokyo 8 diluted to 100 ; Error bars with the same letters are statistically identical according to the N-Keuls test with a threshold of $\alpha=0.05$

The Fig. 15 means soil fertility is fertile. So average fresh ear weight is no different among T0, T1, T2, T3 and T4?

3.2.4 Yield in kg/ha as a function of fresh treatment

The fresh yields obtained from the different treatments are shown in Fig. 16. Analysis of variance revealed a significant difference ($p < 0.05$) between treatments. Treatment with T1 produced the highest yield, with an average of 18888.89 kg/ha. By contrast, the lowest yield was recorded for treatment T2 with 7933.33

kg/ha. Treatments T3, T4 and T0 recorded intermediate yields with 14888.89, 14833.33 and 14000 kg/ha respectively. (Fig. 15)

3.2.5 Dry yield in tone/ha

The mass grain yield (tone/ha) was a function of treatment (Fig. 16). Analysis of variance revealed a significant difference ($p < 0.05$) between treatments. Treatments T1 and T3 gave the highest yields, with an average of 2.98 and 2.63 tone/ha respectively. By contrast, the lowest yield was recorded in treatment T2 with 1.8 tone/ha. Treatments T0 and T4 recorded 2.04 and 2.17 tone/ha respectively.




Fig. 15. Fresh yield as a function of treatment

T0: control without fertiliser, T1: control with NPK 12-22-22 fertiliser, T2: Tokyo 8 diluted to 50%, T3: Tokyo 8 diluted to 75%. T4: Tokyo 8 diluted to 100 ; Error bars with the same letters are statistically identical according to the N-Keuls test with a threshold of $\alpha=0.05$




Fig. 16. Dry yield as a function of treatment

T0: control without fertiliser, T1: control with NPK 12-22-22 fertiliser, T2: Tokyo 8 diluted to 50%, T3: Tokyo 8 diluted to 75%. T4: Tokyo 8 diluted to 100 ; Error bars with the same letters are statistically identical according to the N-Keuls test with a threshold of $\alpha=0.05$

In grain yield of maize, shelling percentage is very important criteria indicating stress tolerance at kernel setting period for final harvest yield. If you have this kind of data, please supplement them.

3.2.6 Correlation between growth parameters and yield components

Principal component analysis (PCA) was carried out on the data collected on the growth

parameters and yield components of maize after application of the different doses of Tokyo 8 biofertiliser. These were mean height, diameter, number of leaves, vigour index, male and female flowering rate, number of ears, mean ear weight and fresh yield (why not included dry grain yield ?). The results showed that axes 1 and 2, which express 91.1% of the inertia, were sufficient to characterise the variables studied. All the variables were strongly positively correlated with axis 1, which accounted for more




Fig. 17. Correlation circle according to PCA of treatments

Ha: height; *Dm*: diameter; *iv*: vigour index, *Ff*: female flowers; *Fm*: male flowers ; *Rend*: yield; *N'épi*: number of ears




Fig. 18. Dispersion of treatments

T0: control without fertiliser, *T1*: control with NPK 12-22-22 fertiliser, *T2*: Tokyo 8 diluted to 50%,
T3: Tokyo 8 diluted to 75%. *T4*: Tokyo 8 diluted to 100

than half (60.93%) of the information, with the exception of yield and number of ears. The latter were strongly positively and strongly negatively correlated respectively with axis 2, which accounts for 30.74%. Height, diameter, vigour index and the percentage of female and male flowers were strongly correlated with each other. The dispersion of the treatments revealed four homogeneous groups. Treatments T1 and T3 produced the best maize growth and yield. Treatment T2 recorded the lowest values and treatments T0 and T4 recorded intermediate values. (Figs. 17 and 18).

4. DISCUSSION

Chemical fertiliser produced the best growth and yield in maize. This is because the chemical fertiliser provided the plants with the mineral elements required for optimum growth and development. It is rich in nitrogen, phosphorus and potassium, the levels of which correspond to the needs of the maize crop. This supply of mineral elements that can be directly assimilated by the plant in sufficient quantities is responsible for the rapid growth and good development observed. Our results are in line with those of [18], who showed that plant nutrition and nitrogen fertilisation are essential factors in determining the yield and quality of agricultural production in tropical environments. According to the work of [19] and [20], the nitrogen (N), phosphorus (P) and potassium (K) contained in chemical fertilisers are involved throughout the plant development cycle. As a result, their absence can lead to unsatisfactory development and yields [18] (But the result of fresh ear weight is no different in this experiment? not yet explained?)

The effects of the various fertilisers on growth parameters were remarkable. The Tokyo 8 biofertiliser improved the growth of the maize plants compared with the untreated control. This result could be explained by the composition of this biofertiliser. It is composed of mineral elements such as nitrogen (NPK, 12-22-22+2SO₃ + MgO + 5CaO), which would have played an active role in the vegetative growth of the maize. Our results are in agreement with those of [20] who showed that nitrogen is a determining factor in plant growth and the determination of plant yield. Also, [21] showed that biofertilisers have the necessary mineral elements for the growth and development of field crops such as maize.

Tokyo 8 biofertiliser is also rich in enzymes and microorganisms. These would have stimulated mechanisms to improve the bioavailability of nutrients in the soil; qualitatively modify soil microbial communities (new equilibrium) and increase soil microbial activity favourable to good plant development. Our results corroborate those of [22], who showed that biofertilisers also improve soil structure by stimulating microbial activities and nitrogen uptake by plants by modifying their morphological and physiological characteristics.

As with the growth parameters, the Tokyo 8 biofertiliser applied to the crop led to an improvement in the components of maize development and yield. This result could be justified on the one hand by the good growth induced by the biofertiliser. Our observation is in line with that of [23], who highlighted the improvement in maize yield by four types of organic fertiliser through the good growth induced by them. Thus, good plant growth is a prerequisite for good crop yields. Tokyo 8 biofertiliser also contains micro-organisms. According to the work of [24], micro-organisms release gibberellin into the environment, which is involved in bud initiation, stem growth, flowering, the breaking of seed dormancy and fruit growth. All these results prove that Tokyo 8 biofertiliser is an organic fertiliser that is very rich in the mineral elements essential for healthy plant development.

The biofertiliser diluted to 75% gave fresh or dry grain yields similar to those of the chemical NPK fertiliser (12 22 22). The 50% dilution slowed down maize plant growth and significantly reduced yields. As for the 100% dilution, it induced growth and yield very close to that of the control. These results show that Tokyo 8 biofertiliser needs to be diluted carefully if it is to be effective. A dilution of 75% seems to be the most suitable for improving the effectiveness of Tokyo 8 biofertiliser in maize crops. This result is similar to those obtained by [25] in tomato crops using seaweed-based biofertilisers. These authors showed that these biofertilisers were effective from the 50% dilution, whereas the 25% dilution was ineffective.

The fact that the Tokyo 8 biofertiliser diluted to 75% induced growth and yield close to that of the NPK chemical fertiliser shows that its composition is sufficient to ensure optimal nutrition of maize plants. Our results are in agreement with those of [26] who show that foliar

spraying of a seaweed extract (SE) biofertiliser improved the yield of baby lettuce under low nitrogen conditions. However, unlike chemical fertilisers, which have a negative impact on the environment, plants and human health, biofertilisers have the advantage of improving the environment. According to the work of [27], biofertilisers improve microbial activity or the physico-chemical properties of the soil, enabling organic compounds to be degraded more effectively.

5. CONCLUSION

This study was carried out with the aim of contributing to sustainable maize production on the farm. To this end, the biofertiliser Tokyo 8 was tested in comparison with the chemical fertiliser NPK 12 22 22 and a control. The results obtained showed that, like NPK, biofertiliser improves the growth parameters and yield components of dry and fresh maize. It provides the mineral elements (No data for this remarks ?) required for maize growth and development. However, for Tokyo 8 biofertiliser to be used effectively, it must be diluted carefully. A dilution of 75% would appear to be ideal for optimum use in maize crops. Tokyo 8 biofertiliser diluted to 75% induces vegetative growth and maize yields very close to those of chemical NPK fertiliser. It could be used to enrich the soil.

In view of this study, it would be advisable to :

- Test the Tokyo 8 biofertiliser on other crops grown by farmers.
- Evaluate the effect of the biofertiliser in combination with NPK
- Test the biofertiliser in other localities.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Yara. Crop nutrition. But. In Ivory Coast. Available:<https://www.yara.ci/fertilisation/fertilisation/mais/mais/> Accessed October 7, 2023
2. Hoopen TEM, Maïga A. Corn production and processing. Pro-Agro Collection. 2012;31.
3. France Agricole; 2023. Available:<https://www.lafranceagricole.fr/condition-cultures/article/848469/la-production-mondiale-de-mais-en-hausse-pour-2023-2024#:~:text=En%20effet%2C%20la%20production%20mondiale,5%20Mt%20en%202022%2D2023> Accessed February 25, 2024
4. Guèye MT, Seck D, Wathelet J-P, Lognay G. Control of pests of cereal and legume stocks in Senegal and West Africa: Bibliographical synthesis. Biotechnology. Agronomy.Soc. Environ. 2011;15(1):183-194.
5. Akanvou L, Akanvou R, Toto K. Effects of maize and legume varieties in the fight against *Striga hermontica* in the savannah zone of Côte d'Ivoire. African Agronomy. 2006;18(1):13-21.
6. Yara. Crop nutrition. But. In Ivory Coast; 2020. Available:<https://www.yara.ci/fertilisation/fertilisation/mais/mais/> Accessed June 7, 2023
7. Sangaré A, Koffi E, Akamou F, Fall MA. National report on the state of plant genetic resources for food and agriculture. 2009;65.
8. Boone, Peter, Stathacos, Charles.JD, Wanzie, Rose.L. Sub-regional assessment of the maize value chain. ATP technical report n°1. Bethesda, MD: ATP Project, Abt Associates Inc. 2008;10.
9. ANSD. Agricultural campaign survey report. 2019-2020; 10.
10. Mbow M. The challenges of Senegalese agriculture in the context of climate change. 2017;50.
11. AKanvou HA, N'da L, Kouakou CK. Local management of the varietal diversity of maize (*Zea Mays* L.) violet Tagouana in the North-Central Ivory Coast. Int. J. Biol. Chem. Sci. 2013;7(5) :2058-2068. DOI: <http://dx.doi.org/10.4314/ijbcs.v7i5.23>
12. Zenabou H, Mahamane S, Bismarck NH, Bado BV, Lompo F, André B. Effect of the combination of organo-mineral fertilizers and the cowpea-millet rotation on nitrogen nutrition and millet yields in the Sahel. In: International Journal of Biological and Chemical Sciences. 2014;8(4): 1620-1632.
13. Akanza KP, Sanogo S, N'Da HA. Combined influence of organic and mineral manure on corn nutrition and yield. Impact on the diagnosis of soil deficiencies. Tropicultura. 2016;34(2):208-220.
14. Somda BB, Ouattara B, Serme I, Pouya MB, Lompo F, Taonda SJB & Sedogo PM.

- Determination of optimal doses of organo-mineral fertilizers in microdose in the Sudanosahelian zone of Burkina Faso. In: International Journal of Biological and Chemical Sciences, 2017;11(2):670-683.
15. Badiane A, Goudiaby AOK, Diop B, Ndour NYB. Characteristics of two compost formulas to valorize organic matter in Casamance (Senegal). Journal of Materials and Environmental Sciences. 2019;8. ISSN: 2028-2508 CODEN: JMESCN.
 16. Mbonigaba. Physical, chemical and microbiological characterization of three tropical acidic soils of Rwanda under natural fallows and constraints on their productivity. 2007;402.
 17. Kouassi B. Analysis of the determinants of the choice and adoption of improved rice varieties in the Gagnoa and Korhogo zones. End of cycle dissertation, INPHB. Ivory Coast. Associates Inc. 2019;117:10.
 18. Boulal H, Zaghouane O, Mourid M, Rezgui S. Practical guide to the management of autumn cereals (wheat and barley) in the Maghreb (Algeria. Morocco. Tunisia). 2007;176.
 19. Abdelmadjid H. Field crops elements of general pyrotechnics 1st ED. T1. Wheat. 2013;256.
 20. Ognalaga M, Odjogui PIO, Lekambou JM, Poligui RN. Effect of sugar cane scum, powder and compost based on *Chromolaena odorata* (L.) King R.M. & H.E. Rob on the growth of Guinea sorrel (*Hibiscus sabdariffa* L.). International Journal of Biological and Chemical Sciences. 2015;9:2507-2519.
 21. Muyayabantu GM, Kadiata BD, Nkongolo KK. Response of maize to different organic and inorganic fertilization regimes in monocrop and intercrop systems in a Sub-Saharan Africa Region. Journal of Soil Science and Environmental Management. 2012;3(2):42-48. DOI: 10.5897/JSSEM11.079
 22. Halpern M, Bar-Tal A, Ofek M, Minz D, Muller T, Yermiyahu U. The use of biostimulants to improve nutrient absorption, in: Advances in Agronomy. Elsevier. 2015;141-174.
 23. Siéné LAC, Doumbouya M, Traoré MS, Condé M, N'Guettia TVF, Koné M. Effect of four types of fertilizers on the growth and productivity of two genotypes of maize (*Zea mays* L.) in case of Late sowing in Korhogo in the North-Central Ivory Coast. Int. J. Biol. Chem. Sci. 2020;14(1): 55-68.
 24. Razaquia Khan M, Masroor A, Khan, Minu Singh, Shafia Nasir, M Naeem, Manzer H. Siddiqui, Firoz Mohammad. Gibberellic acid and triacontanol can improve the opium yield and morphine production in opium poppy (*Papaver somniferum* L); 2007. accessed September 3, 2023.
 25. Abidi L, Snoussi SA, Bradea MS. Effect of a biofertilizer on the technological and nutritional quality of two varieties of tomatoes grown in greenhouses. Revue Agrobiologia. 2016;6(1)5.
 26. Di Mola I, Cozzolino E, Ottaiano L, Giordano M, Roupheal Y, Colla G, Mori M. Effect of biostimulants based on plant extracts and algae on the agronomic traits and leaf quality of young lettuces grown in plastic tunnels under four nitrogen fertilization regimes. Agronomy 2019; 9(10):571.
 27. Povero G, Mejia JF, Di Tommaso D, Piaggese A, Warrior P. Systematic approach to discover and characterize natural plant biostimulants. Frontiers in Plant Science. 2016;7:435.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/113340>