



The Function of RNA Interference (RNAi) in Crop Enhancement-Mechanism and Applications: A Review

Harmanpreet Kaur ^a and Nilesh Talekar ^{a*}

^a *Department of Genetics & Plant Breeding, School of Agriculture, Lovely Professional University, Phagwara-144411, Punjab, India.*

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JEAI/2024/v46i62487

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/116796>

Review Article

Received: 02/03/2024

Accepted: 06/05/2024

Published: 08/05/2024

ABSTRACT

RNA interference (RNAi) is a conserved molecular mechanism that plays a critical role in post-transcriptional gene silencing across diverse organisms. This review delves into the role of RNAi in plant functional genomics and its applications in crop improvement, highlighting its mechanistic insights and practical implications. The overview starts with the fundamental finding of the mechanism of RNAi, following its journey from petunias to its ubiquitous presence in a wide range of species. The discovery of several kinds of regulatory non-coding small RNAs, including as phasiRNAs, miRNAs, and siRNAs, has broadened our knowledge of RNAi-mediated gene regulation beyond what is previously thought. These RNA types have an impact on gene expression through complex post-transcriptional and epigenetic mechanisms. RNA interference (RNAi) has become a potent tool for comprehending gene activities in the context of crop improvement. It has shown to be useful in determining the functions of genes for metabolic pathways, stress tolerance, and more. Furthermore, RNAi-based strategies have potential for sustainable agriculture and integrated pest management, supporting international efforts to ensure

*Corresponding author: E-mail: nilesh.23498@lpu.co.in;

food security. The many uses of RNA interference (RNAi) are covered in this article, including improving crop nutrition, changing plant architecture, and increasing shelf life. The challenges and future prospects of RNAi technology are also explored. The worldwide geography of RNA interference research is emphasized, with notable contributions from North America, Europe, and China. In conclusion, RNA interference (RNAi) continues to be a flexible and essential tool in contemporary plant research, providing new pathways for understanding gene activities and enhancing crop characteristics. Its combination with other biotechnological techniques, such as gene editing, might influence how agriculture and sustainable food production develop in the future.

Keywords: *RNA interference; plant functional genomics; Non-coding small RNAs; crop improvement; gene expression; sustainable agriculture; biotechnological techniques.*

1. INTRODUCTION

RNA interference (RNAi) is a natural defense mechanism against double-stranded RNA (dsRNA), targeting viral and cellular mRNAs. The cleavage products are microRNA (miRNA) and small interfering RNA (siRNA), produced by DICER enzyme. In recent years, researchers have shifted towards biotechnology methods for crop development, allowing for easy manipulation of gene expression in crops for desirable features [1]. Numerous desirable traits, such as nutritional fortifications, allergen reduction or allergen substitution, morphological alterations, male sterility alternation, secondary metabolite enhancement, and defense against a variety of biotic and abiotic stresses in different crop plants, have all been modified successfully through the use of RNA interference [2].

RNA interference (RNAi), which was aptly named Science's "Technology of the Year" in 2002 and Fortune Magazine's "Billion Dollar Breakthrough" in 2003, works by suppressing gene expression by breaking down particular messenger RNAs (mRNA) [3].

Plants provide food, shelter, and other necessities for life, as well as useful products like wood and medicine. However, constraints like limited land and changing climate pose a danger to this crucial partnership. Although artificial gene transfer raises difficulties, genetic engineering provides remedies by increasing agricultural yields and nutrition. Let me introduce you to RNAi, a revolutionary technology that silences genes without the need for transgenics, perhaps lowering dangers and winning over people. This breakthrough improves agricultural resilience and illuminates the control of genes, among other areas, and provides access to new technologies. By impacting plant development and stress

tolerance, RNA interference (RNAi) is paving the way for a future in which artificial manipulation strengthens rather than upsets the delicate equilibrium between people and plants [4].

A putative RdRP, SGS2, SDE1, is necessary for PTGS in plants that are started by transgenes that overexpress an endogenous mRNA; however, transgenes intended to produce dsRNA get around this need. Because sRNAs are not cell autonomous, they can travel throughout the plant and provide mobile silencing signals that silence genes across large distances or from one cell to another [5].

Petunia flowers revealed a curious silencing of color genes, sparking the discovery of RNAi in worms. This potent process uses injected double-stranded RNA to dismantle specific genes, even in offspring. The mystery of its function and RNA transport fuelled research across various organisms, boosted by the power of genomic data. Today, RNAi remains a crucial tool for understanding and manipulating gene activity in plants and animals [6].

Advances in RNA-based gene regulation, specifically RNA interference (RNAi), have led to improved crop quality and reduced biosafety issues. RNAi is a gene silencing phenomenon used for gene function assessment, plant metabolic engineering, and developing stress-tolerant and disease-resistant crops [7].

The discovery of regulatory noncoding RNA (ncRNA) led to the genetic revolution and two new areas of study: understanding ncRNA functions and practical applications like RNA interference (RNAi) to manipulate cellular defenses. RNAi and CRISPR, a bacterial virus defense system, have been suggested as potential magic bullets for medicine and agriculture. RNAi is still used experimentally for

disease management in agriculture, reverse genetics, and functional genomics, despite the use of CRISPR offering new promises and challenges. Recent reviews have explored various aspects of RNAi for plant disease management strategies, including a special issue of *Current Opinion in Virology* [8].

Double-stranded RNA (dsRNA) is a widely used gene function analysis tool due to its specificity and effectiveness. It is found in plants and animals and has been used in genome-wide analysis in worms and flies. RNAi techniques have been developed for high-throughput analysis in cultured cells. Recent research has demonstrated the natural functions of RNA interference, including suppression of transposon activity, resistance to viral infection, post-transcriptional and post-translational regulation of gene expression, and epigenetic control of chromatin structure. RNAi is expected to be useful in genetic engineering crop plants, particularly as a knockdown technique [9].

Antisense oligonucleotide effectiveness is drastically decreased by cellular nucleases, which break down these molecules quickly after they are administered. This barrier might be addressed by using synthetic oligonucleotides that change the phosphodiester bond's composition by substituting sulfur for oxygen. We refer to these altered oligonucleotides as phosphonothioates [10].

Agriculture is being transformed by RNAi technology. Think about fruits without seeds, fruits with a longer shelf life, and fruits with more nutrition—all perfectly accessible! Plant genes can be altered by this novel approach, which can improve tolerance to adverse conditions, regulate biomass, and change the color of flowers. As allergies are reduced, it even permits the addition of beneficial nutrients. The primary factor controlling important plant characteristics is siRNA. A large portion of our protein and 65% of our calories come from cereals like wheat and maize. In influencing the future of cereal crops, this paper examines the fascinating real-world uses of RNA interference [11].

Small interfering RNA was found through more research, which prompted computer scientists to become involved in the prediction and synthesis of precise and artificial RNA strands with greater specificity. This process improves the production of superior transgenic plants with higher market values. A greater and superior quality of crops

with high nutritional content, resistance to disease and abiotic and biotic stress, and improved horticultural features, such as the color of the flowers, can be obtained thanks to RNAi technology and the accessible genetic pool [12].

Since RNAi-based gene silencing mechanisms are conserved in both the plant and animal worlds, they have potential uses in agriculture, such as stress tolerance engineering, as well as functional genomics. Furthermore, advancements in the RNAi technique have unlocked its immense potential to address a number of agricultural difficulties [13].

2. HISTORY OF RNAI

Gene function was previously studied using a variety of techniques, including antisense RNA suppression, mutagen or radiation therapy, T-DNA element and transposon insertion, and more. These techniques, however, were laborious and unreliable. As a result, the RNA interference (RNAi) approach was discovered. Plant scientists first employed antisense RNA suppression as a type of RNA silencing (Williams et al., 2004). However, this strategy did not always completely quiet the target gene, which led to more study into alternate gene silencing techniques. By concurrently introducing sense and antisense strands, Fire et al. (1998) initially reported the RNAi approach in the roundworm *Caenorhabditis elegans*, generating the double-stranded RNA (dsRNA) necessary for initiating the RNAi pathway. When compared to employing simply sense or antisense strands alone, this unexpectedly led to a tenfold increase in targeted gene silencing. This groundbreaking discovery encouraged researchers to explore the complex RNAi phenomenon in greater depth, gaining insight into the molecular workings of RNAi technology and how it affects the activity of particular genes. Plant breeders are now able to use RNAi as a tool for crop enhancement and plant protection thanks to this significant breakthrough [10].

Molecular biologists have been working to prevent gene expression at the mRNA level for the last 20 years. Although antisense approaches utilizing DNA or RNA were relatively straightforward ways to investigate gene activity, their efficacy and specificity were limited. Furthermore, it was sometimes difficult to predict the intended effects, and only modest suppression could be achieved. After the

discovery of post-transcriptional gene silencing (PTGS) in plants in 1998, RNAi function in flies, mammals, plants, and fungi was also seen. The lengthy dsRNA molecules start the process. dsRNA is produced by viruses and transposon activity during natural RNAi processes; it may also be injected into cells for research reasons. The complementary opposite strand is referred to as the antisense strand, while the dsRNA strand that matches a particular area of the target mRNA molecule is referred to as the sense strand. After identifying dsRNA, a Dicer complex that resembles RNase III in *D. melanogaster* cleaves it into segments that are approximately 22 nucleotides long. These fragments, referred to as "small interfering RNAs" (siRNAs), continue to exist as short 3' overhanging double-stranded duplexes. The homologous mRNA is degraded and its synthesis is particularly suppressed by the RNA-induced silencing complex (RISC), which uses these duplexes as templates. This kind of RNAi is called PTGS; in some species, other kinds may also have transcriptional or genomic effects. Nonetheless, Fire and Mello made a substantial contribution to this field's growth. Motivated by the previously reported occurrences, they pondered the potential outcomes of introducing both sense and antisense strands into the worms. Surprisingly, this combination of strands was ten times more efficient than either strand alone at reducing the expression of a particular gene. After identifying the uniqueness of the dsRNA-induced effect, Fire and Mello gave the mechanism the moniker RNA interference (RNAi) [14].

3. MECHANISM OF RNAI

Components of RNAi Machinery: Dicer and the RNA-induced silencing complex (RISC) are the two ribonucleases that take part in the RNAi pathway. Dicer converts dsRNA into active small non-coding RNAs and starts the RNAi pathway, while RISC and the RNase H core enzyme Argonaute (AGO) carry out the gene silencing. The N-terminal helicase domain, a PAZ (Piwi/Argonaute/Zwille) domain, dual RNase III domains, and a dsRNA binding domain make up the Dicer family, which is a class 3 RNase III enzyme. These enzymes' main job is to identify the dsRNA precursor from the RNAi pathway and produce short non-coding RNA with a precise length (between 21 and 24 nt). According to the Dicer catalysis concept, the active core of the multidomain dicer enzyme is formed when two RNase III domains dimerize to produce an intramolecular pseudo-dimer. Additionally, it has

been proposed that every domain forms a new termination by cutting a single strand of dsRNA. The RNA interference process culminates in gene silence by target mRNA degradation, which is carried out by RISC in collaboration with the argonaute (AGO) protein and other effector proteins. Proteins called argonautes are mostly found in eukaryotes, bacteria, and archaea. The Argonaute protein has three important roles: it may identify guide strand termini, use its nuclease ability to cut the target mRNA, or recruit other proteins that are involved in silence. Additionally, RISC with gene silencing takes role in the method of cellular surveillance [7].

3.1 Silencing Genes with RNAi: A Three-Act Play

A potent biological process known as RNA interference (RNAi) targets messenger RNA (mRNA) for destruction in order to control gene expression. The ideal match between the target mRNA and the inserted "guide" sequence is crucial to this process. Here are the three main steps that it takes (Fig. 1):

3.1.1 Act I: Dicer takes the stage

Double-stranded RNA (dsRNA) is either inserted into the cell or arises spontaneously at the start of the play. Dicer, an RNase III family enzyme, intervenes in this situation. Like a proficient cutter, the dicer divides the dsRNA into exact 21-nucleotide segments known as small interfering RNAs (siRNAs). The two-nucleotide overhang on both ends of these siRNAs is a distinctive characteristic.

3.1.2 Act II: RISC assembles and unwinds

Subsequently, the siRNA joins an organization known as the RNA-induced silencing complex (RISC). Argonaute (AGO) protein is the star of the show among the several proteins that make up RISC. The guide strand and the passenger strand are the two strands that remain after the siRNA unwinds inside RISC. The guide strand, nurtured by AGO's PAZ domain, emerges as the play's principal investigator, while the passenger strand frequently deteriorates.

3.1.3 Act III: Guide strand finds and destroys

Equipped with RISC, the guide strand sets out to locate its complementary target within the cellular mRNA pool. The AGO protein in RISC flexes its muscle when a perfect match is detected.




Fig. 1. Plant RNAi pathway mechanism

Its PIWI domain cleaves the target mRNA at a precise place, usually ten nucleotides distant from the match, similar to a pair of molecular shears. This successfully silences the specific gene by making the mRNA unintelligible.

Sometimes an enzyme known as RNA-dependent RNA polymerase (RDRP) can be employed to produce even more siRNAs from the passenger strand rather than discarding it. This increases the effect of silence, but it's not a constant feature of the performance.

Through the strategic placement of complementary guide strands, RNA interference (RNAi) enables cells to precisely control gene expression, resulting in a well-tuned cellular symphony [12].

Dicer cleaves dsRNA to create siRNAs, which are then processed and added to RISC in conjunction with the AGO protein. The target sequence is subsequently identified by the RISC complex, which cleaves it to prevent the desired gene from being expressed [12].

4. GENE CONSTRUCT FOR RNAi

Gene expression is inactivated by RNA interference in a sequence-specific way. One efficient RNAi trigger is double-stranded RNA (dsRNA). Direct introduction of dsRNAs can produce RNAi in worms, flies, and mammalian cells. On the other hand, hairpin RNA-producing construct transformation is typically used to establish RNAi in plants. Chuang and Meyerowitz (2000) were the first to show that an RNAi construct could effectively transcribe RNA in *Arabidopsis thaliana*. They demonstrated how the insertion of an RNA interference vector led to the silence of a homeobox gene that was identical to the corresponding null mutant. An inverted repeat containing target sequences under the direction of a potent promoter makes up the plant RNA interference vector. A spacer fragment lies between the inverted repeat sequences. The substrate of Dicer is a dsRNA structure formed by the inverted repeat region against the target gene. Following transcription, a spacer creates a hairpin RNA's loop and inverted repeat sequences produce a dsRNA structure known as the stem (Fig. 2) Since *Escherichia coli* cannot replicate inverted repeat sequences without a spacer, the spacer sequence makes it easier to produce an RNAi vector [6,15-20].




Fig. 2. Gene construct for RNAi

5. APPLICATIONS OF RNA INTERFERENCE TO ENHANCE AGRICULTURAL YIELD

5.1 Seedless Fruit Development

Fruits without seeds are often preferred by customers because they improve fruit quality and prolong their shelf life [21,22]. Parthenocarp, which is the process by which fruit develops directly from the ovary without fertilization, can produce fruits without seeds. In eggplant, the absence of seeds inhibits browning and pulp texture deterioration [23]. It is possible to intentionally induce the generation of seedless fruits by interfering with the genes responsible for seed formation and seed set. The phytohormones control the process of seed development both geographically and temporally. Fruit that is seedless and produced by phytohormone mutations typically exhibits pleiotropic consequences, such as altered flavor and smaller fruit size. Hence, new, more effective techniques should be used for the development of parthenocarpic fruits [23]. Cucumber (*Cucumis sativus*) [24,25] and tomato (*Solanum lycopersicum*) [26] yields are limited by the development of their seeds. Therefore, edible fruit tissue is a widely desired and appreciated alternative to seed and seed cavities for

customers, breeding firms, and production enterprises. Auxin-dependent plant developmental processes are regulated by transcription factors encoded by auxin response factors (ARFs). It was discovered that mature, unpollinated ovaries had high expression levels of the tomato (*S. lycopersicum*) ARF7 factor, also known as SIARF7. Additional investigation showed that 48 hours after pollination, the expression of *sIARF7* falls. This expression is elevated from the beginning stages of flower development to the production of mature blooms. Fruits that are parthenocarpic were produced via RNA interference-mediated development of transgenic tomato lines that had a downregulated *sIARF7* gene [27]. By suppressing the chalcone synthase (CHS) gene, the initial gene involved in the flavonoid production pathway, using RNA interference (RNAi), Schijlen et al. [28] created seedless tomatoes. Similarly, tobacco (*Nicotiana tabacum cv. xanthi*) produced fruits with fewer or no seeds when the *flavonol synthase* (FLS) gene was silenced by post-transcription, a crucial enzyme for the synthesis of flavonols [29].

In tomato parthenocarpic flower buds, auxin production is specifically expressed by auxin genes. When these genes were silenced using RNA interference, tomatoes developed parthenocarpic fruits along with a few other

auxin-related abnormalities [30]. Small parthenocarpic fruit and flower (spff) mutants in a tomato cultivar were identified and described by Takei et al. [31]. Male sterility with parthenocarpic fruit set development was seen upon linkage analysis and RNAi-based silencing of the *Solyco4g077010* gene, which encodes the receptor-like protein kinase.

5.2 Increasing Nutritional Value

The primary source of necessary nutrients for human diets is plants. Nonetheless, deficiencies in one or more important mineral elements affect about two thirds of the global population [32]. By altering a number of biochemical and physiological processes, RNA interference (RNAi) can be used to raise the quantities of nutrients that crops need. Under the polyunsaturated fatty acid synthesis route, the enzyme omega-3 fatty acid desaturase (FAD3) catalyzes the production of α -linolenic acid (18:3). Soybean (*Glycine max*) and other seed oils become unstable due to α -linolenic acid. By using siRNA to silence FAD3 in soybeans, Flores et al. [33] were able to drastically lower the quantity of α -linolenic acid by 1–3% when compared to other non-transgenic lines. Artificial microRNA (amiFATB) has been used to downregulate the fatty acyl-ACPthioesterase (FATB) gene in *Camelina sativa*, improving the oilseed quality. In comparison to wild-type seeds, the results demonstrated a significant drop in the concentration of total saturated fatty acids, with a 45% reduction in palmitic acid (16:0) and a 38% reduction in stearic acid (18:0) [34].

By using hairpin RNA to silence the *gluB* gene, Kusaba et al. [35] created an *Oryza sativa* cultivar with minimal glutenin levels, known as LGC-1. An elevated concentration of amylose Two starch branching enzyme (SBE) II (SBEIIa and SBEIIb) have been suppressed in *Triticum* endosperm using RNAi to create the *Triticum* cultivar [36]. Starch phosphorylation and dephosphorylation are essential steps in the breakdown of starch in plants. The buildup of starch in the leaves of *Arabidopsis thaliana* and *Zea mays* was caused by the downregulation of phosphoglucan phosphatase (SEX4) and glucan water dikinase (GWD) by RNA interference [37]. *Brassica napus* increased its carotenoid content by RNA-mediated silencing of ϵ -Cyclases (ϵ -CYC). It was discovered that seeds produced from RNAi transgenic *Brassica* lines have high levels of lutein, violaxanthin, zeaxanthin, and β -carotene [38].

RNAi can potentially be used to help crops accumulate minerals. Aggarwal et al. [39] produced *Triticum* grains with high Zn and Fe content and low levels of the antinutrient phytic acid (PA) by RNAi-mediated downregulation of inositol pentakisphosphate kinase (IPK1).

5.3 Abiotic Stress Resistance

Abiotic stress conditions, such as salinity, drought, floods, and temperature changes, are the main causes of the global decline in the production of many important crops. Scientists are under tremendous pressure to produce crop kinds that are resistant to stress due to the ever-changing environment and the increasing need for food from a growing population. The capacity of non-coding RNAs (ncRNAs), specifically microRNAs (miRNAs), to create crops resistant to abiotic stress. ncRNAs regulate critical functions like as transcription, detoxification, and development by adjusting gene expression at various levels. This affects how crops react to oxidative stress, salt, cold, and drought. This process is shown by the overexpression of miR393 in *Arabidopsis* seedlings subjected to different stimuli.

Researchers can create cultivars that are more resilient to stress by using RNA interference (RNAi) technology to target particular non-coding RNAs (ncRNAs) and the genes that are linked to them. Under difficult environmental circumstances, this strategy has great potential to increase the production and sustainability of important crops like rice, wheat, lentils, and sugarcane. plant responses to environmental stressors, particularly salt and drought, and the function of microRNAs (miRNAs) [1].

It indicates that particular miRNAs are expressed by a variety of plant species, including important crops like rice, wheat, and maize, and that these miRNAs affect the expression of these plant species in response to salinity. Important participants in this adaptation are identified as members of notable miRNA families, such as miR156 and miR398, respectively. The paragraph also highlights studies that exploit genome-wide methods to investigate miRNA-mediated gene regulation in drought-stressed plants, including rice and sorghum as examples. This work highlights the possibility of comprehending miRNA activities to enhance plant resilience and stress tolerance overall [1].

Two effective methods for enhancing rice's tolerance to drought through natural processes

are:- In the first method, superoxide dismutase (SOD) activity is elevated and drought tolerance is eventually raised as a result of RNA interference (RNAi) silencing the receptor for activated C-kinase 1 (RACK1). The second technique uses maize short interfering RNA (siRNA) to inhibit rice's farnesyltransferase/squalene synthase (SQS), which raises sterol levels, decreases stomata-mediated water loss, and improves rice's resistance to drought. These techniques greatly increase rice's resistance to drought in both the vegetative and reproductive phases, providing important resources for creating crops that can tolerate changing climate conditions. The *OstZTF1* gene's critical function in rice stress tolerance. Surprisingly, silencing this gene increased tolerance to low water and high salt environments. It's interesting to note that low *OstZTF1* expression regulates hormone synthesis at the molecular and cellular levels in response to salt stress, aiding plants in preserving homeostasis. The study emphasizes the importance of microRNAs (miRNAs) in plant stress responses beyond *OstZTF1*. Various plant species use distinct miRNAs that are sensitive to heat and cold to control target genes, which increases resistance to a range of abiotic stresses. The ability to develop stress-resistant crops using *OstZTF1* and miRNA manipulation [1].

5.4 Biotic Stress Tolerance

Two core lncRNAs, GhIncNAT-ANX2- and GhIncNAT-RLP7-silenced seedlings, demonstrated greater resistance against *Verticillium dahliae* and *Botrytis cinerea*, according to functional analysis. This enhanced resistance may have been caused by increased expression of the LOX1 and LOX2 genes. Cotton leaf curl virus, which contains a sense transcript BC1 that may cause pathogenicity symptoms, affects cotton, one of the major cash crops. A BC1 antisense construct was inserted into the *Nicotiana tabacum* plant, and upon expression, the BC1 anti-transgenic tobacco exhibited resistance against the Begomovirus-causing viruliferous whitefly (*Bemisia tabaci*) [15].

5.4.1 Bacterial disease resistance

Bacteria are basic tiny creatures that infiltrate plants by rain, wind, soil, seed dissemination, or other mechanisms. Unlike viruses, bacteria do not require insects as their vectors. Numerous plant diseases caused by bacterial pathogens

have a significant negative impact on crop productivity and result in significant yearly losses for the world's farmers. One of the main problems in the field of crops like tomatoes, soybeans, and bananas is bacterial illness [1].

5.4.2 Viral disease resistance

Numerous plant viruses are highly transmissible and frequently have severe impacts on plants. Due to their quick growth and dissemination, several of them significantly reduce crop output and quality. Plant resistance to viruses is largely dependent on RNA silencing, with a variety of silencing components contributing to antiviral defense. The small RNA-directed RNA degradation mechanism targets both DNA and RNA viruses. RNA silencing, a key antiviral mechanism in plants, is dependent on DICER-like enzymes cleaving viral dsRNA into virus-derived small interfering RNAs (vsiRNAs). Even now, lines of virus-resistant squash (*Cucurbita*) and papaya (*Carica papaya*) that were deregulated in the 1990s are still grown. Even though it was previously believed that viruses infect plants through the development of a virus-derived protein (coat protein-mediated resistance, or CPMR) before the viruses really settle and begin to disrupt the plant's RNA silencing mechanism. The first use of gene silencing was to create virus-resistant plant types. Plants have developed antiviral tactics that resemble the RNA silencing processes found in nature. This was initially shown when researchers created the potato virus Yresistan, which caused plants to produce the viral proteinase gene's RN transcripts [1,5].

Plants can also be engineered to produce dsRNAs that silence vital genes in insect pests and parasitic nematodes. Immunity to other viruses, such as the Cucumber and Tobacco Mosaic Virus, Tomato Spotted Wilt Virus, Bean Golden Mosaic Virus, Banana Bract Mosaic Virus, Rice Tungro Bacilliform Virus, and many others, has also been demonstrated [5].

5.4.3 Fungal diseases

Fungal diseases pose a serious hazard to crops because they can produce toxic mycotoxins that are dangerous to human and animal health in addition to causing enormous yield losses. Utilizing RNA interference (RNAi) technology to build resistance to fungal infections has been studied by researchers. According to one study, tobacco plants that had their glutathione S-transferase (GST) gene silenced by RNA interference (RNAi) were more resilient to the

Table 1. RNA interference to control fungal diseases

Fungal pathogen	Disease	Target gene	Outcome
<i>Phytophthora parasitica</i> var. <i>nicotianae</i>	Black shank disease	Glutathione S-transferase (GST)	Increased resistance in tobacco plants
<i>Blumeria graminis</i>	Powdery mildew of barley and wheat	Not specified	RNAi can be an effective way to control the pathogen

fungal pathogen *Phytophthora parasitica* var. *nicotianae*, which causes black shank disease. In different research, the fungus *Blumeria graminis*, which causes barley and wheat powdery mildew, was targeted via RNA interference (RNAi). The outcomes demonstrated that using RNA interference to control fungal diseases can be successful [1].

5.4.4 Insect and nematode resistance

Insect functional genomics research and insect pest management both benefit from the use of RNA interference (RNAi). Insects should be able to absorb dsRNA by grazing on plants. According to recent findings, dsRNA that was given to insects as part of their food efficiently down-regulated the genes that the insects were targeting. More importantly, it has been demonstrated that the hdRNAi-1 approach silences insects that feed on plants, paving the path for the creation of a new breed of insect-resistant crops. Western corn rootworm (WCR) resistance was demonstrated using transgenic corn plants designed to produce dsRNAs for the tubulin or vacuolar ATPase genes. In this study, resistance manifested itself as decreased mortality, stunting, and damage to plants. Larval development of cotton bollworm (*Helicoverpa armigera*) has been seen to be slowed when they feed on plant material producing dsRNA specific to the cytochrome P450 gene (CYP6AE14). Therefore, species-specific insecticide, or hdRNAi-1 technology, is another name for it and a possible substitute for conventional pesticides. In addition to stable RNAi transgenic plants, insect damage can be reduced by applying dsRNA to plant leaves and dsRNA-enriched soil. The preceding evaluation covered studies evaluating the effectiveness of dsRNA spray against insect pests. However, further knowledge of the RNAi process in insects and their genetic data is required to fully grasp the promise of this technique [13].

Some crucial questions that need to be answered are: can an insect's dsRNA be transferred back to plants through insect saliva? Can an insect's RNAi mechanism be passed

from one feeding stage (larvae or adults) to the next growth stage or generation (pupa or egg)? How long does it take an insect to effectively silence its genes and die from its first meal on an RNAi transgenic plant? Furthermore, dsRNA in vertebrates has the ability to trigger immunological responses on its own, perhaps providing resistance to insect progeny. Further investigation is necessary to gain a deeper comprehension and make appropriate adjustments to the hd-RNAi technique [13].

Engineering plants to be resistant to nematodes is a crucial component of agricultural progress. Two methods may be used to achieve this: (i) utilizing the hdRNAi-1 approach to increase plant resistance; (ii) using HGS-hpRNAi to silence specific plant genes. When it comes to these two methods, hdRNAi-1 works best. Through the consumption of transgenic plant tissue, dsRNA or its siRNAs are transferred from plant to nematode in this manner. Compared to cyst nematode (CN), transgenic plants expressing hdRNAi-1 have demonstrated greater efficacy against root-knot nematode (RKN). It has been demonstrated that regulating this nematode is possible with transgenic tobacco plants that produce dsRNA targeting a putative zinc finger transcription factor gene (MjTis11) in RKN (*Meloidogyne javanica*). hdRNAi-1 was utilized to target up to four important CN (*Heterodera schachtii*) genes by expressing several hpRNA constructs in *Arabidopsis*, therefore increasing the efficiency of CN nematode control. As a result, there were less adult CN nematode females eating on this plant. Higher gene silencing effectiveness in CN will be possible with a deeper comprehension of the RNAi system in plant parasitic nematodes [13].

5.4.5 Improving resistance against parasitic weeds

It has been challenging to manage plant parasitic weeds using genetic engineering since no genetic loci granting resistance in agricultural plants have yet to be discovered. Therefore, current study has looked at the possible

applications of RNA interference in this field. The transmission of signals from the host plant to the parasite weeds is required for hdsRNAi-1 to function. Thus, the distribution of the RNAi signal was studied. Transgenic lettuce roots producing hpRNA carrying a portion of the GUS gene were made available for parasitisation by the *Triphysaria versicolor* parasite, which expresses the GUS reporter gene. *Triphysaria* roots connected to non-transgenic lettuce displayed complete GUS activity when stained for GUS activity, whereas those parasitizing RNAi transgenic lettuce lacked GUS activity. This suggests that a silencing signal can quiet a

specific gene by traveling from host to parasite [13].

Mannose 6-phosphate reductase gene (M6PR) hpRNA constructed from *Orobanche aegyptiaca* is present in transgenic tomato plants, and this construct has been shown to be efficient against the parasite by killing its tubercles that are connected to tomato plants. Additional research also demonstrated that parasites may use their haustorium to absorb particular RNAi from their host. These investigations unequivocally showed the potential of hdsRNAi-1 technology for controlling plant parasites in the future [13].

Table 2. Crops with improved stress tolerance through RNAi

Trait(s)	Crop Improved	Resistance against	Targeted Gene(s)	References
Bacterial resistance	<i>A. thaliana</i>	<i>Agrobacterium tumefaciens</i>	iaaM	[40]
	Citrus limon	<i>Pseudomonas syringae</i>	ipt	[41]
		<i>Xanthomonas citri</i>	PPRL CalS1	[42]
Abiotic stress tolerance	<i>A. thaliana</i>	Osmotic tolerance	WZY2	[43]
	<i>T. aestivum</i>	Salt tolerance	TaPUB-1	[44]
	<i>N. tabacum</i>	Salt tolerance	Nt ε-LCY	[45]
	<i>O. sativa</i>	Salt tolerance	OsPEX11	[46]
	<i>B. rapa</i>	Salt tolerance	GIGANTEA (GI)	[47]
	<i>A. thaliana</i>	Drought tolerance	PAD4 LSD1 EDS1	[48]
	<i>O. sativa</i>	Drought tolerance	OsGRXS17	[49]
	<i>O. sativa</i>	Drought tolerance	OsDSR-1	[50]
	<i>O. sativa</i>	Drought tolerance	OsERF101	[51]
	<i>S. lycopersicum</i>	Drought and salt tolerance	SlbZIP1	[52]
<i>N. tabacum</i>	Drought tolerance	BrDST71	[53]	
Virus resistance	<i>N. bethamiana</i>	Tomato yellow leaf curl Thailand virus (TYLCTV)	GSA	[54]
	<i>Glycine max</i>	Soybean mosaic virus (SMV)	eIF4E1	[55]
	<i>O. sativa</i>	Rice tungroo bacilliform virus (RTBV) Rice tungroo spherical virus (RTSV)	Coat protein 3 CP3	[56]
	<i>Nicotiana bethamiana</i>	Chilli-infecting begomoviruses	AC1 AC2 βC1	[57]
	<i>Triticum spp.</i>	Triticum mosaic virus (TMV)	Coat protein (CP)	[58]
	<i>Oryza sativa</i>	Rice black streak dwarf virus (RBSDV)	S7-2 S8	[59]
	<i>Glycine max</i>	Soybean mosaic virus (SMV)	SMV P3 cistron	[60]
		Mungbean yellow mosaic virus (MYMIV)	CP	[61]
	<i>Arachis hypogaea</i>	Tobacco streak virus (TSV)	CP	[62]
	<i>Solanum tuberosum</i>	Potato virus X (PVX), Potato virus Y (PVY)	CP	[63]

Trait(s)	Crop Improved	Resistance against	Targeted Gene(s)	References
		Potato virus S (PVS)		
Virus resistance	<i>Brassica rapa</i>	<i>Tetranychus urticae</i>	COPB2	[64]
	<i>A. thaliana</i>	<i>Myzus persicae</i>	MyCP	[65]
	Lettuce	<i>B. tabaci</i>	V-ATPase	[66]
	<i>N. tabacum</i>	<i>Bemisia tabaci</i>	AChE	[67]
	<i>Nicotiana bethamiana</i>	Chilli-infecting begomoviruses	EcR AC1 AC2 βC1	[68]
	<i>S. lycopersicum</i>	<i>Helicoverpa armigera</i>	HaCHI	[69]
Fungal resistance	<i>Glycine max</i>	<i>Phytophthora sojae</i>	GmSnRK1.1	[70]
	<i>S. lycopersicum</i>	<i>F. oxysporum</i>	ODC	[71]
	<i>S. tuberosum</i>	<i>Phytophthora infestans</i>	Avr3a	[72]
	<i>T. aestivum</i>	<i>Fusarium graminearum</i>	Chs 3b	[73]
	<i>Musa spp.</i>	<i>F. oxysporum f. sp. cubense (Foc)</i>	Foc velvet protein	[74]
	<i>N. tabacum</i>	<i>Sclerotinia sclerotiorum</i>	Chs	[75]
	<i>S. lycopersicum</i>	<i>F. oxysporum</i>	Fow2 chs V	[76]
	<i>Z. mays</i>	<i>A. flavus</i>	Amy1	[77]
	<i>Zea mays</i>	<i>Aspergillus flavus</i>	ZmPRms	[78]
	<i>S. tuberosum</i>	<i>Phytophthora infestans</i>	PVS1 PVS2 PVS3	[79]
		<i>Alternaria solani</i>	PVS4	
	<i>S. lycopersicum</i>	<i>F. oxysporum</i>	Fmk1 Hog1 Pbs2	[80]
	<i>O. sativa</i>	<i>Magnaporthe oryzae</i>	MoABC1 MoMAC1	[81]
		<i>Rhizoctonia solani</i>	MoPMK1 RPMK1-1 RPMK1-2	[82]
	Nematodes Resistance	<i>A. thaliana</i>	<i>M. incognita</i>	Mi-msp3 Mi-msp 5 Mi-msp18 Mi-msp24
<i>S. lycopersicum</i>		<i>Meloidogyne incognita</i>	Mi-cpl1	[84]
<i>N. benthamiana</i>		<i>Radopholus similis</i>	Rs-cps	[85]
<i>S. lycopersicum</i>		<i>M. incognita</i>	PolA1	[86]
<i>Glycine max</i>		<i>Heterodera glycines</i>	Hg16B09 HgY25 HgPrp17	[87] [88]

5.5 Development of Male Sterile Lines

Male sterility has also been produced by RNA interference, which is useful for the hybrid seed market. With RNA interference (RNAi), genes that are expressed only in tissues involved in pollen generation may be targeted. For example, TA29 gene expression, which is required for pollen formation, can be inhibited in order to create male sterile tobacco strains. Additionally, RNA interference (RNAi) was utilized to stop Msh1 from expressing in tobacco and tomatoes, which led to changes in the mitochondrial DNA linked to cytoplasmic male sterility that occurs spontaneously [5].

5.6 Modification of Flower Color and Scent by RNAi-Mediated Gene Silencing

A branch of horticulture called "flower farming" is dedicated to the production of flowers and other decorative plants. The need for flowers in a variety of hues and designs for smells and decorating has grown in the modern day. This may be done by employing RNA interference technology to silence the genes that code for pigment. By utilizing RNA interference (RNAi) to repress a cDNA encoding the chalcone isomerase (CHI) gene that was obtained from *N. tabacum* petals, flavonoid components in flower

petals were altered and pigmentation was reduced [89]. Similarly, the RNAi construct TrCHS1 targeting chalcone synthase (CHS) has been shown to cause flower color modification in liliaceous ornamental *Tricyrtis* sp. [90]. Ornamental gentian plants' flower colors changed as a result of RNA interference-mediated silencing of three anthocyanin biosynthesis genes: chalcone synthase (CHS), anthocyanidin synthase (ANS), and flavonoid 3O,5-O-hydroxylase (F3O5 OH) [91]. The bloom of gentian plants is naturally bright blue in hue. The pigment known as "gentiodelphin," a polyacrylate delphinidin, accumulates in the petals of gentian plants and imparts color to the flowers. Important enzymes for gentiodelphin production are flavonoid 3O,5O-hydroxylase (F3O5 OH) and anthocyanin 5,3O-aromatic acyltransferase (5/3OAT), whose downregulation by RNA interference modifies floral color [92]. In order to reconstruct the delphinidin pathway, He et al. [93] initially identified two chrysanthemum cultivars and separated seven genes involved in anthocyanin biosynthesis: CmCHS, CmF3H, CmF3OH, CmDFR, CmANS, CmCHI, and Cm3GT. Furthermore, cyanidin concentration was elevated in chrysanthemum flowers with brighter red petals due to the overexpression of the *Senecio cruentus* F3O5OH (PCFH) gene and the inhibition of the CmF3OH gene; however, no reports of delphinidin accumulation have been made.

5.7 Enhanced Fruit Shelf Life

Fruits and vegetables are more prone to spoiling owing to their nature and composition than grains. India loses between thirty and forty percent of the overall yield of fruits and vegetables to spoiling, according to the Agricultural Research Data Book 2004. Consequently, as another essential agronomic attribute that may reduce fruit and vegetable spoiling and, in turn, lessen agricultural loss, we must use strategies that might extend the shelf life of fruits and vegetables. By focusing on the genes that code for the ethylene production pathway, RNA interference technology (RNAi) extended the tomato's shelf life. When a tomato was given a dsRNA unit, the ACC oxidase gene's expression was inhibited. It was discovered that the transgenic plants' matured fruits significantly decreased the rate at which ethylene was produced, extending the tomato's shelf life. Using RNA interference technology, two ripening-specific N-glycoprotein modifying enzymes were identified in tomatoes: α -

mannosidase (α -Man) and β -d-N-acetylhexosaminidase (β -Hex). This illustrated how inhibiting these genes prolongs the tomato's storage life by delaying the ripening process [1].

5.8 Secondary Metabolite Production

Secondary metabolites from plants are utilized in medications, food additives, perfumes, colors, and insecticides. Many different genes control the biosynthesis of secondary metabolites, however occasionally certain unwanted substances can impede this process. In addition to being useful for manipulating secondary metabolites, RNA interference (RNAi) may be employed to effectively decrease the production of those molecules [94]. In the opium poppy (*Papaver somniferum*), Allen et al. [95] revealed that RNA interference (RNAi)-mediated silencing of numerous genes involved at different points in a complicated biochemical process replaced morphine with the non-narcotic alkaloid (S)-reticuline. To inhibit the expression of every member of the codeine reductase (COR) gene family, they created hpRNA. As a result, morphine, codeine, and opium were substituted with (S)-reticuline, a non-narcotic alkaloid precursor, to create transgenic lines. The central nervous system, respiratory system, and circulatory system are all naturally stimulated by caffeine. Moreover, it lowers the risk of throat, mouth, and liver cancer. In addition, consuming too much of it can lead to health problems including tremors in the muscles, anxiety, upset stomach, sleeplessness, and restlessness. Three enzymes, namely CaXMT1, CaMXMT1 (theobromine synthase), and CaDXMT1 (caffeine synthase), are involved in the manufacture of caffeine in coffee plants. The CaMXMT1 gene was silenced by RNA interference (RNAi) and the resulting 70% reduction in caffeine concentration suggests that RNAi technology can be used to produce decaffeinated coffee beans [96]. Similarly, RNA interference (RNAi) was used to downregulate the caffeine synthase (CS) gene in order to create transgenic tea (*Camellia sinensis*) lines with low caffeine content [97].

A well-known Chinese herb that is also utilized in other Asian nations is salvia miltiorrhiza. By downregulating the Chalcone synthase (CHS) gene, the first enzyme in flavonoid biosynthesis, and treating it with salicylic acid as an elicitor, phenolic acid production was increased. The synthesis of flavonoids was significantly reduced while the level of phenolic acid increased, according to the results [98]. Peltate glandular

trichomes (PGTs) are small, specialized structures that are present in various aromatic plants, including spearmint (*Mentha spicata*), and are involved in the generation of secondary metabolites. In their investigation into the activity of transcription factors in the pathway leading to the manufacture of secondary metabolites, Wang et al. [99] identified and described the MsYABBY5 gene that is expressed in PGT. Following the inhibition of the MsYABBY5 gene, terpene synthesis increased, indicating that transcription factors with encodes function as a negative regulator of secondary metabolite production. The DE-ETIOLATED-1 (DET1) gene, a negative regulator of photomorphogenesis, was suppressed through RNA interference (RNAi) in the embryonic callus of papaya (*Carica papaya* L.) plants in order to examine its effects on the expression of a gene involved in the secondary metabolite biosynthesis pathway. The findings suggested a connection between the photo-regulated pathway and secondary metabolite synthesis [100].

5.9 Deletion of Allergens from Food Crops

Additionally, plants have a large number of toxic and undesired substances, the removal of which is an expensive and time-consuming procedure. An effective technique for ridding the plants of toxins is the RNAi approach. By downregulating three different methylation stages of the caffeine biosynthesis pathway and suppressing the expression of the 7-N-methylxanthine-methyltransferase gene (CaMXMT1), the amount of caffeine in transgenic coffee beans is lowered by up to 70%. Tuber of cassava (*Manihot esculenta*) contain harmful substances called cyanogenic glucosides. The amount of cyanogenic glucosides can be decreased by about 90% by down-regulating the enzymes CYP79D1 and CYP79D2, which are involved in the production of linamarin and lotaustraline. By simultaneously suppressing the ASPARAGINE SYNTHETASE genes (StAs1 and StAs2), commercial potatoes with much decreased asparagine—a primary precursor of the brain toxin acrylamide—can be created using a similar strategy. A hypersensitive reaction to ordinarily safe protein components in food, primarily via the immunoglobulin (IgE) mechanism, is known as a food allergy. Technology based on RNA interference is also helpful in reducing these allergies. It has been utilized to suppress Mal-d-1, an apple allergen that exhibits IgE antibodies. Intron spliced hpRNA constructs containing the Mal-d-1

inverted repeat sequence were used to alter apple plants. This resulted in a 10-fold reduction in Mal-d-1 leaf expression without affecting normal plant development. Tearless onions may also be created by downregulating the conversion of 1-propenylsulfenic acid to propanthial S-oxide, a factor that causes tears and is lachrymatory, using RNA silencing technology [10].

5.10 Change/Alteration in Plant Architecture

A plant's success is largely dependent on its architecture, or the blueprint for its form and structure. Every plant, no matter how big or small, has a distinct design that determines important agronomic characteristics including height, branching, yield, and even stress tolerance. Scientists can enhance crops by fine-tuning plant architecture by comprehending this relationship. A potent technology that provides fine control over genes determining architecture is RNA interference (RNAi). For example, rice that has had the OsDWARF4 gene silenced has stronger, shorter plants with erect leaves, which facilitates harvest. RNAi that targets the ODC gene in tobacco change the structure of the plant, decreasing its ability to withstand stress while providing information about its function in development. Plant architecture is likewise shaped by microscopic regulators called microRNAs (miRNAs). In maize, modifying Cg1 miRNA postpones blooming, which may be advantageous for breeding techniques. It's interesting to note that changing Cg1 in poplar trees improves the generation of biofuel by encouraging desirable features like shorter internodes and less lignin. Additionally, RNAi can meet certain demands.

Plants with low lignin content that is simpler to convert to bioethanol are produced when lignin production genes are silenced. In a similar vein, RNAi can enhance leaf plucking in mulberry and tea plants and make roses thornless. This intriguing look into the field of manipulating plant architecture emphasizes the possibilities of RNA interference. Through adjustments to the hidden architect, crops with enhanced stress tolerance, yields, and customized traits may be produced, guaranteeing food security and sustainable agriculture in an evolving global landscape [1].

5.11 Quality Improvement

The RNAi technology not only increases the food's nutritional value but also fortifies plants

against biotic and abiotic stress. One of the most crucial indicators of food quality is thought to be the amount of amino acids it contains. A significant advancement in the use of RNA interference (RNAi) in plants has been made possible by research into the metabolic processes involved in the production of the main amino acids. The model tobacco plant containing the *Arabidopsis* S-adenosyl-L-methionine synthase gene was repressed with RNA interference (RNAi), which led to an increase in L-methionine content in transgenic plants. Maize plants by inhibiting the lysine-degrading gene in maize plants led to transgenic plants having a greater lysine content than wild-type plants. Numerous additional investigations have also noted the silencing impact on plastid genes that increase the levels of aspartate, glutamate, and lysine in plants, such as NbAsp5 and NbPAT [12].

6. LIMITATIONS AND CHALLENGES ASSOCIATED WITH THE UTILIZATION OF RNAI TECHNOLOGY

- **Delivery:** One of the biggest challenges still is getting siRNA molecules into the target cells. Since siRNA is fragile, enzymes may break it down before it reaches its intended target. Moreover, siRNA has trouble passing across cell membranes. Researchers are looking at several techniques to increase the specificity and effectiveness of delivery, such as using nanoparticles and viral vectors.
- **Off-target effects:** siRNA may inadvertently quiet genes that are not the intended target, which might have unforeseen consequences. To reduce this danger, highly precise siRNA molecules must be designed.
- **Immune response:** The immune system may be activated by the introduction of foreign RNA molecules, which might lessen the efficacy of RNAi treatment. Methods to lessen immunogenicity are being researched.
- **Short half-life:** The body breaks down siRNA molecules quickly, reducing the amount of time they may be active. SiRNA can be chemically altered to increase its stability.
- **Specificity:** SiRNA can silence unwanted genes with similar sequences even when it is well-designed. Unwanted adverse effects might result from this.

7. CONCLUSION

Malnutrition is a serious issue, particularly in underdeveloped nations (WHO 2000). Approximately 792 million people worldwide—including 20% of those living in developing nations are chronically malnourished (FAO, 2000). It is imperative to create bio-fortified fruits, vegetables, and cereals by enriching them with nutritionally necessary components and compounds including vitamins, fatty acids, critical minerals, antioxidants, and amino acids in order to assure a healthy diet for a healthy world.

Crop plants should be produced with stress (biotic and abiotic) tolerance as one of their most essential agronomical features. Consequently, it is necessary to create plant types resistant to pests and phytopathogens as well as to shifting climatic factors including high temperatures, flooding, droughts, oxidative stress, and modifications to the composition of the soil (heavy metal buildup, salinity, declining fertility, etc.). In order to feed the growing population, these plant kinds will be essential for food security. Studies using RNA interference have demonstrated the promise of this technology to enhance crops and address issues with hunger, malnutrition, and food security.

Extensive investigations have been conducted to identify tiny non-coding regulatory RNAs, uncover their biogenesis, and evaluate their impact on gene suppression; nonetheless, several details concerning the intricacy of the RNA interference pathway and its constituent parts remain to be clarified. But the uses of short non-coding RNA research are setting new standards for agricultural advancement, which is enhancing living standards.

RNA interference (RNAi) is a process that results in heterochromatin development through chromatin remodeling and transcriptional gene silence. These changes might get inherited and have unfavorable consequences that put biosafety at risk. Crop plants created by RNA interference should have their risks to the environment and food safety evaluated. Thus, vectors must be specifically customized in accordance with the requirement for crop development.

However, for the functional genomics and biotechnology of perennial plants, RNAi can be quite useful. Improved agricultural output is required to fulfill the world's demand for food,

fuel, and fiber; these needs will be met by the development of RNAi technology. Applications of RNA interference (RNAi) in crop science offer several ways to improve crop quality, including increased nutritional content and stress tolerance. Therefore, RNAi has enormous potential to greatly increase agricultural productivity. Future RNA interference practice and research should concentrate on finding new targets, investigating combination treatments, improving the safety, efficacy, specificity, and delivery of RNAi medicines, and broadening their uses across a range of illnesses and domains. Unlocking RNAi's full potential as an effective tool for fundamental research and therapeutic intervention will require addressing these issues.

8. IMPLICATIONS FOR FUTURE RESEARCH AND PRACTICE

- **Efficiency and Specificity:** Improving RNAi methods' efficiency and specificity should be the focus of future study. Better delivery methods must be created to target particular tissues or cell types with the fewest possible side effects.
- **Delivery Systems:** The development of delivery systems is essential to the practical use of RNA interference treatments. Enhancing delivery vehicles including lipid nanoparticles, viral vectors, and exosomes in terms of stability, targeting, and safety profiles should be the main goal of research.
- **Target Identification:** It's critical to find new RNA interference targets for a range of disorders. The functions of non-coding RNAs, circular RNAs, long non-coding RNAs, and other RNA species in disease pathogenesis should be investigated in future studies.
- **Combination Therapies:** Researching the potential benefits of combining RNA interference (RNAi) with other therapeutic modalities, such as antibodies, small compounds, or immunotherapies, may help improve treatment outcomes and get beyond resistance mechanisms.
- **Off-Target Effects:** The safety and effectiveness of RNAi treatments depend on their capacity to recognize and reduce these side effects. To precisely measure off-target impacts, future research should concentrate on enhancing computational prediction algorithms and experimental validation methods.

- **Immunogenicity:** For RNAi therapies to remain safe and effective over the long term, it is critical to investigate their immunogenicity. For clinical trials to be successful, it will be crucial to understand how RNA interference (RNAi) elicits immune responses and to devise methods to reduce immunogenicity.
- **Disease Applications:** It may be possible to improve patient outcomes by applying RNA interference (RNAi) therapeutics to a larger spectrum of diseases than those that are presently being targeted, such as cancer, genetic disorders, viral infections, neurodegenerative diseases, and metabolic disorders.
- **RNAi in Agriculture:** Exploring the application of RNAi technology for disease resistance, pest management, and crop enhancement in agriculture has the potential to transform sustainable farming methods and food security.
- **Regulatory Approval:** For RNAi treatments to be widely used in clinical practice, it will be essential to streamline the regulatory processes for them and solve issues with manufacturing, quality control, and scalability.
- **Patient Stratification:** By creating methods for patient stratification based on biomarkers and molecular profiles, treatment results for RNA interference (RNAi) treatments may be enhanced.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Gautam P, Kumar R, Feroz Z, Vijayaraghavalu S, Kumar M. RNA interference technology in plants: mechanisms and applications in crop improvement. In *Plant genomics for sustainable agriculture*. Singapore: Springer Nature Singapore. 2022 Jun;265-290.
2. Saurabh S, Vidyarthi AS, Prasad D. RNA interference: concept to reality in crop improvement. *Planta*. 2014 Mar;239:543-64.
3. Williams M, Clark G, Sathasivan K, Islam AS. RNA interference and its application in crop improvement. *Plant tissue culture and Biotechnology*. 2004;1:18.

4. Jagtap UB, Gurav RG, Bapat VA. Role of RNA interference in plant improvement. *Naturwissenschaften*. 2011 Jun;98:473-92.
5. Meena AK, Verma LK, Kumhar BL. RNAi, it's mechanism and potential use in crop improvement: A review. *International Journal of Pure and Applied Bioscience*. 2017;5(2):294-311.
6. Dheemanth TL, Ramachandra RK, Chavan MO, Prakash BG. RNA interference in crop improvement. *Trends Biosci*. 2017;10(46):9322-35.
7. Rajput M, Choudhary K, Kumar M, Vivekanand V, Chawade A, Ortiz R, Pareek N. RNA interference and CRISPR/Cas gene editing for crop improvement: Paradigm shift towards sustainable agriculture. *Plants*. 2021 Sep 14;10(9):1914.
8. Zhu KY, Palli SR. Mechanisms, applications, and challenges of insect RNA interference. *Annual Review of Entomology*. 2020 Jan 7;65:293-311.
9. Kusaba M. RNA interference in crop plants. *Current opinion in Biotechnology*. 2004 Apr 1;15(2):139-43.
10. Chakrabarti J, Das M. RNA Interference (RNAi): An Approach to Plant Protection and Crop Improvement A Review.
11. Dubrovna OV, Mykhalska SI, Komisarenko AG. Use of RNA Interference Technology for Improving Economically Valuable Traits of Cereal Crops. *Cytology and Genetics*. 2023 Dec;57(6):587-610.
12. Singh A, Roychoudhury A. RNA Interference (RNAi) Technology: An Effective Tool in Plant Breeding. In *Advanced Crop Improvement, Theory and Practice*. Cham: Springer International Publishing. 2023 Aug 2;1:309-320.
13. Senthil-Kumar M, Mysore KS. RNAi in plants: Recent developments and applications in agriculture. *Gene Silencing: Theory, Techniques and Applications*. 2010:183-99.
14. Pathak K, Gogoi B. RNA interference (RNAi): Application in crop improvement: A review. *Agricultural Reviews*. 2016;37(3):245-9.
15. Tilahun T, Bezie Y, Kerisew B, Taye M. The application of antisense technology for crop improvement: A review. *Cogent Food & Agriculture*. 2021 Jan 1;7(1):1910157.
16. Ali N, Datta SK, Datta K. RNA interference in designing transgenic crops. *GM Crops*. 2010 Jul 28;1(4):207-13.
17. Campbell TN, Choy FY. RNA interference: past, present and future. *Current Issues in Molecular Biology*. 2005 Jan;7(1):1-6.
18. Xu W, Jiang X, Huang L. RNA interference technology. *Comprehensive Biotechnology*. 2019:560.
19. Das S, Marwal A, Choudhary DK, Gupta VK, Gaur RK. Mechanism of RNA interference (RNAi): current concept. *Int Proc Chem Biol Environ Eng*. 2011;9:240-5.
20. Dykxhoorn DM, Lieberman J. The silent revolution: RNA interference as basic biology, research tool, and therapeutic. *Annu. Rev. Med*. 2005 Feb 18;56:401-23.
21. Alam I, Salimullah M. Genetic engineering of eggplant (*Solanum melongena* L.): Progress, controversy and potential. *Horticulturae*. 2021 Apr 11;7(4):78.
22. Pandolfini T. Seedless fruit production by hormonal regulation of fruit set. *Nutrients*. 2009 Dec;1(2):168-77.
23. Varoquaux F, Blanvillain R, Delseny M, Gallois P. Less is better: new approaches for seedless fruit production. *Trends in Biotechnology*. 2000 Jun 1;18(6):233-42.
24. Denna DW. Effects of genetic parthenocarpy and gynocercious flowering habit on fruit production and growth of cucumber (*Cucumis sativus* L.).
25. Tiedjens VA. Sex ratios in cucumber flowers as affected by different conditions of soil and light.
26. Falavigna A, Soressi GP. Influence of the pat-sha gene on plant and fruit traits in tomato (*L. esculentum* Mill.). *Modern trends in tomato genetics and breeding*. 1987 Sep 2:128.
27. De Jong M, Wolters-Arts M, Feron R, Mariani C, Vriezen WH. The *Solanum lycopersicum* auxin response factor 7 (SIARF7) regulates auxin signaling during tomato fruit set and development. *The Plant Journal*. 2009 Jan;57(1):160-70.
28. Schijlen EG, de Vos CR, Martens S, Jonker HH, Rosin FM, Molthoff JW, Tikunov YM, Angenent GC, van Tunen AJ, Bovy AG. RNA interference silencing of chalcone synthase, the first step in the flavonoid biosynthesis pathway, leads to parthenocarpic tomato fruits. *Plant Physiology*. 2007 Jul 1;144(3):1520-30.
29. Mahajan M, Ahuja PS, Yadav SK. Post-transcriptional silencing of flavonol synthase mRNA in tobacco leads to fruits with arrested seed set. *PLoS One*. 2011 Dec 1;6(12):e28315.

30. Molesini B, Pandolfini T, Rotino GL, Dani V, Spena A. Aucsia gene silencing causes parthenocarpic fruit development in tomato. *Plant Physiology*. 2009 Jan 1;149(1):534-48.
31. Takei H, Shinozaki Y, Yano R, Kashojiya S, Hernould M, Chevalier C, et al. Loss-of-function of a tomato receptor-like kinase impairs male fertility and induces parthenocarpic fruit set. *Frontiers in Plant Science*. 2019 Apr 16;10:436826.
32. White PJ, Broadley MR. Biofortification of crops with seven mineral elements often lacking in human diets—iron, zinc, copper, calcium, magnesium, selenium and iodine. *New Phytologist*. 2009 Apr;182(1):49-84.
33. Flores T, Karpova O, Su X, Zeng P, Bilyeu K, Slepser DA, Nguyen HT, Zhang ZJ. Silencing of Gm FAD3 gene by siRNA leads to low α -linolenic acids (18: 3) of fad3-mutant phenotype in soybean [*Glycine max* (Merr.)]. *Transgenic Research*. 2008 Oct;17:839-50.
34. Ozseyhan ME, Li P, Na G, Li Z, Wang C, Lu C. Improved fatty acid profiles in seeds of *Camelina sativa* by artificial microRNA mediated FATB gene suppression. *Biochemical and Biophysical Research Communications*. 2018 Sep 5;503(2):621-4.
35. Kusaba M, Miyahara K, Iida S, Fukuoka H, Takano T, Sassa H, et al. Low glutelin content1: A dominant mutation that suppresses the glutelin multigene family via RNA silencing in rice. *The Plant Cell*. 2003 Jun 1;15(6):1455-67.
36. Regina A, Bird A, Topping D, Bowden S, Freeman J, Barsby T, et al. High-amylose wheat generated by RNA interference improves indices of large-bowel health in rats. *Proceedings of the National Academy of Sciences*. 2006 Mar 7;103(10):3546-51.
37. Weise SE, Aung K, Jarou ZJ, Mehrshahi P, Li Z, Hardy AC, Carr DJ, Sharkey TD. Engineering starch accumulation by manipulation of phosphate metabolism of starch. *Plant Biotechnology Journal*. 2012 Jun;10(5):545-54.
38. Yu B, Lydiate DJ, Young LW, Schäfer UA, Hannoufa A. Enhancing the carotenoid content of *Brassica napus* seeds by downregulating lycopene epsilon cyclase. *Transgenic Research*. 2008 Aug;17:573-85.
39. Aggarwal S, Kumar A, Bhati KK, Kaur G, Shukla V, Tiwari S, Pandey AK. RNAi-mediated downregulation of inositol pentakisphosphate kinase (IPK1) in wheat grains decreases phytic acid levels and increases Fe and Zn accumulation. *Frontiers in Plant Science*. 2018 Mar 6;9:332156.
40. Escobar MA, Civerolo EL, Summerfelt KR, Dandekar AM. RNAi-mediated oncogene silencing confers resistance to crown gall tumorigenesis. *Proceedings of the National Academy of Sciences*. 2001 Nov 6;98(23):13437-42.
41. Katiyar-Agarwal S, Morgan R, Dahlbeck D, Borsani O, Villegas Jr A, Zhu JK, et al. A pathogen-inducible endogenous siRNA in plant immunity. *Proceedings of the National Academy of Sciences*. 2006 Nov 21;103(47):18002-7.
42. Enrique R, Siciliano F, Favaro MA, Gerhardt N, Roeschlin R, Rigano L, et al. Novel demonstration of RNAi in citrus reveals importance of citrus callose synthase in defence against *Xanthomonas citri* subsp. *citri*. *Plant Biotechnology Journal*. 2011 Apr;9(3):394-407.
43. Yu Z, Wang X, Mu X, Zhang L. RNAi mediated silencing of dehydrin gene WZY2 confers osmotic stress intolerance in transgenic wheat. *Functional Plant Biology*. 2019 Jun 14;46(10):877-84.
44. Wang W, Wang W, Wu Y, Li Q, Zhang G, Shi R, et al. The involvement of wheat U-box E3 ubiquitin ligase TaPUB1 in salt stress tolerance. *Journal of Integrative Plant Biology*. 2020 May;62(5):631-51.
45. Shi Y, Liu P, Xia Y, Wei P, Li W, Zhang W, et al. Downregulation of the lycopene ϵ -cyclase gene confers tolerance to salt and drought stress in *Nicotiana tabacum*. *Acta Physiologiae Plantarum*. 2015 Oct;37:1-5.
46. Cui P, Liu H, Islam F, Li L, Farooq MA, Ruan S, Zhou W. OsPEX11, a peroxisomal biogenesis factor 11, contributes to salt stress tolerance in *Oryza sativa*. *Frontiers in Plant Science*. 2016 Sep 15;7:208031.
47. Kim JA, Jung HE, Hong JK, Hermand V, Robertson McClung C, Lee YH, et al. Reduction of GIGANTEA expression in transgenic *Brassica rapa* enhances salt tolerance. *Plant Cell Reports*. 2016 Sep;35:1943-54.
48. Szechyńska-Hebda M, Czarnocka W, Hebda M, Karpiński S. PAD4, LSD1 and EDS1 regulate drought tolerance, plant biomass production, and cell wall

- properties. *Plant Cell Reports*. 2016 Mar;35:527-39.
49. Hu Y, Wu Q, Peng Z, Sprague SA, Wang W, Park J, et al. Silencing of OsGRXS17 in rice improves drought stress tolerance by modulating ROS accumulation and stomatal closure. *Scientific Reports*. 2017 Nov 21;7(1): 15950.
 50. Yin X, Huang L, Wang M, Cui Y, Xia X. OsDSR-1, a calmodulin-like gene, improves drought tolerance through scavenging of reactive oxygen species in rice (*Oryza sativa* L.). *Molecular Breeding*. 2017 Jun;37:1-3.
 51. Jin Y, Pan W, Zheng X, Cheng X, Liu M, Ma H, Ge X. OsERF101, an ERF family transcription factor, regulates drought stress response in reproductive tissues. *Plant Molecular Biology*. 2018 Sep;98:51-65.
 52. Zhu M, Meng X, Cai J, Li G, Dong T, Li Z. Basic leucine zipper transcription factor SlbZIP1 mediates salt and drought stress tolerance in tomato. *BMC Plant Biology*. 2018 Dec;18:1-4.
 53. Park JS, Yu JG, Lee GH, Park YD. Drought tolerance induction in transgenic tobacco through RNA interference of BrDST71, a drought-responsive gene from Chinese cabbage. *Horticulture, Environment, and Biotechnology*. 2018 Oct;59:749-57.
 54. Tzean Y, Chang HH, Tu TC, Hou BH, Chen HM, Chiu YS, et al. Engineering plant resistance to tomato yellow leaf curl Thailand virus using a phloem-specific promoter expressing hairpin RNA. *Molecular Plant-Microbe Interactions*. 2020 Jan 16;33(1):87-97.
 55. Gao L, Luo J, Ding X, Wang T, Hu T, Song P, et al. Soybean RNA interference lines silenced for eIF4E show broad potyvirus resistance. *Molecular Plant Pathology*. 2020 Mar;21(3): 303-17.
 56. Malathi P, Muzammil SA, Krishnaveni D, Balachandran SM, Mangrauthia SK. Coat protein 3 of Rice tungro spherical virus is the key target gene for development of RNAi mediated tungro disease resistance in rice. *Agri Gene*. 2019 Jun 1;12:100084.
 57. Sharma VK, Basu S, Chakraborty S. RNAi mediated broad-spectrum transgenic resistance in *Nicotiana benthamiana* to chilli-infecting begomoviruses. *Plant Cell Reports*. 2015 Aug;34:1389-99.
 58. Shoup Rupp JL, Cruz LF, Trick HN, Fellers JP. RNAi-Mediated, Stable Resistance to Triticum mosaic virus in Wheat. *Crop Science*. 2016 Jul;56(4):1602-10.
 59. Ahmed MM, Bian S, Wang M, Zhao J, Zhang B, Liu Q, et al. RNAi-mediated resistance to rice black-streaked dwarf virus in transgenic rice. *Transgenic Research*. 2017 Apr;26:197-207.
 60. Yang X, Niu L, Zhang W, Yang J, Xing G, He H, et al. RNAi-mediated SMV P3 cistron silencing confers significantly enhanced resistance to multiple Potyvirus strains and isolates in transgenic soybean. *Plant Cell Reports*. 2018 Jan;37:103-14.
 61. Kumari A, Hada A, Subramanyam K, Theboral J, Misra S, Ganapathi A, Malathi VG. RNAi-mediated resistance to yellow mosaic viruses in soybean targeting coat protein gene. *Acta Physiologiae Plantarum*. 2018 Feb;40:1-2.
 62. Senthilraja C, Gurivi Reddy M, Rajeswaran J, Kokiladevi E, Velazhahan R. RNA interference-mediated resistance to Tobacco streak virus in transgenic peanut. *Australasian Plant Pathology*. 2018 Mar;47:227-30.
 63. Hameed A, Tahir MN, Asad S, Bilal R, Van Eck J, Jander G, Mansoor S. RNAi-mediated simultaneous resistance against three RNA viruses in potato. *Molecular Biotechnology*. 2017 Mar;59:73-83.
 64. Shin YH, Lee SH, Park YD. Development of mite (*Tetranychus urticae*)-resistant transgenic Chinese cabbage using plant-mediated RNA interference. *Horticulture, Environment, and Biotechnology*. 2020 Apr;61:305-15.
 65. Dutta TK, Papolu PK, Banakar P, Choudhary D, Sirohi A, Rao U. Tomato transgenic plants expressing hairpin construct of a nematode protease gene conferred enhanced resistance to root-knot nematodes. *Frontiers in Microbiology*. 2015 Apr 1;6:260.
 66. Ibrahim AB, Monteiro TR, Cabral GB, Aragão FJ. RNAi-mediated resistance to whitefly (*Bemisia tabaci*) in genetically engineered lettuce (*Lactuca sativa*). *Transgenic Research*. 2017 Oct;26(5):613-24.
 67. Malik HJ, Raza A, Amin I, Scheffler JA, Scheffler BE, Brown JK, Mansoor S. RNAi-mediated mortality of the whitefly through transgenic expression of double-stranded RNA homologous to acetylcholinesterase and ecdysone receptor in tobacco plants. *Scientific Reports*. 2016 Dec 8;6(1):38469.

68. Sharma VK, Basu S, Chakraborty S. RNAi mediated broad-spectrum transgenic resistance in *Nicotiana benthamiana* to chilli-infecting begomoviruses. *Plant Cell Reports*. 2015 Aug;34:1389-99.
69. Mamta, Reddy KR, Rajam MV. Targeting chitinase gene of *Helicoverpa armigera* by host-induced RNA interference confers insect resistance in tobacco and tomato. *Plant Molecular Biology*. 2016 Feb;90:281-92.
70. Wang L, Wang H, He S, Meng F, Zhang C, Fan S, et al. GmSnRK1. 1, a sucrose non-fermenting-1 (SNF1)-related protein kinase, promotes soybean resistance to *Phytophthora sojae*. *Frontiers in Plant Science*. 2019 Aug 2;10:996.
71. Singh N, Mukherjee SK, Rajam MV. Silencing of the ornithine decarboxylase gene of *Fusarium oxysporum* f. sp. *lycopersici* by host-induced RNAi confers resistance to Fusarium wilt in tomato. *Plant Molecular Biology Reporter*. 2020 Sep;38:419-29.
72. Sanju S, Siddappa S, Thakur A, Shukla PK, Srivastava N, Pattanayak D, Sharma S, Singh BP. Host-mediated gene silencing of a single effector gene from the potato pathogen *Phytophthora infestans* imparts partial resistance to late blight disease. *Functional & Integrative Genomics*. 2015 Nov;15:697-706.
73. Cheng W, Song XS, Li HP, Cao LH, Sun K, Qiu XL, et al. Host-induced gene silencing of an essential chitin synthase gene confers durable resistance to Fusarium head blight and seedling blight in wheat. *Plant Biotechnology Journal*. 2015 Dec;13(9):1335-45.
74. Ghag SB, Shekhawat UK, Ganapathi TR. Host-induced post-transcriptional hairpin RNA-mediated gene silencing of vital fungal genes confers efficient resistance against Fusarium wilt in banana. *Plant Biotechnology Journal*. 2014 Jun;12(5): 541-53.
75. Andrade CM, Tinoco ML, Rieth AF, Maia FC, Aragão FJ. Host-induced gene silencing in the necrotrophic fungal pathogen *Sclerotinia sclerotiorum*. *Plant Pathology*. 2016 May;65(4):626-32.
76. Bharti P, Jyoti P, Kapoor P, Sharma V, Shanmugam V, Yadav SK. Host-induced silencing of pathogenicity genes enhances resistance to *Fusarium oxysporum* wilt in tomato. *Molecular Biotechnology*. 2017 Aug;59:343-52.
77. Gilbert MK, Majumdar R, Rajasekaran K, Chen ZY, Wei Q, Sickler CM, et al. RNA interference-based silencing of the alpha-amylase (*amy1*) gene in *Aspergillus flavus* decreases fungal growth and aflatoxin production in maize kernels. *Planta*. 2018 Jun;247:1465-73.
78. Majumdar R, Rajasekaran K, Sickler C, Lebar M, Musungu BM, Fakhoury AM, et al. The pathogenesis-related maize seed (PRms) gene plays a role in resistance to *Aspergillus flavus* infection and aflatoxin contamination. *Frontiers in Plant Science*. 2017 Oct 17;8:289597.
79. Yoshioka M, Adachi A, Sato Y, Doke N, Kondo T, Yoshioka H. RNAi of the sesquiterpene cyclase gene for phytoalexin production impairs pre-and post-invasive resistance to potato blight pathogens. *Molecular Plant Pathology*. 2019 Jul;20(7):907-22.
80. Pareek M, Rajam MV. RNAi-mediated silencing of MAP kinase signalling genes (*Fmk1*, *Hog1*, and *Pbs2*) in *Fusarium oxysporum* reduces pathogenesis on tomato plants. *Fungal Biology*. 2017 Sep 1;121(9):775-84.
81. Zhu L, Zhu J, Liu Z, Wang Z, Zhou C, Wang H. Host-induced gene silencing of rice blast fungus *Magnaporthe oryzae* pathogenicity genes mediated by the brome mosaic virus. *Genes*. 2017 Sep 26;8(10):241.
82. Zhu L, Zhu J, Liu Z, Wang Z, Zhou C, Wang H. Host-induced gene silencing of rice blast fungus *Magnaporthe oryzae* pathogenicity genes mediated by the brome mosaic virus. *Genes*. 2017 Sep 26;8(10):241.
83. Joshi I, Kumar A, Kohli D, Singh AK, Sirohi A, Subramaniam K, Chaudhury A, Jain PK. Conferring root-knot nematode resistance via host-delivered RNAi-mediated silencing of four *Mi-msp* genes in *Arabidopsis*. *Plant Science*. 2020 Sep 1;298:110592.
84. Dutta TK, Papolu PK, Banakar P, Choudhary D, Sirohi A, Rao U. Tomato transgenic plants expressing hairpin construct of a nematode protease gene conferred enhanced resistance to root-knot nematodes. *Frontiers in Microbiology*. 2015 Apr 1;6:260.
85. Li Y, Wang K, Lu Q, Du J, Wang Z, Wang D, et al. Transgenic *Nicotiana benthamiana* plants expressing a hairpin RNAi construct of a nematode *Rs-cps* gene exhibit enhanced resistance to

- Radopholus similis*. Scientific Reports. 2017 Oct 13;7(1):13126.
86. Chukwurah PN, Poku SA, Yokoyama A, Fukuda H, Shishido M, Nakamura I. Expression of *Meloidogyne incognita* PolA1 hairpin RNA reduced nematode multiplication in transgenic tomato. Plant Biotechnology Reports. 2019 Dec;13(6):591-601.
87. Hu Y, You J, Li C, Pan F, Wang C. The *Heterodera glycines* effector Hg16B09 is required for nematode parasitism and suppresses plant defense response. Plant Science. 2019 Dec 1;289:110271.
88. Tian B, Li J, Vodkin LO, Todd TC, Finer JJ, Trick HN. Host-derived gene silencing of parasite fitness genes improves resistance to soybean cyst nematodes in stable transgenic soybean. Theoretical and Applied Genetics. 2019 Sep 1;132:2651-62.
89. Nishihara M, Nakatsuka T, Yamamura S. Flavonoid components and flower color change in transgenic tobacco plants by suppression of chalcone isomerase gene. FEBS Letters. 2005 Nov 7;579(27):6074-8.
90. Kamiishi Y, Otani M, Takagi H, Han DS, Mori S, Tatsuzawa F, Okuhara H, Kobayashi H, Nakano M. Flower color alteration in the liliaceous ornamental *Tricyrtis* sp. by RNA interference-mediated suppression of the chalcone synthase gene. Molecular Breeding. 2012 Aug;30:671-80.
91. Nakatsuka T, Mishiba KI, Abe Y, Kubota A, Kakizaki Y, Yamamura S, Nishihara M. Flower color modification of gentian plants by RNAi-mediated gene silencing. Plant Biotechnology. 2008 Mar 1;25(1):61-8.
92. Nakatsuka T, Mishiba KI, Kubota A, Abe Y, Yamamura S, Nakamura N, Tanaka Y, Nishihara M. Genetic engineering of novel flower colour by suppression of anthocyanin modification genes in gentian. Journal of Plant Physiology. 2010 Feb 15;167(3):231-7.
93. He H, Ke H, Keting H, Qiaoyan X, Silan D. Flower colour modification of chrysanthemum by suppression of F3'H and overexpression of the exogenous *Senecio cruentus* F3'5'H gene. PLoS One. 2013 Nov 8;8(11):e74395.
94. Borgio JF. RNA interference (RNAi) technology: A promising tool for medicinal plant research. J Med Plant Res. 2009 Dec 1;3(13):1176-83.
95. Allen RS, Millgate AG, Chitty JA, Thisleton J, Miller JA, Fist AJ, Gerlach WL, Larkin PJ. RNAi-mediated replacement of morphine with the nonnarcotic alkaloid reticuline in opium poppy. Nature Biotechnology. 2004 Dec;22(12):1559-66.
96. Ogita S, Uefuji H, Yamaguchi Y, Koizumi N, Sano H. Producing decaffeinated coffee plants. Nature. 2003 Jun 19;423(6942):823.
97. Mohanpuria P, Kumar V, Ahuja PS, Yadav SK. Producing low-caffeine tea through post-transcriptional silencing of caffeine synthase mRNA. Plant Molecular Biology. 2011 Aug;76:523-34.
98. Zhang S, Li H, Liang X, Yan Y, Xia P, Jia Y, Liang Z. Enhanced production of phenolic acids in *Salvia miltiorrhiza* hairy root cultures by combing the RNAi-mediated silencing of chalcone synthase gene with salicylic acid treatment. Biochemical Engineering Journal. 2015 Nov 15;103:185-92.
99. Wang Q, Reddy VA, Panicker D, Mao HZ, Kumar N, Rajan C, Venkatesh PN, Chua NH, Sarojam R. Metabolic engineering of terpene biosynthesis in plants using a trichome-specific transcription factor Ms YABBY 5 from spearmint (*Mentha spicata*). Plant Biotechnology Journal. 2016 Jul;14(7):1619-32.
100. Jamaluddin ND, Rohani ER, Mohd Noor N, Goh HH. Transcriptome-wide effect of DE-ETIOLATED1 (DET1) suppression in embryogenic callus of *Carica papaya*. Journal of Plant Research. 2019 Mar 1;132:181-95.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/116796>