



# Assessment of Physico-Chemical Properties of Soil from Different Blocks of Eluru District, Andhra Pradesh, India

Chillumuntha Vishnu <sup>a\*</sup>, Ram Bharose <sup>a</sup>  
and Nithya Krishnan <sup>a</sup>

<sup>a</sup> Department of Soil Science and Agricultural Chemistry, Naini Agricultural Institute Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj- (211 007) Uttar Pradesh, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ijpss/2024/v36i74780>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/119063>

Original Research Article

Received: 11/04/2024

Accepted: 14/06/2024

Published: 26/06/2024

## ABSTRACT

The experiment was conducted in 2023 to assess the physical and chemical properties of different blocks of Eluru district, Andhra Pradesh. Three distinct villages were chosen from each block for the study. The physical properties of Eluru district soils, soil colour, texture, bulk density ( $\text{mg m}^{-3}$ ), particle density ( $\text{mg m}^{-3}$ ), percent pore space and percent water retaining capacity and chemical properties pH, EC ( $\text{dSm}^{-1}$ ) at 25°C, percent organic carbon, available nitrogen, phosphorus and potassium ( $\text{kg ha}^{-1}$ ), calcium C mol ( $\text{p}^+$ )  $\text{kg}^{-1}$  and magnesium C mol ( $\text{p}^+$ )  $\text{kg}^{-1}$  at 0-15, 15-30 and 30-45 cm depth. Findings showed that in terms of depth, there was a minor rise in both bulk and particle density. The water retaining capacity of the soil and the overall physical condition of the soil was

\*Corresponding author: E-mail: [vishnuch225@gmail.com](mailto:vishnuch225@gmail.com);

found good and supportive for plant growth. The soils of selected villages of different blocks in Eluru district were found to acidic to basic in nature due to soil minerals producing sodium carbonate ( $\text{Na}_2\text{CO}_3$ ) and sodium bicarbonate ( $\text{NaHCO}_3$ ) upon weathering. The organic carbon, nitrogen and phosphorus were observed as medium to high. The potassium content was low in all locations. There was enough calcium present, however there were several regions where magnesium was lacking. The deficiency of the nutrients can be mitigated by the use of organic and inorganic fertilizers.

**Keywords:** Eluru; physico-chemical properties of soil; Andhra Pradesh; weathering.

## 1. INTRODUCTION

Soil is an important element essential for the survival of living organisms [1]. Soil Testing is well recognized as an important technique to assess inherent nutrient supplying capacity of soil to the growing plants [2]. Andhra Pradesh is a state in the south-eastern coastal region of India. It is the seventh largest state by area covering an area of 162,975 km<sup>2</sup> (62,925 sq. mi) [3]. The city of Eluru lies at 16.7107°N 81.0952°E. The average rainfall is 119.02 mm yearly. The Soils in the district are made up of Alluvial, Black, Red and sandy soils. Paddy, banana, sugarcane, and coconut are the main agricultural products cultivated, and the Cashew nut, mango and tobacco are other important production from the district [4,5,6,7,8].

The study aims to evaluate the physico-chemical status of soils at both district and village levels. It seeks to provide a detailed analysis of soil properties, offering significant insights and guidelines to help farmers improve soil management practices and enhance agricultural productivity.

## 2. MATERIALS AND METHODS

Soil samples were collected from three different block i.e., Kaikaluru, Eluru and Denduluru of Eluru district. From each block soil samples were taken at a profile depth of 0-15, 15-30, 30-45 cm. A total 27 soil samples were collected from nine different villages. In each soil sampling bag 500 gm of soil was collected. Each soil samples were spread on a clean white paper sheet in the shade and air dried at room temperature. The large lumps/clods were crushed/broken to its ultimate soil particle using a wooden mallet. The powdered soil was sieved through 2 mm sieve. The soil material was collected in a clean cloth or polythene bag and was labelled properly for laboratory analysis. The physical properties of soils, soil colour, texture, bulk density ( $\text{mg m}^{-3}$ ), particle density ( $\text{mg m}^{-3}$ ), percent pore space and

percent water retaining capacity were analysed with the following standard procedure [9,10,11,] and chemical properties pH, EC ( $\text{dSm}^{-1}$ ) at 25°C, percent organic carbon, available nitrogen, phosphorus and potassium ( $\text{kg ha}^{-1}$ ), calcium C mol ( $\text{p}^+$ )  $\text{kg}^{-1}$  and magnesium C mol ( $\text{p}^+$ )  $\text{kg}^{-1}$  were analyzed by following [12,13,14,15,16,17] and [18] at 0-15, 15-30 and 30-45cm depth. The data recorded during the course of the investigation was completely randomized design, as per the method "Analysis of Variance technique" as given by [19].

## 3. RESULTS AND DISCUSSION

The results (Table 1) indicates that the soil colour of dry condition varies from greyish brown to yellowish red in 0-15 cm depth, from light brownish to red in 15-30 cm depth and from grey to red in 30-45 cm depth. The soil colour of wet condition varies from dark brown to dark red in 0-15 cm depth, from very dark to red in 15-30 cm depth and from very dark grey to red in 30-45 cm depth. In comparison to dry soil, wet soils are typically darker. The results (Table 2) indicates that the soil texture Eluru district is dominantly found as sandy loam, clay and sandy clay. The sand, silt and clay percentage varied from 25.14% to 67.12%, 16.50% to 23.15% and 11.49% to 54.21%. Comparable outcomes were discovered by [20]. The results (Table 3) indicates that the bulk density of soil varies from 1.06 to 1.33  $\text{mg m}^{-3}$  in selected blocks of Eluru district, Andhra Pradesh. It is clear that the bulk density increases with the increase in soil depth due to soil compactness. Comparable outcomes were discovered by [21], and the particle density of soil varied from 2.15 to 2.59  $\text{mg m}^{-3}$ . According to depth, the particle density likewise rises. Particle density varies according to mineral content of soil particles. Comparable outcomes were discovered by [22]. The findings (Table 4) indicate that the pore space (%) of soil ranged from 41.56 to 50.27 %. The pore space found to decrease with increase in depth at attributed to increase in compaction in the sub surface.

**Table 1. Soil colour in dry and wet condition of different block of Eluru district, Andhra Pradesh**

Block	Village	Dry condition			Wet condition		
		0-15 cm depth	15-30 cm depth	30-45 cm depth	0-15 cm depth	15-30 cm depth	30-45 cm depth
Kaikaluru	Varahapatnam	10YR 5/2 greyish brown	10YR 6/2 light brownish grey	10YR 5/1 grey	7.5YR 4/2 dark brown	10YR 3/0 very dark grey	10YR 3/1 very dark grey
	Tamarakollu	10YR 5/1 grey	10YR 7/1 light grey	5Y 5/2 olive grey	5YR 4/1 dark grey	5YR 4/1 dark grey	5Y 3/2 dark olive grey
	Achavaram	10YR 6/2 light brownish grey	10YR 7/1 light grey	10YR 6/1 grey	2.5YR 4/4 olive brown	10YR 3/3 dark brown	10YR 4/2 dark greyish brown
Eluru	Sreeparru	2.5Y 4/2 dark greyish brown	2.5YR 5/4 light olive brown	2.5Y 5/6 light olive brown	2.5YR 4/2 dark greyish brown	2.5Y 4/2 dark greyish brown	2.5Y 4/2 dark greyish brown
	Madepalli	10YR 7/2 light grey	5Y 5/6 olive	5Y 6/6 olive yellow	5Y 4/3 olive	5Y 4/4 olive	5Y 5/3 olive
	Malkapuram	10YR 6/3 pale brown	2.5Y 4/4 olive brown	5Y 5/6 olive	5Y 4/4 olive	2.5Y 4/2 dark greyish brown	5Y 4/2 olive grey
Denduluru	Denduluru	7.5YR 4/4 dark brown	2.5Y 6/4 light yellowish brown	10YR 6/8 brownish yellow	2.5Y 5/6 light olive brown	2.5Y 5/6 light olive brown	10YR 5/8 yellowish brown
	Challa chintalapudi	5YR 3/3 dark reddish brown	2.5YR 3/6 dark red	10R 4/6 red	5YR 4/3 reddish brown	2.5YR 4/4 reddish brown	10R 3/6 dark red
	Ramarao gudem	5YR 4/6 yellowish red	2.5YR 4/6 red	2.5YR 5/8 red	2.5YR 3/6 dark red	2.5YR 4/8 red	2.5YR 4/8 red

**Table 2. Soil texture of different block of Eluru district, Andhra Pradesh at 0-15, 15-30 and 30-45cm**

S. No.	Block	Village	Sand %	Silt %	Clay %	Textural Class
1	Kaikaluru	Varahapatnam B <sub>1</sub> V <sub>1</sub>	67.12	20.08	12.80	Sandy Loam
		Tamarakollu B <sub>1</sub> V <sub>2</sub>	64.48	21.36	14.16	Sandy Loam
		Achavaram B <sub>1</sub> V <sub>3</sub>	65.36	23.15	11.49	Sandy Loam
2	Eluru	Sreeparru B <sub>2</sub> V <sub>1</sub>	28.05	18.27	53.68	Clay
		Madepalli B <sub>2</sub> V <sub>2</sub>	25.14	20.65	54.21	Clay
		Malkapuram B <sub>2</sub> V <sub>3</sub>	30.36	17.53	52.11	Clay
3	Denduluru	Denduluru B <sub>3</sub> V <sub>1</sub>	46.15	17.04	36.81	Sandy Clay
		Challa chintalapudi B <sub>3</sub> V <sub>2</sub>	48.38	16.50	35.12	Sandy Clay
		Ramarao gudem B <sub>3</sub> V <sub>3</sub>	45.10	19.34	35.56	Sandy Clay

**Table 3. Bulk density ( $\text{mg m}^{-3}$ ) and Particle Density ( $\text{mg m}^{-3}$ ) of soil at different block of Eluru district, Andhra Pradesh**

Block	Village	Bulk Density ( $\text{mg m}^{-3}$ )		Particle Density ( $\text{mg m}^{-3}$ )	
		Range	Mean	Range	Mean
Kaikaluru	Varahapatnam B <sub>1</sub> V <sub>1</sub>	1.27 – 1.31	1.29	2.53 – 2.59	2.56
	Tamarakollu B <sub>1</sub> V <sub>2</sub>	1.25 – 1.29	1.27	2.46 – 2.50	2.48
	Achavaram B <sub>1</sub> V <sub>3</sub>	1.28 – 1.30	1.29	2.43 – 2.47	2.44
Eluru	Sreeparru B <sub>2</sub> V <sub>1</sub>	1.17 – 1.22	1.19	2.50 – 2.56	2.52
	Madepalli B <sub>2</sub> V <sub>2</sub>	1.25 – 1.29	1.26	2.50 – 2.53	2.51
	Malkapuram B <sub>2</sub> V <sub>3</sub>	1.30 - 1.33	1.31	2.35 – 2.39	2.37
Denduluru	Denduluru B <sub>3</sub> V <sub>1</sub>	1.17 – 1.22	1.19	2.22 – 2.24	2.23
	Challa chintalapudi B <sub>3</sub> V <sub>2</sub>	1.14 – 1.19	1.16	2.19 – 2.27	2.23
	Ramarao gudem B <sub>3</sub> V <sub>3</sub>	1.06 – 1.12	1.09	2.15 – 2.23	2.19

**Table 4. Pore Space (%) and Water Retaining Capacity (%) of soil at different block of Eluru district, Andhra Pradesh**

Block	Village	Pore space (%)		Water Retaining Capacity (%)	
		Range	Mean	Range	Mean
Kaikaluru	Varahapatnam B <sub>1</sub> V <sub>1</sub>	44.61 – 49.68	47.05	43.82 – 48.66	45.59
	Tamarakollu B <sub>1</sub> V <sub>2</sub>	45.29 – 49.37	47.39	42.81 – 47.54	45.16
	Achavaram B <sub>1</sub> V <sub>3</sub>	43.92 – 47.43	45.37	41.14 – 45.67	43.38
Eluru	Sreeparru B <sub>2</sub> V <sub>1</sub>	45.11 – 50.27	47.73	43.97 – 49.85	46.27
	Madepalli B <sub>2</sub> V <sub>2</sub>	44.32 – 48.78	46.93	41.71 – 46.69	44.68
	Malkapuram B <sub>2</sub> V <sub>3</sub>	41.56 – 46.66	44.36	40.36 – 45.62	42.85
Denduluru	Denduluru B <sub>3</sub> V <sub>1</sub>	42.67 – 47.05	44.74	41.50 – 46.70	43.78
	Challa chintalapudi B <sub>3</sub> V <sub>2</sub>	43.29 – 48.57	45.80	40.33 – 47.96	44.36
	Ramarao gudem B <sub>3</sub> V <sub>3</sub>	45.59 – 50.26	48.19	44.11 – 49.60	46.89

**Table 5. Soil pH of soil at different block of Eluru district, Andhra Pradesh**

Block	Village	Soil pH	
		Range	Mean
Kaikaluru	Varahapatnam B <sub>1</sub> V <sub>1</sub>	6.98 – 7.20	7.08
	Tamarakollu B <sub>1</sub> V <sub>2</sub>	5.87 – 6.38	6.05
	Achavaram B <sub>1</sub> V <sub>3</sub>	6.70 – 7.00	6.80
Eluru	Sreeparru B <sub>2</sub> V <sub>1</sub>	7.54 – 7.93	7.69
	Madepalli B <sub>2</sub> V <sub>2</sub>	6.18 – 8.05	7.35
	Malkapuram B <sub>2</sub> V <sub>3</sub>	7.59 – 7.93	7.80
Denduluru	Denduluru B <sub>3</sub> V <sub>1</sub>	7.45 – 7.71	7.55
	Challa chintalapudi B <sub>3</sub> V <sub>2</sub>	7.36 – 7.61	7.51
	Ramarao gudem B <sub>3</sub> V <sub>3</sub>	6.10 – 6.83	6.40

**Table 6. EC (dS m<sup>-1</sup>) and Organic Carbon (%) of soil at different block of Eluru district, Andhra Pradesh**


Block	Village	EC of soil water suspension (dS m <sup>-1</sup> )		Organic Carbon (%)	
		Range	Mean	Range	Mean
Kaikaluru	Varahapatnam B <sub>1</sub> V <sub>1</sub>	0.46 – 0.58	0.50	0.52 – 1.60	1.01
	Tamarakollu B <sub>1</sub> V <sub>2</sub>	0.18 – 0.28	0.21	0.65 – 0.75	0.68
	Achavaram B <sub>1</sub> V <sub>3</sub>	0.32 – 0.87	0.52	0.30 – 0.52	0.40
Eluru	Sreeparru B <sub>2</sub> V <sub>1</sub>	0.57 – 0.73	0.67	0.49 – 0.52	0.51
	Madepalli B <sub>2</sub> V <sub>2</sub>	0.44 – 0.58	0.51	0.49 – 0.52	0.50
	Malkapuram B <sub>2</sub> V <sub>3</sub>	0.75 – 0.80	0.77	0.65 – 1.17	0.82
Denduluru	Denduluru B <sub>3</sub> V <sub>1</sub>	0.14 – 0.25	0.18	0.52 – 0.78	0.60
	Challa chintalapudi B <sub>3</sub> V <sub>2</sub>	0.21 – 0.23	0.21	0.39 – 0.60	0.50
	Ramarao gudem B <sub>3</sub> V <sub>3</sub>	0.19 – 0.35	0.25	0.52 – 0.91	0.69

**Table 7. Available Nitrogen (kg ha<sup>-1</sup>), Phosphorus (kg ha<sup>-1</sup>) and Potassium (kg ha<sup>-1</sup>) of soil at different block of Eluru district, Andhra Pradesh**


Block	Village	Available Nitrogen (kg ha <sup>-1</sup> )		Available Phosphorus (kg ha <sup>-1</sup> )		Available Potassium (kg ha <sup>-1</sup> )	
		Range	Mean	Range	Mean	Range	Mean
Kaikaluru	Varahapatnam B <sub>1</sub> V <sub>1</sub>	250.0 – 325.0	283.3	70.0 – 94.0	84.6	26.0 – 31.0	28.3
	Tamarakollu B <sub>1</sub> V <sub>2</sub>	175.0 – 300.0	250.0	45.0 – 94.0	73.6	14.0 – 21.0	16.6
	Achavaram B <sub>1</sub> V <sub>3</sub>	250.0 – 350.0	283.3	35.0 – 62.0	50.3	27.0 – 30.0	28.6
Eluru	Sreeparru B <sub>2</sub> V <sub>1</sub>	120.0 – 250.0	203.3	37.0 – 83.0	57.0	64.0 – 77.0	69.0
	Madepalli B <sub>2</sub> V <sub>2</sub>	156.0 – 376.0	294.0	51.0 – 83.0	67.0	54.0 – 65.0	58.0
	Malkapuram B <sub>2</sub> V <sub>3</sub>	200.0 – 275.0	241.6	22.0 – 69.0	48.3	58.0 – 61.0	60.0
Denduluru	Denduluru B <sub>3</sub> V <sub>1</sub>	200.0 – 250.0	216.6	26.0 – 53.0	39.3	30.0 – 37.0	34.0
	Challa chintalapudi B <sub>3</sub> V <sub>2</sub>	200.0 – 300.0	233.3	29.0 – 52.0	37.6	20.0 – 34.0	26.0
	Ramarao gudem B <sub>3</sub> V <sub>3</sub>	200.0 – 250.0	216.6	37.0 – 86.0	60.3	40.0 – 49.0	43.0

**Table 8. Exchangeable Calcium and Magnesium (meq 100g<sup>-1</sup>) of soil at different block of Eluru district, Andhra Pradesh**

Block	Village	Exchangeable Calcium (meq 100g <sup>-1</sup> )		Exchangeable Magnesium (meq 100g <sup>-1</sup> )	
		Range	Mean	Range	Mean
Kaikaluru	Varahapatnam B <sub>1</sub> V <sub>1</sub>	6.90 – 7.80	7.30	0.20 – 0.30	0.26
	Tamarakollu B <sub>1</sub> V <sub>2</sub>	1.90 – 2.90	2.43	0.50 – 0.90	0.70
	Achavaram B <sub>1</sub> V <sub>3</sub>	2.00 – 3.40	2.86	0.60 – 1.00	0.80
Eluru	Sreeparru B <sub>2</sub> V <sub>1</sub>	3.51 – 5.18	4.26	0.80 – 1.30	1.06
	Madepalli B <sub>2</sub> V <sub>2</sub>	2.36 – 5.02	3.78	1.38 – 2.47	1.98
	Malkapuram B <sub>2</sub> V <sub>3</sub>	2.11 – 4.99	3.64	1.20 – 2.60	2.08
Denduluru	Denduluru B <sub>3</sub> V <sub>1</sub>	6.40 – 8.10	7.43	0.60 – 0.90	0.76
	Challa chintalapudi B <sub>3</sub> V <sub>2</sub>	8.10 – 9.60	8.86	2.70 – 3.90	3.26
	Ramarao gudem B <sub>3</sub> V <sub>3</sub>	4.90 – 6.90	5.73	0.20 – 0.30	0.23




**Fig. 1. The soil of textural triangle at different blocks of Eluru district**




**Fig. 2. Graphical representation of bulk density (mg m<sup>-3</sup>) and particle density (mg m<sup>-3</sup>) of soil in different block of Eluru district, Andhra Pradesh**

Surface soils are having high amount of macro and micro pores compared to sub surface soil due to presence of high organic matter. Comparable outcomes were discovered by [22], and the water retaining capacity of soil (%) ranged from 40.33 to 49.85 %. The water retaining capacity value decrease with increase in depth because of soil compaction and reduction in pore space. Soils vary in their water holding capacity according to their structure, texture and bulk density relationship to total pore size distribution. Comparable outcomes were discovered by [23]. The findings (Table 5)

indicate that the pH value of soil ranged from 5.87 to 8.05. The soils of selected villages of different blocks in Eluru district were found to acidic to basic in nature. The causes of soil alkalinity can be natural or man-made: The natural cause is the presence of soil minerals producing sodium carbonate (Na<sub>2</sub>CO<sub>3</sub>) and sodium bicarbonate (NaHCO<sub>3</sub>) upon weathering. Comparable outcomes were discovered by [24]. The findings (Table 6) indicate that the electrical conductivity of soil ranged from 0.14 to 0.87 dS m<sup>-1</sup>.



**Fig. 3. Graphical representation of soil Pore Space (%) and Water Retaining Capacity (%) in different block of Eluru district, Andhra Pradesh**




**Fig. 4. Graphical representation of Soil pH in different block of Eluru district, Andhra Pradesh**


The soil samples are clearly acceptable for most crops and satisfy the permitted EC level. Comparable outcomes were discovered by [20], and the value of total organic carbon (%) of soil varied from 0.30 to 1.60 %. The block that was chosen has a medium level of organic matter because to microbial activity and the deposition of plant residue. Comparable outcomes were discovered by [25]. The findings (Table 7) indicate that the available nitrogen content of soil

ranged from 120.0 to 350.0 kg ha<sup>-1</sup>. The district of Eluru villages all have medium nitrogen levels. Available nitrogen decreases with the increasing depth due to the fact it is positively correlated with organic matter content which decreases with depth and might be due to higher pH to depth. Comparable outcomes were discovered by [26]. The findings (Table 7) indicate that the available phosphorus content of soil ranged from 22.0 to 94.0 kg ha<sup>-1</sup>.






**Fig. 5. Graphical representation of EC ( $\text{dS m}^{-1}$ ) and Organic Carbon (%) different block of Eluru district, Andhra Pradesh**



**Fig. 6. Graphical representation of available Nitrogen ( $\text{kg ha}^{-1}$ ), Phosphorus ( $\text{kg ha}^{-1}$ ) and Potassium ( $\text{kg ha}^{-1}$ ) of Soil in different block of Eluru district, Andhra Pradesh**

Available phosphorus decreases with the increasing depth. Higher level of available phosphorus in surface soil could be attribute of favorable soil pH and organic matter content. Comparable outcomes were discovered by [27]. The findings (Table 7) indicate that available potassium content of soil ranged from 14.0 to 77.0  $\text{kg ha}^{-1}$ . Available potassium decreases with the increasing depth. The high content of available potassium on surface soil may be attributed to the release of available K form

organic residues and application of potassium fertilizers. Comparable outcomes were discovered by [27]. The findings (Table 8) indicate that exchangeable calcium and magnesium content of soil ranged from 1.90 to 9.60 ( $\text{cmol (p}^+) \text{ kg}^{-1}$ ), 0.20 to 3.90 ( $\text{cmol (p}^+) \text{ kg}^{-1}$ ) respectively in selected blocks. The calcium and magnesium decrease with the increasing in depth due to the attributes of high pH towards the depth. Comparable outcomes were discovered by [28].



**Fig. 7. Graphical representation of Exchangeable Calcium and Magnesium (meq100g<sup>-1</sup>) of soil in different block of Eluru district, Andhra Pradesh**

#### 4. CONCLUSION

From the above study the relationship between various physical and chemical parameters and their effect on soil quality was analysed. The soils of selected villages of different blocks in Eluru district were found to acidic to basic in nature. The soil colour of dry condition varies from greyish brown to yellowish red in 0-15 cm depth, from light brownish to red in 15-30 cm depth and from grey to red in 30-45 cm depth. The soil colour of wet condition varies from dark brown to dark red in 0-15 cm depth, from very dark to red in 15-30 cm depth and from very dark grey to red in 30-45 cm depth. The soil textural class was identified as sandy loam, clay and sandy clay. The bulk density and particle density increase with the increase in soil depth due to soil compactness. The water retaining capacity of the soil and the overall physical condition of the soil was found good and supportive for plant growth. It is clear that the soils of selected villages of three blocks are acidic to basic in nature. The electrical conductivity of the soil ranged from 0.14 to 0.87 which is suitable for crop growth. The organic carbon content were observed as medium. Nitrogen and phosphorous content of the soil was medium to high. The potassium content were observed as low. Calcium and magnesium are sufficient in this soil. The deficiency of the nutrients can be

mitigated by the use of organic and inorganic fertilizers.

#### 5. FUTURE PERSPECTIVES OF THE STUDY

Future studies should focus on monitoring soil changes over time and seasonal variations to understand the impact of agricultural practices and climate change. High-resolution soil mapping and spatial distribution analysis can identify areas needing targeted interventions. Evaluating different agricultural practices and soil conservation techniques will help recommend best practices. Precision agriculture and organic amendments can improve nutrient management and soil fertility. Research on soil-plant interactions, crop diversification, and resilience to climate change will enhance soil and crop productivity. Identifying soil pollutants and developing remediation strategies are crucial for maintaining soil health. Engaging farmers through training and participatory research ensures practical implementation, while policy recommendations and incentive programs support sustainable soil management.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image

generators have been used during writing or editing of manuscripts.

## ACKNOWLEDGMENT

I express my gratitude to my advisor Dr. Ram Bharose, for his guidance, support and expertise the Department of Soil Science and Agricultural Chemistry, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, U.P. and Krishi Vigyan Kendra(KVK), Undi, West Godavari, Andhra Pradesh for providing necessary equipment, laboratory for analysis and other basic infrastructure.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Abo-Habaga MM, Ismail ZE, Okasha MH. Effect of tillage systems on a soil moisture content and crops productivity. *Journal of Soil Sciences and Agricultural Engineering*. 2022;13(7):231-235.
2. Bhatt R, Sharma M. Importance of soil testing and techniques of soil sampling. Lap Lambert Academic publishing. 2014
3. Sahu, Ranjith Kumar, Tarence Thomas, Anurag Kumar Singh, and Indar Raj Naga. Assessment of Physico-Chemical Properties of Soil from Different Blocks of Visakhapatnam District, Andhra Pradesh. *International Journal of Plant & Soil Science*. 2023;35(15):322-335.
4. District Administration West Godavari. Developed and hosted by national informatics centre, ministry of electronics & information technology, Government of India; 2024.
5. Kumari S, Swaroop N, Thomas T, Kumar R. Assessment of Physico-Chemical Properties of Soil from Different Blocks of Sahibganj District, Jharkhand, India. *Int. J. Environ. Clim. Change*. 2023 Jun. 20 [cited 2024 Jun. 13];13(8):1998-2004. Available: <https://journalijecc.com/index.php/IJECC/article/view/2157>
6. Raina TK, Kumar Salgotra R, Dey T, Singh B, Sharma S, Johar P. Assessment of Physicochemical Properties of Rhizosphere Soil Samples from Basmati and Non-basmati Rice Growing Areas of Jammu Region, Jammu and Kashmir, India. *J. Exp. Agric. Int.* 2021 Dec. 21 [cited 2024 Jun. 13];43(11):200-9. Available: <https://journaljeai.com/index.php/JEAI/article/view/1912>
7. Singh S, Singh J, Vig AP. Earthworm as ecological engineers to change the physico-chemical properties of soil: soil vs vermicast. *Ecological Engineering*. 2016 May 1;90:1-5.
8. Berihun T, Tadele M, Kebede F. The application of biochar on soil acidity and other physico-chemical properties of soils in southern Ethiopia. *Journal of plant nutrition and soil Science*. 2017 Jun; 180(3):381-8.
9. Munsell AH. *Munsell Soil Colour Charts*. Munsell Colour Company Inc., Baltimore; 1954.
10. Bouyoucos GJ. The hydrometer as new method of mechanical analysis of soil. *Soil Science*. 1927;23:343-353.
11. Muthuvel P, Udayasoorian C, Natesan R, Ramaswami PR. *Introduction to soil analysis*. Coimbatore: Tamil Nadu Agricultural University; 1992.
12. Jackson ML. The pH was determined in 1:2 soil water suspensions using digital pH meter. *Soil Chemical Analysis*, Prentice Hall, Inc. Englewood. cliffe. NJ.; 1958.
13. Wilcox et al. Electrical conductivity, Amer. Water Works Assoc. J., 1950;42:775-776.
14. Walkley A, Black. A critical examination of a Rapid method for determining organic carbon in soils: Effect of variations in digestion conditions and of Organic soil constituents. *Soil Sci*. 1947;63:251- 263.
15. Subbiah BV, Asija GL. A Rapid Procedure for the Determination of Available Nitrogen in Soils. *Current Science*, Vol. 1956;25: 259-260.
16. Olsen SR, Cole CV, Watanabe. F.S. And Dean. LA. Estimation of Available Phosphorus in Soils by Extraction with Sodium Bicarbonate. U. S. Department of Agriculture, Circular No. 1954;939.
17. Toth SJ, rince AL. Estimate of Cation Exchange capacity and exchangeable Ca, K, Na, Content of soil by flame photometer technique. *Soil Sci*. 1949;67:439-445.
18. Jackson ML. *Soil Chemical Analysis*. Prentice Hall of India Pvt. Ltd., New Delhi; 1973
19. Fisher RA. Statistical methods and scientific induction. *Journal of the royal statistical society series*. 1960;17:69-78.
20. Sathyanarayana E, Padmaja G, Saranya S, Bharghavi J, Santhosh Kumar M, Rajashekhar M, Veeranna J, Kumari Sunita Soil fertility status of

- soybean growing soils of Adilabad district, Telangana, The Pharma Innovation Journal. 2021;SP-10(10):1112-1120.
21. Gangothri N, Dadhich AS. A study on the availability of micro and macro nutrients in red and black agricultural soils of Eklaskhampeta, Telangana, India, Poll Res. 2021;40(May Suppl. Issue):104-111.
  22. Verma C, Lal A, David ADM, Rao PS. Determination of Physico-chemical properties in soil samples of Prayagraj (Allahabad) District, Uttar Pradesh, India. Asian Journal of Applied Chemistry Research. 2019;4(2):1-8.
  23. Singh AK, Singh AK, Singh AK. Study of available potassium by different extracting reagents and fertility status of soils of Bairia sub-division of Ballia UP. Asian Journal of Science and Technology. 2019;9(4):9584-9586.
  24. Kumari AK, Rao PC, Padmaja G, Madhavi M. Effect of Physico- Chemical Properties on Soil Enzyme Acid Phosphatase Activity of Some Soils in Vegetable Growing research station, Warangal, Telangana, Journal of Pharmacognosy and Phytochemistry. 2017;7(6):1852-1856.
  25. Singh YP, Raghubanshi BPS, Tiwari RJ, Motsara S. Distribution of Available Macro and Micronutrients in Soils of Morena District of Madhya Pradesh, Journal of Multidisciplinary Advance Research. 2014;3:1-8.
  26. Rajamani K, Hari N, Rajashekar M. Soil Fertility Evaluation and GPS- GIS Based Soil Nutrient Mapping of Krishi Vigyan Kendra, Palem, Telangana, International Research Journal of Pure & Applied Chemistry. 2020;21(23):139-145. ISSN: 2231-3443.
  27. Wani SA, Najjar GR, Padder BA, Akhter F, Chand S. Altitudinal and depth-wise variation of soil physico-chemical properties and available nutrients of pear orchards in Jammu & Kashmir, India. Chem. Sci. Rev. Lett. 2017;6(23):1638-1645.
  28. Malavath R, Mahesh C, Balaguruvaiah D, Vidyasagar GECH. Land use options and site suitability for sugarcane growing red soils, red laterite soils and black soils of Medak district of Telangana. Journal of Pharmacognosy and Phytochemistry. 2018;7(2):409-416.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/119063>