



Advances in Research

Volume 25, Issue 4, Page 207-218, 2024; Article no.AIR.119197
ISSN: 2348-0394, NLM ID: 101666096

Mental Health and Quality Sleep among Brazilian Medical Students during Covid-19 Pandemic: Prevalence and Associated Factors

Gabriel S. Mendonça ^a, Gabriel R. Medeiros ^a,
Gilsiane S. Sampaio ^a, Eduarda R. Rabello ^a,
Daniela R. Freitas ^a, Gabriella M. L. Oliveira ^a
and Paulo M. A. Lima ^{a*}

^a Núcleo de Pesquisa da Faculdade de Medicina, Universidade de Rio Verde – Campus Goiânia, Goiânia, Goiás, Brazil.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/air/2024/v25i41098>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/119197>

Original Research Article

Received: 20/04/2024
Accepted: 24/06/2024
Published: 28/06/2024

ABSTRACT

Aims: Due the pandemic of the new coronavirus (COVID-19), stricter measures to limit unessential interactions were adopted, including the suspension of classroom lessons and the implementation of the distance learning regime. As these restrictions change lifestyles, this study aimed to assess

*Corresponding author: E-mail: paulo.lima@unirv.edu.br, lima.pma@outlook.com;

Cite as: Mendonça, Gabriel S., Gabriel R. Medeiros, Gilsiane S. Sampaio, Eduarda R. Rabello, Daniela R. Freitas, Gabriella M. L. Oliveira, and Paulo M. A. Lima. 2024. "Mental Health and Quality Sleep Among Brazilian Medical Students During Covid-19 Pandemic: Prevalence and Associated Factors". *Advances in Research* 25 (4):207-18. <https://doi.org/10.9734/air/2024/v25i41098>.

the impacts of social distancing on mental health and sleep quality in medical students during the COVID-19 pandemic.

Study Design: Cross-sectional study.

Place and Duration of Study: Medical school in Goiânia, Goiás, Brazil. Data collection took place from June 1 to July 1, 2020.

Methodology: This study used self-administered questionnaires answered by students from a Public University in Brazil. The collection instrument consisted of four parts: socioeconomic variables, self-perception of health, Self-Report Questionnaire (SRQ-20), and Pittsburgh Sleep Quality Index (PSQI). Absolute and relative frequencies described the sample, Chi-square test and Student's t-test tested associations between the variables, Pearson's correlation tested the relationship between PSQI and SRQ-20.

Results: Of the 210 participants, 138 students (65.7%) had scores suggestive of mental distress by the SRQ-20; women had a higher risk when compared to men (women 67.7% vs. men 49%, $p < .01$). 186 students (88.6%) had poor sleep quality according to the PSQI; women also had higher sleep disturbance (women 1.26 ± 0.06 vs. men 1.35 ± 0.04 , $p < .05$). Mental distress measures were positively correlated with sleep quality ($r = 0.611$, $p < .001$).

Conclusion: The strict preventive measures, and the radical changes in the lifestyle of medical students had a negative impact on the quality of their sleep and mental state. The students showed high levels of mental distress and poor sleep quality, and women appear to be an important risk factor. Implementation of interventions for well-being in pandemic condition by universities, and special focus to the most affected groups (such as women), should be stimulated.

Keywords: Coronavirus; medicine; quarantine; PSQI; SRQ-20.

1. INTRODUCTION

Coronavirus Disease 19 (COVID-19) pandemic, caused by the new coronavirus related to the severe acute respiratory syndrome (SARS-CoV-2), has resulted in an ongoing global public health crisis [1]. First detected in Wuhan, China, in December 2019, the World Health Organization declared the COVID-19 outbreak a public health emergency of international concern on January 30, 2020, and officially characterized it as a pandemic on March 11, 2020 [2].

SARS-CoV-2 is highly transmissible, spreading primarily through direct contact or respiratory droplets from an infected individual [1]. Common symptoms include a runny nose, cough, sore throat, body aches, nausea, vomiting, diarrhea, fever, and breathing difficulties [3]. To contain further spread and prevent healthcare system collapse, many governments implemented stringent measures such as social distancing, movement restrictions, curfews, mask mandates, contact tracing, virus detection tests, and quarantine [4]. On 15th March 2020, Brazil declared a strict nationwide quarantine.

While these measures limited non-essential interactions, movement, and work activities to control disease transmission, they also increased risk factors for non-communicable diseases [5].

These restrictions significantly altered lifestyles and social relationship, leading to increased psychological distress and mental health issues, including sadness, fear, anger, anxiety, depression, and sleep disorders [6].

University students, particularly medical students, were significantly affected by these measures. Most universities worldwide suspended face-to-face learning, transitioning to remote classes and exams [7]. For medical students, clinical practice and laboratory exercises were paused, causing major disruptions in their education and daily life [8].

Medical students are particularly vulnerable to poor mental well-being and sleep quality, experiencing higher rates of mental illness and psychological distress compared to the general student population [9-11]. Contributing factors include adjustment to the medical school environment, ethical conflicts, and exposure to death, human suffering [12,13].

Studies examining the impact of COVID-19 on the quality of life of the Brazilian population remain scarce. Therefore, this study aimed to assess mental health distress and sleep quality among medical students at a public University in Goiás, Brazil, and to identify sociodemographic variables associated with psychological distress.

2. MATERIALS AND METHODS

2.1 Study Design and Participants

This cross-sectional study aimed to evaluate the mental health and sleep quality among medical student during the COVID-19 pandemic and identify associated factors.

The participants were undergraduate students enrolled at a medical school in Goiânia, Goiás, Brazil. In Brazil, medical school consists of 12 periods (semesters), classified as follows: from the 1st to the 4th (Basic Course) and from the 5th semester onwards (Professional Cycle). All students from the first to the fourth year who were 18 years or older were invited to participate. The students were contacted via their mobile numbers due to the strict social distancing measures during the COVID-19 pandemic. Informed consent and the questionnaires were provided online through the Google Forms platform and could be accessed via a link sent to all participants. The students first accessed the informed consent form on the initial screen of the electronic questionnaire and could only proceed after consenting. The questionnaires were available only in Portuguese. Participation was entirely voluntary, with no incentives provided, and individuals were not identifiable in the research.

The COVID-19 pandemic was officially declared in Brazil on 15th March of 2020; the campus was closed, and most of the curriculum was offered remotely. Data collection took place from June 1 to July 1, 2020, immediately after the first wave of COVID-19 pandemic in Brazil.

The sample size calculation was based on the population of approximately 480 students enrolled from the first to the fourth year. With a margin of error of 5% and a confidence interval of 95%, it was determined that 218 volunteers needed to be interviewed.

The experimental protocol was approved by the National Research Ethics Commission of Brazil Platform (CAAE 32563220.0.0000.0008).

2.2 Data collection and questionnaires

The participants answered four questionnaires: sociodemographic information, self-perception of health (SPH), the self-reporting questionnaire (SRQ-20), and the Pittsburgh sleep quality index (PSQI).

Sociodemographic information. Adapted from the Brazilian Institute of Geography and Statistics, this section included questions about gender, age, skin color, religion, marital status, family income, academic level, and housing information.

Self-perception of health (SPH). Developed by the researchers, this questionnaire asked students to assess their own health, report any pre-existing disease or disorders that impair their well-being, and indicate if they were currently receiving treatments or support for a mental health concern.

Self-reporting questionnaire (SRQ-20). Mental distress was evaluated using the Brazilian validated version of SRQ-20 [14], a questionnaire developed by the World Health Organization to investigate non-psychotic psychiatric disorders. Originally, SRQ was constructed to screen psychotic (4 items) and non-psychotic (20 items) disorders; because of the low sensitivity and unreliability of the 4 psychotic items, the 20 items are widely used in epidemiologic studies, hence referred as SRQ-20 [15]. SRQ-20 aims to evaluate symptoms and screens for common mental disorders. It comprises 20 items evaluating depressive/anxious and somatic symptoms, reduced vital energy, and depressive thoughts over the past month. The possible answers are 'yes/no', with each affirmative answer equating to one point in the final score. The cut-off point for significant symptom severity are ≥ 8 for women and ≥ 6 for men [14].

Pittsburgh Sleep Quality Index (PSQI). Sleep quality was assessed using the PSQI, a self-administered questionnaire evaluating sleep quality over the past month. It includes 24 items grouped into 7 components: subjective sleep quality, sleep latency, sleep duration, sleep efficiency, sleep disturbance, use of sleep medication, and daytime dysfunction. Each component is scored from 0 to 3, with a total score ranging from 0 to 21 points. The lower the score, the better the quality of sleep. A score > 5 indicates poor sleep quality [16]. The Brazilian validated version of PSQI was used in this study [17].

A pilot study with a small number of students was previously conducted to assess the reliability and validity of the questionnaires, especially the first two which are not officially standardized. The construction and content aspects were validated by a healthcare professional, while clarity was

assessed with students. Participants were invited to participate voluntarily and provide feedback on the clarity, readability, and comprehensibility of the questionnaire. Feedback was analyzed, and a finalized version was created. Participants reported that the questionnaire were clear and easy to complete. Data from the pilot study were not included in the current study.

2.3 Statistical Analysis

Variables were created for each questionnaire based on the total score: presence (MD+) or absence (MD-) of mental distress (SRQ-20), and good (SQ+) and poor (SQ-) sleep quality (PSQI), according to the specified cut-off points. Initially, a descriptive analysis of the population was performed according to the variables (sleep quality and mental distress, sociodemographic variables, and self-perception of health). Categorical variables were represented by their absolute frequencies and percentages, and the Kolmogorov-Smirnov distribution test were used to analyze the data. Responses scores for the SRQ-20 and PSQI are reported as mean \pm standard errors of the mean. To evaluate the outcome variable (presence or absence of mental distress or good or poor sleep quality) and possible exposures (presence or absence of each predictive factor), the Chi-square test was used. Pearson's correlation analysis was used to establish the relationship between PSQI and SRQ-20 scores. Comparisons between sociodemographic characteristics (between sexes and need for mental health support) and scores on the SRQ-20 and PSQI were made using the two-tailed Student's t-test. The significance level was set at $P < .05$.

3. RESULTS AND DISCUSSION

The present study evaluated the effects of COVID-19 pandemic restrictive measures on sleep quality and mental health among university medical students. Additionally, demographic and socioeconomic variables, and perceived stress related to school and the COVID-19 pandemic were assessed to identify potential influencing factors.

A total of 210 undergraduate students participated in the study, with a uniform distribution across different periods of their medical education. The participants were predominantly women (76.7%) and men

comprised 23.3%. The ages ranged from 18 to 60 years, with the majority (73.4%) aged between 20 to 25 years. Most participants were single (91.4%), identified as Catholic (41.4%), had a normal body mass index (67.7%), and identified their skin color as white (72.4%). Regarding living conditions, most reported a family income between 4 and 10 minimum wages (34.3%), lived with their family (83.3%), and belonged to a family group of 3 to 5 people (78.1%). A majority (57.6%) claimed to be in good health and had not sought support or treatment for mental disorders (69.1%) (Table 1).

Our findings provide new evidence that the Brazilian lockdown significantly impacted both sleep and psycho-emotional well-being among medical students.

In the assessment of mental health, 133 out of the 210 students (63.3%) had scores suggesting mental distress according to the SRQ-20 questionnaire. A significant difference was observed in relation to gender, with women showing higher mental distress scores compared to men (women: 9.87 ± 0.37 vs. men: 5.98 ± 0.62 ; $P < .001$). The proportion of women experiencing mental distress was 67.7%, compared to 49% among men ($P < .01$). As expected, mental distress was higher among those who reported poor health in the SPH questionnaire ($P < .001$). Additionally, students receiving treatments or support for mental health concerns had higher distress scores, with 86.1% of them showing mental distress compared to 56.1% of those not receiving any treatment ($P < .001$) (Table 1).

The PSQI questionnaire revealed that 186 out of the 210 students (88.6%) had poor sleep quality. No significant differences were found between the sociodemographic variables evaluated (Table 1).

Although the overall PSQI scores did not differ between men and women, gender was significantly associated with mental health as assessed by the SRQ-20. Therefore, PSQI component scores were also compared between genders. Differences were found only in sleep disturbances, with women reporting greater sleep disturbances compared to men (women: 1.35 ± 0.04 vs. men: 1.26 ± 0.06 ; $P < .05$) (Table 2).

Table 1. Distribution of students according to demographic and socioeconomic variables, and prevalence of mental distress and poor sleep quality

Variable	N (210)	%	% MD+	P	% SQ-	P
Gender						
Men	49	23.3	49.0	< 0.01	91.8	0.57
Women	161	76.7	67.7		87.6	
Age (years)						
< 19	24	11.4	54.2	0.13	79.2	0.28
20-22	80	38.1	71.2		87.5	
23-25	74	35.3	68.9		93.2	
> 26	32	15.2	53.1		87.5	
Course period						
1 ^o	12	5.7	83.3	0.64	91.7	0.58
2 ^o	20	9.5	70.0		80.0	
3 ^o	25	11.9	60.0		84.0	
4 ^o	30	14.3	63.3		93.3	
5 ^o	33	15.7	75.8		97.0	
6 ^o	27	12.9	55.6		88.9	
7 ^o	32	15.2	62.5		88.9	
8 ^o	31	14.8	64.5		87.1	
Body mass index						
Underweight (< 18.5)	7	3.3	85.7	0.46	87.5	0.68
Normal (18.5-24.9)	142	67.7	63.4		87.2	
Overweight (25.0-29.9)	53	25.2	67.9		90.6	
Obese (> 30.0)	8	3.8	75.0		100.0	
Ethnicity						
White	152	72.4	65.8	0.46	87.5	0.61
Brown	52	24.8	69.2		92.3	
Black	3	1.4	33.3		66.7	
Yellow	2	0.9	0.0		100.0	
Indigenous	1	0.5	100.0		100.0	
Religion						
Catholic	87	41.4	63.2	0.47	86.2	0.13
Evangelical	50	23.8	74.0		84.0	
Spiritist	25	11.9	64.0		88.0	
No religion	43	20.5	65.1		100.0	
Other	5	2.4	40.0		80.0	
Marital status						
Single	192	91.4	67.7	0.03	88.5	0.93
Married	15	7.1	53.3		86.7	
Other	3	1.5	0.0		100.0	
Family income						
< 2 minimum wages	4	1.9	50.0	0.72	100.0	0.71
2-4 minimum wages	20	9.5	70.0		95.0	
4-10 minimum wages	72	34.3	70.8		87.5	
10-20 minimum wages	68	32.4	63.2		89.7	
> 20 minimum wages	46	21.9	60.9		84.8	
Living condition						
With family members	175	83.3	66.3	0.92	90.3	0.16
Friends	3	1.4	66.7		81.3	
Alone	32	15.3	62.5		66.7	
Number of people in the residence						
I live alone	8	3.8	87.5		100.0	

Variable	N (210)	%	% MD+	P	% SQ-	P
< 2 people	34	16.2	61.8	0.55	88.2	0.65
3-5 people	164	78,1	65.2		87.8	
> 5 people	4	1.9	75.0		100.0	
Health self-perception						
Too bad	2	0.9	100.0	< 0.001	100.0	0.70
Bad	3	1.4	100.0		100.0	
Regular	43	20.6	81.4		90.7	
Good	121	57.6	66.9		89.3	
Very good	41	19.5	41.5		82.9	
Treatment/support for a mental health concern						
Yes	65	30.9	86.1	< 0.001	93.8	0.17
No	145	69.1	56.5		86.2	

Table 2. Comparative scores obtained for each component of the PSQI between men and women students

PSQI	MALE (n = 49)	FEMALE (n = 161)	P
Sleep quality	1.33 ± 0.12	1.44 ± 0.06	0.39
Sleep latency	1,55 ± 0.14	1.55 ± 0.07	0.55
Sleep duration	1.06 ± 0.16	1.14 ± 0.09	0.55
Sleep efficiency	2.43 ± 0.16	2.20 ± 0.10	0.41
Sleep disturbances	1.26 ± 0.06	1.35 ± 0.04	< 0.05
Sleeping medications	0.45 ± 0.13	0.37 ± 0.07	0.72
Daytime dysfunction	1.53 ± 0.11	1.61 ± 0.06	0.10
Global score	9.17 ± 0.49	9.32 ± 0.24	0.24




Fig. 1. The relationship between sleep quality (PSQI scores) and mental distress (SRQ-20 scores)

A significant positive correlation was found between the PSQI and SRQ-20 scores ($r = 0.611$, $P < .001$) (Fig. 1).

In summary, the prevalence of common mental disorders reached 63% of students, while poor sleep quality affected nearly 90% of them, with a greater impact observed in female students compared to males.

Common mental disorders, as evaluated by the SRQ-20, include minor psychiatric disorders such as anxiety, depressive, and somatoform disorders, characterized by symptoms like insomnia, fatigue, irritability, forgetfulness, difficulty concentrating, and somatic issues [14]. Previous studies using the same questionnaire estimated the prevalence of common mental disorders in the Brazilian population to be between 22% and 35% [18-21]; however, our results showed a prevalence of 63% among medical students, exceeding the national average.

The medicine course itself appears to be a significant risk factor for developing mental disorders, with prevalence rates higher than the national average [9,10,22]. This higher prevalence is also observed when comparing medical students to those in other health-related fields, such as dentistry, nursing, and physical education [23]. The stressful situations to which medical students are exposed during their course may contribute to the higher prevalence of mental distress among them [9,10,22]. Consequently, it is expected that the current pandemic would exacerbate their mental health issues, given the direct exposure of medical students to changes in health care systems during the COVID-19 pandemic.

A key limitation of our study is the absence of pre-pandemic records for comparison. However, we observed that the prevalence of mental distress is significantly higher compared to previous studies that used the same SRQ-20 among medical students before the pandemic, where the prevalence ranged between 34 and 45% [23-25]. The COVID-19 pandemic has heightened the prevalence of mental disorders in the general population, with factors such as decreased social interactions, income reduction, economic stress, the threat of contracting the virus, and uncertainty about the future contributing to increased stress, depression, and anxiety. These factors, coupled with social distancing and confinement, have intensified

these symptoms [5,6,26,27]. Considering that medical students already represent a higher risk group for mental disorders, these factors strengthen the hypothesis that social isolation during the pandemic has been crucial in increasing mental disorders cases among these students.

Female students in our study exhibited higher SRQ-20 scores than their male counterparts, with 68% of women affected compared to 49% of the men. Extensive literature indicates that women generally have higher rates of stress and mental disorders, both in the general population and among medical students [28-35]. Traditional gender roles may lead to differences in how attitudes and emotions towards life experiences are expressed, with women more likely to report higher levels of anxiety and being more affected by traumatic events [28-30]. Additionally, sex hormones may play a role in the higher prevalence of depression, anxiety stress-related disorders among women [36,37]. It is also plausible that the SRQ-20 measures symptoms more commonly experienced by women, leading to higher scores among them. Using a scale more focused on men's health issues might reveal different patterns [38]. Moreover, it is also important to note that although our sample is representative of the gender distribution among the respondents (with the academic population consisting of approximately 70% women and 30% men), the smaller number of male participants might also suggest a potential bias due to this result.

Sleep is a crucial mechanism for regulating and recovering biological and cognitive functions. Poor sleep quality or lack of sleep affects the quality of life at any age, negatively impacting cognitive performance, mood, memory, concentration, learning, logical reasoning, and creativity [39]. This underscores the importance of assessing the sleep quality among university students, especially during the stressful situation of the COVID-19 pandemic [40].

Our study found that 88.6% of students had poor sleep quality, with a mean PSQI global score of 9.245. These findings align with a previous study conducted with medical students in Saudi Arabia, where 85.5% reported poor sleep quality with a mean PSQI score of 8.356 [41]. In 2012, another study in the same college found that only 36.3% of students had poor sleep quality, highlighting a significant increase during the pandemic [41,42]. This alarming trend emphasizes the severe

impact of the pandemic on sleep quality among medical students, who were already vulnerable to poor sleep quality compared to general population [43-46]. Similar studies using PSQI questionnaire to evaluate sleep quality found a prevalence between 44% and 72% to poor quality sleep among undergraduate medical students [47-51].

Several studies have cited reasons for the high prevalence of sleep disorders during the pandemic period, beyond psychological stress. These include adjustment to modified teaching-learning formats, loss of social connectedness, increased smartphone and social media use close to bedtime, and circadian misalignment [48,52-54]. Studies have shown that insomnia prevalence increased significantly during the lockdown, with university students reporting longer sleep times and more time in bed [53, 55-58]. Factors such as online lessons and the elimination of commuting times to clinical, university, and hospitals, may have allowed for more flexible schedules and additional sleep time [55,57,58].

In summary, sleep disturbances among university students present an alarming trend with significant consequences on their mental health. Our results showed a significant correlation between sleep quality and common mental disorders, reinforcing the relationship between sleep and emotions well-being [47,59-62].

Women, in addition to having a higher prevalence of common mental disorders, also reported more sleep disturbances, consistent with literature findings that female students are more likely to be poor sleepers compared to their men counterparts [33,50,63-66]. Studies conducted in the general population also report that sleep problems are more common in women [67,68].

Usually, self-assessing and reporting health problems are more frequent among females than males; therefore, they use health services more often; thus, this result confirms information of a recent study that points to this population as one of the most affected by these conditions during the pandemic [63]. These findings highlight the need for sleep awareness programs with gender-specific approaches.

Still in this regard, interesting results were found by Fowler and colleagues (2022); before the

stay-at-home order, women reported significantly lower sleep quality than men, but post-order, men reported significantly worse sleep quality, reaching levels similar to women [48].

It is important to acknowledge the limitations of this study. First, the cross-sectional design limits our ability to establish causal relationships between sleep quality and mental health issues. Longitudinal data are needed to better define these relationships throughout the pandemic. Second, our data relied on subjective self-reported information without clinical or instrumental examinations, which may introduce recall bias. Future research should include objective measures of sleep and structured interviews to investigate mental health problems for confirmation of our findings. Lastly, data collection occurred during the pandemic, preventing inferences about post-pandemic associations.

4. CONCLUSION

In conclusion, this study has provided insights into the impact of COVID-19 on medical students' well-being and provided an opportunity to address their concerns. The strict preventive measures, and the radical changes in the lifestyle of medical students due COVID-19 pandemic had a negative impact on the quality of their sleep and on their mental state. Medical students showed alarming levels of poor sleep quality and psychiatric disorders during the COVID-19 pandemic, with women being more affected than men. The findings could be helpful in designing appropriate implementation of some interventions for well-being in pandemic condition, and a special focus must be given to the most affected groups (such as female students).

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

CONSENT

All authors declare that online written consent was obtained from the voluntaries. The students first accessed the written consent form on the initial screen of the electronic questionnaire and could only proceed after consenting.

ETHICAL APPROVAL

All experimental procedures received approval from the National Research Ethics Commission of Brazil Platform (CAAE 32563220.0.0000.0008).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Rothan HA, Byrareddy SN. The epidemiology and pathogenesis of coronavirus disease (COVID-19) outbreak. *J Autoimmun.* 2020;109:102433.
2. Cucinotta D, Vanelli M. WHO Declares COVID-19 a Pandemic. *Acta Biomed.* 2020;91(1):157-160.
3. Jiang F, et al. Review of the clinical characteristics of coronavirus disease 2019 (COVID-19). *J Gen Intern Med.* 2020;35(5):1545-1549.
4. Wilder-Smith A, Freedman DO. Isolation, quarantine, social distancing and community containment: Pivotal role for old-style public health measures in the novel coronavirus (2019-nCoV) outbreak. *J Travel Med.* 2020;27(2).
5. Brooks SK, et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet.* 2020;395(10227):912-920.
6. Xiong J, et al. Impact of COVID-19 pandemic on mental health in the general population: A systematic review. *J Affect Disord.* 2020;277:55-64.
7. Sahu P. Closure of universities due to coronavirus disease 2019 (COVID-19): Impact on education and mental health of students and academic staff. *Cureus.* 2020;12(4):e7541.
8. Chandratre S. Medical students and Covid-19: Challenges and supportive strategies. *J Med Educ Curric Dev.* 2020;7:2382120520935059.
9. Pacheco JP, et al. Mental health problems among medical students in Brazil: A systematic review and meta-analysis. *Braz J Psychiatry.* 2017;39(4):369-378.
10. Rotenstein LS, et al. Prevalence of depression, depressive symptoms, and suicidal ideation among medical students: A systematic review and meta-analysis. *JAMA.* 2016;316(21):2214-2236.
11. Quek TT, et al. The global prevalence of anxiety among medical students: A meta-analysis. *Int J Environ Res Public Health.* 2019;16(15).
12. Dyrbye LN, Thomas MR, Shanafelt TD. Medical student distress: Causes, consequences, and proposed solutions. *Mayo Clin Proc.* 2005;80(12):1613-22.
13. Bore M, Kelly B, Nair B. Potential predictors of psychological distress and well-being in medical students: A cross-sectional pilot study. *Adv Med Educ Pract.* 2016;7:125-35.
14. Mari JJ, Williams P. A validity study of a psychiatric screening questionnaire (SRQ-20) in primary care in the city of Sao Paulo. *Br J Psychiatry.* 1986;148:23-6.
15. Beusenbergh M, Orley JH, Health WHODOM. A User's guide to the self reporting questionnaire (SRQ / compiled by Beusenbergh M, Orley J. World Health Organization; 1994.
16. Buysse DJ, et al. The pittsburgh sleep quality index: A new instrument for psychiatric practice and research. *Psychiatry Res.* 1989;28(2):193-213.
17. Bertolazi AN, et al. Validation of the brazilian portuguese version of the pittsburgh sleep quality index. *Sleep Med.* 2011;12(1):70-5.
18. Coledam DHC, et al. Prevalence of common mental disorders among Brazilian workers: Systematic review and meta-analysis. *Cien Saude Colet.* 2022;27(2):579-591.
19. Lima MS, et al. Stressful life events and minor psychiatric disorders: An estimate of the population attributable fraction in a Brazilian community-based study. *Int J Psychiatry Med.* 1996;26(2):211-22.
20. Ludermir AB, Melo Filho DA. Living conditions and occupational organization associated with common mental disorders. *Rev Saude Publica.* 2002;36(2):213-21.
21. Anselmi L, et al. Prevalence and early determinants of common mental disorders in the 1982 birth cohort, Pelotas, Southern

- Brazil. *Rev Saude Publica*. 2008;42 Suppl 2(Suppl 2):26-33.
22. Soares SJB, et al. Common mental disorders among medical students: Systematic review and meta-analysis of Brazilian studies. *Sao Paulo Med J*. 2022;140(4):615-622.
 23. Facundes VL, Ludermir AB. Common mental disorders among health care students. *Braz J Psychiatry*. 2005;27(3):194-200.
 24. Costa EF, et al. Common mental disorders among medical students at Universidade Federal de Sergipe: A cross-sectional study. *Braz J Psychiatry*. 2010;32(1):11-9.
 25. Lima MC, Domingues Mde S, Cerqueira AT. Prevalence and risk factors of common mental disorders among medical students. *Rev Saude Publica*. 2006;40(6):1035-41.
 26. Usher K, Bhullar N, Jackson D. Life in the pandemic: Social isolation and mental health. *J Clin Nurs*. 2020;29(15-16):2756-2757.
 27. Lima CKT, et al. The emotional impact of Coronavirus 2019-nCoV (New Coronavirus disease). *Psychiatry Res*. 2020;287:112915.
 28. Batra K, et al. Assessing the psychological impact of covid-19 among college students: An evidence of 15 countries. *Healthcare (Basel)*. 2021;9(2).
 29. Chaplin TM, et al. Gender differences in response to emotional stress: An assessment across subjective, behavioral, and physiological domains and relations to alcohol craving. *Alcohol Clin Exp Res*. 2008;32(7):1242-50.
 30. Alegria M, et al. Social determinants of mental health: Where we are and where we need to go. *Curr Psychiatry Rep*. 2018;20(11):95.
 31. Riecher-Rossler A. Sex and gender differences in mental disorders. *Lancet Psychiatry*. 2017;4(1):8-9.
 32. Beutel ME, et al. Mental health and loneliness in the German general population during the COVID-19 pandemic compared to a representative pre-pandemic assessment. *Sci Rep*. 2021;11(1):14946.
 33. Harries AJ, et al. Effects of the COVID-19 pandemic on medical students: A multicenter quantitative study. *BMC Med Educ*. 2021;21(1):14.
 34. Huang Y, Zhao N. Generalized anxiety disorder, depressive symptoms and sleep quality during COVID-19 outbreak in China: A web-based cross-sectional survey. *Psychiatry Res*. 2020;288:112954.
 35. Paludo AC, et al. Female students are the most psychologically affected by the COVID-19 outbreak: A case study in an academic community in Brazil. *Rev Assoc Med Bras*. 2021;67(5):741-746.
 36. Kuehner C. Why is depression more common among women than among men? *Lancet Psychiatry*. 2017;4(2):146-158.
 37. Li SH, Graham BM. Why are women so vulnerable to anxiety, trauma-related and stress-related disorders? The potential role of sex hormones. *Lancet Psychiatry*. 2017;4(1):73-82.
 38. Pereira MB, et al. Mental health of medical students before and during covid-19 pandemic: A 3-year prospective study. *Med Sci Educ*. 2022;32(4):873-881.
 39. Pagel JF, Kwiatkowski CF. Sleep complaints affecting school performance at different educational levels. *Front Neurol*. 2010;1:125.
 40. Semyachkina-Glushkovskaya O, et al. Brain mechanisms of Covid-19-sleep disorders. *Int J Mol Sci*. 2021;22(13).
 41. Meo SA, et al. Impact of COVID-19 pandemic on sleep quality among medical and general science students: King Saud University Experience. *Pak J Med Sci*. 2022;38(3Part-I):639-644.
 42. Alfawaz HA, et al. Psychological well-being during COVID-19 lockdown: Insights from a Saudi State University's Academic Community. *J King Saud Univ Sci*. 2021;33(1):101262.
 43. Azad MC, et al. Sleep disturbances among medical students: A global perspective. *J Clin Sleep Med*. 2015;11(1):69-74.
 44. Ahmed N, Sadat M, Cukor D. Sleep knowledge and behaviors in medical students: Results of a single center survey. *Acad Psychiatry*. 2017;41(5):674-678.
 45. Ayala EE, et al. A cross-sectional snapshot of sleep quality and quantity among us medical students.

- Acad Psychiatry. 2017;41(5):664-668.
46. Seoane HA, et al. Sleep disruption in medicine students and its relationship with impaired academic performance: A systematic review and meta-analysis. *Sleep Med Rev.* 2020;53:101333.
 47. Mishra J, et al. Sleep quality and associated factors among undergraduate medical students during Covid-19 confinement. *Clin Epidemiol Glob Health.* 2022;15:101004.
 48. Fowler LA, Kumte N. The effect of COVID-19 pandemic stay-at-home orders on sleep deprivation in medical students: A retrospective study. *SN Soc Sci.* 2022;2(3):29.
 49. Saguem BN, et al. Predictors of sleep quality in medical students during COVID-19 confinement. *Encephale.* 2022;48(1):3-12.
 50. Sundas N, et al. Sleep quality among medical students of a tertiary care hospital: A descriptive cross-sectional study. *JNMA J Nepal Med Assoc.* 2020;58(222):76-79.
 51. Puteikis K, Mameniskyte A, Mameniskiene R. Sleep quality, mental health and learning among high school students after reopening schools during the Covid-19 pandemic: Results of a cross-sectional online survey. *Int J Environ Res Public Health.* 2022;19(5).
 52. Altena E, et al. Dealing with sleep problems during home confinement due to the COVID-19 outbreak: Practical recommendations from a task force of the European CBT-I Academy. *J Sleep Res.* 2020;29(4):e13052.
 53. Cellini N, et al. Changes in sleep pattern, sense of time and digital media use during COVID-19 lockdown in Italy. *J Sleep Res.* 2020;29(4):e13074.
 54. Marelli, S., et al., Impact of COVID-19 lockdown on sleep quality in university students and administration staff. *J Neurol*, 2021. 268(1): p. 8-15.
 55. Luciano F, et al. COVID-19 lockdown: Physical activity, sedentary behaviour and sleep in Italian medicine students. *Eur J Sport Sci.* 2021;21(10):1459-1468.
 56. Staller N, Kalbacher L, Randler C. Impact of pandemic lockdown on learning behaviour and sleep quality in German students: Results of an online survey before and during the pandemic. *Somnologie (Berl).* 2022;26(2):98-105.
 57. Staller N, Randler C. Changes in sleep schedule and chronotype due to Covid-19 restrictions and home office. *Somnologie (Berl).* 2021;25(2):131-137.
 58. Wright KP, Jr, et al. Sleep in university students prior to and during Covid-19 Stay-at-Home orders. *Curr Biol.* 2020;30(14):R797-R798.
 59. Peixoto KO, et al. Association of sleep quality and psychological aspects with reports of bruxism and TMD in Brazilian dentists during the Covid-19 pandemic. *J Appl Oral Sci.* 2021;29:e20201089.
 60. Becker SP, et al. Sleep in a large, multi-university sample of college students: Sleep problem prevalence, sex differences, and mental health correlates. *Sleep Health.* 2018;4(2):174-181.
 61. Zhang Y, Peters A, Chen G. Perceived stress mediates the associations between sleep quality and symptoms of anxiety and depression among college nursing students. *Int J Nurs Educ Scholarsh.* 2018;15(1).
 62. Choueiry N, et al. Insomnia and relationship with anxiety in university students: A cross-sectional designed study. *Plos One.* 2016;11(2):e0149643.
 63. Bigalke JA, Greenlund IM, Carter JR. Sex differences in self-report anxiety and sleep quality during Covid-19 stay-at-home orders. *Biol Sex Differ.* 2020;11(1):56.
 64. Almojali AI, et al. The prevalence and association of stress with sleep quality among medical students. *J Epidemiol Glob Health.* 2017;7(3):169-174.
 65. Eleftheriou A, et al. Sleep quality and mental health of medical students in greece during the Covid-19 pandemic. *Front Public Health.* 2021;9:775374.
 66. Xie J, et al. Depressive symptoms, sleep quality and diet during the 2019 novel coronavirus epidemic in china: A survey of medical students. *Front Public Health.* 2020;8:588578.

67. Mallampalli MP, Carter CL. Exploring sex and gender differences in sleep health: A Society for Women's Health Research Report. J Womens Health (Larchmt). 2014;23(7):553-62.
68. Tang J, et al. Gender and regional differences in sleep quality and insomnia: A general population-based study in hunan province of china. Sci Rep. 2017;7:43690.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/119197>