



# Phytobiomes: Harnessing the Power of Microbial Communities for Plant Health

**Ashwini Kumar <sup>a++</sup>, R Manisha <sup>b#\*</sup>,  
Haralakal Keerthi Kumari <sup>c#</sup>, Neha <sup>d</sup>, P. Kalaiselvi <sup>e†</sup>,  
Saloni Jaswal <sup>f‡</sup>, Devansh Gupta <sup>g^</sup> and Saksham Tinna <sup>h‡</sup>**

<sup>a</sup> R. A. K. College of Agriculture, RVSKVV, Sehore, (M.P.), India.

<sup>b</sup> Department of Plant Pathology, Faculty of Agriculture, Annamalai University, Chidambaram, India.

<sup>c</sup> Department of Agricultural Microbiology, University of Agricultural Sciences Bangalore, India.

<sup>d</sup> Department of Floriculture and Landscape Architecture, Chaudhary Charan Singh Haryana Agricultural University, Hisar, Haryana, India.

<sup>e</sup> ICAR- Krishi Vigyan Kendra, Tamil Nadu Agricultural University, Salem, Tamil Nadu, India.

<sup>f</sup> Department of Soil Sciences, College of Agriculture, Punjab Agricultural University, Punjab, India.

<sup>g</sup> Department of Soil Science, Punjab Agricultural University, Ludhiana, Punjab, India.

<sup>h</sup> Department of Soil Sciences, College of Agriculture, Punjab Agricultural University, Punjab, India.

## Authors' contributions

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## Article Information

DOI: <https://doi.org/10.9734/mrji/2024/v34i101486>

### Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/122866>

**Review Article**

**Received: 13/07/2024**

**Accepted: 16/09/2024**

**Published: 19/09/2024**

<sup>++</sup> Assistant Professor;

<sup>#</sup> Ph.D. Scholar;

<sup>†</sup> Associate Professor (ENS);

<sup>‡</sup> B.Sc. (Hons) Agriculture;

<sup>^</sup> B.Sc. (Hons) Horticulture;

\*Corresponding author: E-mail: [manisharamesh222@gmail.com](mailto:manisharamesh222@gmail.com);

**Cite as:** Kumar, Ashwini, R Manisha, Haralakal Keerthi Kumari, Neha, P. Kalaiselvi, Saloni Jaswal, Devansh Gupta, and Saksham Tinna. 2024. "Phytobiomes: Harnessing the Power of Microbial Communities for Plant Health". *Microbiology Research Journal International* 34 (10):1-19. <https://doi.org/10.9734/mrji/2024/v34i101486>.

## ABSTRACT

Phytobiomes encompass the dynamic ecosystems surrounding plants, which include a diverse range of microbial communities such as fungi, bacteria, algae, viruses, and nematodes. These microorganisms interact intricately with plants, influencing their health, growth, and resilience. Beneficial fungi form symbiotic relationships with plant roots, enhancing nutrient absorption, while bacteria play a role in nitrogen fixation and disease suppression. Algae contribute to soil health and organic matter decomposition, and viruses, though often seen as harmful, can provide resistance to certain pathogens. Nematodes, both beneficial and parasitic, regulate soil nutrient cycles and influence plant growth. Harnessing the potential of these microbial communities offers promising solutions for sustainable agriculture. By promoting beneficial interactions within the phytobiome, it is possible to improve plant productivity, enhance nutrient use efficiency, and increase tolerance to environmental stresses such as drought, salinity, and pathogens. Phytobiome research aims to uncover the complex relationships within these microbial networks to develop novel agricultural practices that reduce reliance on chemical inputs. This holistic approach not only promotes plant health but also ensures ecological balance, contributing to the long-term sustainability of agroecosystems in the face of environmental challenges. Understanding the power of microbial communities within phytobiomes is key to advancing future agricultural innovations.

*Keywords: Phytobiomes; biofertilizers; chemical; viruses; biology; crops; biocontrol.*

## 1. INTRODUCTION

Phytobiomes encompass the intricate communities of microorganisms—bacteria, fungi, viruses, archaea, and other microscopic life forms—that live in close association with plants. These microbial communities are found on the surface of leaves and roots, within plant tissues, and in the surrounding soil. Far from being mere inhabitants of the plant environment, these microorganisms engage in complex interactions with their host plants, influencing critical aspects of plant health, growth, and resilience [1]. The concept of phytobiomes extends beyond just the microbiome of a plant; it includes the entire biological and physical environment in which plants grow, encompassing soil, climate, other plants, insects, and agricultural practices. This holistic view recognizes that plant health is not determined solely by the plant's genetics but is also shaped by the interactions within its surrounding ecosystem [2]. Recent advances in molecular biology, high-throughput sequencing, and bioinformatics have significantly expanded our understanding of phytobiomes. These technologies have revealed that phytobiomes play essential roles in nutrient cycling, enhancing plant growth, protecting against pathogens, and helping plants withstand environmental stresses such as drought and salinity [3]. The ability of phytobiomes to influence these processes opens up new avenues for enhancing agricultural productivity in sustainable ways. As global agriculture faces increasing challenges from climate change, soil degradation, and the need

to produce more food with fewer resources, the potential of phytobiomes offers a promising path forward [4]. By harnessing and optimizing these microbial communities, we can develop innovative strategies for crop management that reduce the reliance on chemical inputs, improve crop resilience, and support the long-term sustainability of agricultural systems [5].

A wide variety of microorganisms, including as bacteria, archaea, Protistas, fungus, and viruses, are able to flourish and multiply in the niches that plants offer for them. These bacteria are very important to the overall health and production of plants when they are exposed to regular environmental conditions [6]. Plants and the microbiota that are linked with them make up a holobiont, which is a system in which evolutionary selection contributes to the stability of the overall system. Bacteria and fungus are the species that have received the greatest attention from researchers; however, other taxa such as archaea, viruses, protistas, and nematodes also play significant roles [7]. Through the use of genome-wide association studies (GWAS), plant loci that are associated with microbial diversity and structure have been discovered. Microbial genomics has made it possible to identify genes that are connected with plant-associated lifestyles. These genes have the potential to assist the surrounding microbiota in coping with the defences of the host plant, which is necessary for successful colonization [8]. The methods and techniques used in plant microbiome research play an

essential part in the study of the whole microbiota of host plants and the functional characteristics of that microbiota, with the goal of improving plant growth and production in conjunction with biotic and abiotic stress. Recent developments in technologies like as metagenomics, transcriptomics, and genome editing have the potential to contribute to the intensification of the link between legumes and microbiomes, therefore directing advances towards improved nitrogen nutrition [9]. The primary objectives of this review are to investigate recent information concerning the plant microbiome and the direct and indirect plant-beneficial mechanisms that it possesses, to discuss the gaps in research, and to highlight the current knowledge and understanding of plant-associated microbial communities and the functional attributes that they possess for the purpose of enhancing sustainable agricultural productivity [10].

## 2. HOW HOST FUNCTIONS

In the same way that bacteria in the human body interact with one another and have an impact on the health of plants, these bacteria also have an effect on the soil. There is a correlation between the two. At the level of the community, each of the three forms of microbiomes—the rhizomicrobiome, the phyllosphere, and the endomicrobiome—is unique from the other two [11]. Furthermore, phytomicrobiomes that are related with various plant components are distinct from one another on account of their individuality. The composition of the phytomicrobiome is determined by a variety of variables, such as the types of plants, the phases of growth that the plants go through, and the circumstances of the environment itself [12]. These factors take into account the kinds of plants that are there. Actinobacteria, proteobacteria, planctomycetes, verrucomicrobia, firmicutes, acidobacteria, bacteria, and gemmatimonadetes are the most common types of microorganisms that may be found in root microbiomes. Root microbiomes are home to a vast range of microorganisms. Core microbiomes, which are present in the majority of samples taken from a single plant species, are either directly inherited by the plant species or are actively chosen by the plant species [13]. Examples of both of these processes are described below. Core microbiomes may be found in the vast majority of tissue samples taken from plants. Additionally, they are vulnerable to being altered by

genotypes, climatic circumstances, and management approaches. This is in addition to the fact that they co-evolved with many other plant species. This is due to the fact that they are often capable of being shaped concurrently [14]. The fundamental microbiome of a number of different crop species may be characterized with the intention of doing more research and development. The possibility of this happening is not inconceivable. This may entail the cultivation of endophytic microorganisms as well as the genetic modification of plants in order to improve the nutritional content of the plants and boost their general health. The goal is to make the plants healthier overall [15]. The relationship that exists between the core microbiomes of various plant species and genotypes is an example of the microbiome assemblage that is driven by the host. This connection can be observed in the core microbiomes of different plant species [16]. The 'hub microbiota approach' is used in order to get an understanding of the interaction that takes place between the core group of the microbiome and the functional qualities that it has. This is done with the intention of achieving the objective of improving the health of plants. When it comes to the control of all of the functional networks that are included within the plant microbiome, certain microbial taxa and genera, in addition to populations, are responsible for the responsibility. This microbiome is a network that is very intricate and connects with other networks [17, 18].

## 3. IMPORTANCE OF MICROBIAL COMMUNITIES IN PLANT HEALTH

Microbial communities, often referred to as the plant microbiome, play a crucial role in maintaining and enhancing plant health. These communities are composed of bacteria, fungi, archaea, viruses, and other microorganisms that live in and around plants, including the rhizosphere (soil surrounding the roots), phyllosphere (above-ground plant parts), and endosphere (inside plant tissues) [19]. Their importance in plant health can be understood through several key functions:

### 3.1 Nutrient Acquisition

Microbial communities are vital for nutrient cycling and availability. For instance:

- **Nitrogen Fixation:** Certain bacteria, such as *Rhizobium* species, fix atmospheric nitrogen into a form that plants can absorb

and use, which is critical for plant growth [20].

- **Phosphorus Solubilisation:** Microbes such as phosphate-solubilizing bacteria and mycorrhizal fungi convert insoluble phosphorus in the soil into forms that plants can take up, thereby improving nutrient availability.
- **Decomposition of Organic Matter:** Microbes decompose organic matter, releasing nutrients back into the soil and making them available to plants [21].

### 3.2 Disease Suppression

Microbial communities help protect plants from pathogens through various mechanisms:

- **Competition:** Beneficial microbes outcompete harmful pathogens for space and resources, thereby preventing disease [22].
- **Antimicrobial Production:** Some microbes produce antibiotics, antifungals, or other substances that inhibit the growth of pathogens.
- **Induced Systemic Resistance (ISR):** Certain beneficial microbes trigger plant defence mechanisms, making plants more resistant to a broad range of pathogens [23].

### 3.3 Stress Tolerance

Microbial communities enhance plant resilience to abiotic stresses such as drought, salinity, and temperature extremes:

- **Drought Resistance:** Some microbes improve water uptake or produce substances that help plants retain water under drought conditions.
- **Salinity Tolerance:** Certain bacteria can mitigate the negative effects of high salt levels on plants, allowing them to grow in saline environments [24].
- **Temperature Extremes:** Microbes can produce proteins and other compounds that help plants survive extreme temperatures.

### 3.4 Growth Promotion

Microbes directly promote plant growth by:

- **Hormone Production:** Many microbes produce plant hormones like auxins,

gibberellins, and cytokines', which stimulate root development, shoot growth, and overall plant vigour [25].

- **Nutrient Uptake Enhancement:** Mycorrhiza fungi extend the root system's reach, allowing plants to access nutrients and water more efficiently [26].

### 3.5 Ecosystem Stability and Biodiversity

Microbial communities contribute to the stability and biodiversity of ecosystems by:

- **Maintaining Soil Health:** They improve soil structure, enhance water retention, and increase soil fertility.
- **Supporting Plant Diversity:** Microbes promote the coexistence of different plant species by mediating plant-microbe and plant-plant interactions, which can lead to more resilient ecosystems [27].

### 3.6 Bioremediation

Microbial communities play a role in bioremediation by breaking down pollutants and detoxifying soils, thereby enabling plants to grow in contaminated environments. This is particularly important in restoring degraded or polluted agricultural lands [28].

### 3.7 Plant-Microbe Symbiosis

Symbiotic relationships between plants and microbes, such as those between legumes and nitrogen-fixing bacteria, or between plants and mycorrhiza fungi, are fundamental to plant health. These partnerships are essential for nutrient exchange, growth, and survival in various environments [29].

### 3.8 Adaptation to Changing Environments

Microbial communities help plants adapt to changing environmental conditions by facilitating rapid responses to stressors, such as climate change, pests, and diseases. This adaptability is crucial for the long-term sustainability of agricultural systems [30].

## 4. THE ROLE OF PHYTOBIOMES IN MODERN AGRICULTURE

Phytobiomes play a pivotal role in modern agriculture by serving as natural allies in

enhancing crop productivity, sustainability, and resilience. These complex microbial communities, encompassing bacteria, fungi, viruses, and other microorganisms, interact intimately with plants, facilitating essential processes such as nutrient acquisition, disease suppression, and stress tolerance [31]. By harnessing the beneficial functions of phytobiomes, farmers can improve soil health and fertility, reduce dependence on chemical fertilizers and pesticides, and promote environmentally friendly farming practices [32]. Additionally, phytobiomes contribute to the development of crops that are more resilient to abiotic stresses like drought and salinity, as well as biotic challenges such as pathogens and pests. Integrating phytobiome management into agricultural systems not only boosts crop yields and quality but also supports sustainable agriculture by fostering ecosystem stability and biodiversity. As the agricultural sector faces increasing pressures from climate change and the need for sustainable food production, leveraging the power of phytobiomes offers a promising pathway to achieve more efficient, resilient, and ecologically balanced farming practices [33].

## 5. COMPONENTS OF PHYTOBIOMES

Phytobiomes are complex ecosystems composed of various biological and environmental components that interact with plants to influence their health, growth, and resilience. The key components of phytobiomes include:

### 5.1 Plants

The central component of the phytobiome, plants serve as the primary habitat and host for various microorganisms. They influence the composition and activity of microbial communities through the release of root exudates, leaf exudates, and other organic compounds that attract or repel specific microbes [34].

### 5.2 Microorganisms

Microbial communities within phytobiomes consist of diverse groups of organisms, each playing specific roles:

- **Bacteria:** These are abundant in the rhizosphere (soil surrounding roots), phyllosphere (above-ground parts), and endosphere (inside plant tissues). They contribute to nutrient cycling, disease

suppression, and plant growth promotion [35].

- **Fungi:** Fungi, including mycorrhiza fungi, are essential for nutrient acquisition, particularly phosphorus, and play roles in disease resistance and stress tolerance [36].
- **Archaea:** Though less studied, archaea are involved in nutrient cycling, particularly in extreme environments where they contribute to nitrogen and carbon cycling.
- **Viruses:** Plant-associated viruses can impact both the plant and microbial communities, sometimes in beneficial ways by controlling pathogen populations [37].
- **Protista and Nematodes:** These are part of the soil food web and influence microbial populations by preying on bacteria and fungi, thus regulating microbial dynamics [38].

### 5.3 Soil

Soil serves as the foundation for the phytobiome, providing a medium for root growth and a habitat for a vast array of microorganisms. Soil properties such as texture, structure, pH, organic matter content, and moisture levels directly influence the composition and function of microbial communities within the phytobiome [39].

### 5.4 Rhizosphere

The rhizosphere is the narrow region of soil directly influenced by root exudates and associated microbial activity. It is a hotspot for plant-microbe interactions, where beneficial microbes like nitrogen-fixing bacteria and mycorrhiza fungi thrive, contributing to nutrient acquisition and plant health [40].

### 5.5 Phyllosphere

The phyllosphere refers to the above-ground parts of the plant, including leaves, stems, and flowers, which host microbial communities adapted to more variable and sometimes harsh environmental conditions. These microbes can protect plants from foliar pathogens and contribute to nutrient cycling and overall plant health [41].

### 5.6 Endosphere

The endosphere consists of the interior tissues of the plant, where endophytic microorganisms

reside. These microbes can promote plant growth, enhance stress tolerance, and protect against pathogens by producing antimicrobial compounds or triggering plant defence mechanisms [42].

### 5.7 Abiotic Factors

Environmental factors such as temperature, humidity, light, and water availability significantly impact the composition and function of phytobiomes. Abiotic stresses like drought, salinity, and extreme temperatures can alter microbial communities, sometimes leading to shifts that either harm or benefit plant health [43].

### 5.8 Plant Exudates

Plants release a variety of organic compounds through their roots (root exudates) and leaves (leaf exudates) that serve as food sources or signalling molecules for microbes. These exudates shape the microbial community structure and function, influencing which microbes thrive and how they interact with the plant [44].

### 5.9 Pests and Pathogens

Pests and pathogens, including insects, fungi, bacteria, and viruses, are also part of the phytobiome. Their interactions with plants and other microbes can lead to diseases or trigger defence mechanisms that shape the overall health and resilience of the plant [45].

### 5.10 Agricultural Practices

Human interventions, such as the use of fertilizers, pesticides, crop rotation, and tillage, influence the phytobiome by altering the soil environment and microbial communities. Sustainable practices that support phytobiome diversity and health are increasingly recognized as critical for maintaining productive and resilient agricultural systems [46].

## 6. INTERACTIONS BETWEEN PLANTS AND MICROBIAL COMMUNITIES

Interactions between plants and microbial communities are complex and multifaceted, forming the foundation of plant health, growth, and resilience. These interactions can be mutualistic, commensal, or antagonistic, and

they occur in various plant-associated environments, such as the rhizosphere, phyllosphere, and endosphere. The key interactions between plants and microbial communities include:

### 6.1 Nutrient Exchange

One of the most important interactions is the exchange of nutrients between plants and microbes:

- **Nitrogen Fixation:** Certain bacteria, such as *Rhizobium* in legumes, form symbiotic relationships with plants by fixing atmospheric nitrogen into a form that plants can use. In return, the plants provide the bacteria with carbohydrates produced through photosynthesis [47].
- **Mycorrhiza Symbiosis:** Mycorrhiza fungi form associations with plant roots, extending their hyphae into the soil to increase the surface area for nutrient absorption, particularly phosphorus. In exchange, the fungi receive sugars and other organic compounds from the plant [48].
- **Phosphate Solubilisation:** Some soil bacteria and fungi solubilize phosphorus from insoluble compounds in the soil, making it available to plants. This interaction enhances plant nutrient uptake and [49].

### 6.2 Promotion of Plant Growth

Microbes promote plant growth through various mechanisms:

- **Production of Plant Hormones:** Certain microbes produce hormones such as auxins, cytokine's, and gibberellins, which stimulate plant growth by enhancing root development, shoot elongation, and overall plant vigour [50].
- **Improvement of Root Architecture:** Beneficial microbes can influence root morphology, increasing root branching and length, which enhances the plant's ability to absorb water and nutrients [51].
- **Enhanced Stress Tolerance:** Microbes can help plants tolerate abiotic stresses like drought, salinity, and temperature extremes by producing stress-related compounds, such as osmoprotectants, or

by altering the plant's stress response pathways [52].

### 6.3 Disease Suppression

Microbial communities play a crucial role in protecting plants from diseases:

- **Antagonism against Pathogens:** Beneficial microbes can suppress plant pathogens through competition for space and resources, production of antimicrobial compounds, or direct parasitism of the pathogen [53].
- **Induced Systemic Resistance (ISR):** Certain beneficial microbes trigger a plant-wide immune response known as ISR, which enhances the plant's ability to resist a broad spectrum of pathogens. This interaction primes the plant's defence mechanisms, making it more resistant to future infections [54].
- **Biocontrol:** Microbial biocontrol agents, such as *Trichoderma* and *Pseudomonas*, are used to control soil-borne and foliar pathogens, reducing the need for chemical pesticides [55].

### 6.4 Signalling and Communication

Plants and microbes engage in intricate communication through signalling molecules:

- **Root Exudates:** Plants secrete a variety of organic compounds through their roots, including sugars, amino acids, and secondary metabolites, which attract or repel specific microbes. These exudates can shape the composition of the microbial community in the rhizosphere [56].
- **Quorum Sensing:** Microbes use quorum sensing, a form of cell-to-cell communication, to coordinate their activities in response to the density of their population. This can influence microbial behaviour, such as biofilm formation, virulence factor production, and the activation of beneficial traits [57].
- **Symbiotic Signalling:** In mutualistic relationships, such as those between legumes and *Rhizobium* bacteria, plants and microbes exchange specific signals to establish and maintain symbiosis. For example, flavonoids released by plant roots induce the expression of nod genes in *Rhizobium*, leading to the formation of nitrogen-fixing nodules [58].

### 6.5 Competition and Antagonism

While many plant-microbe interactions are beneficial, some are competitive or antagonistic:

- **Pathogen Attack:** Pathogenic microbes can invade plant tissues, causing diseases that lead to reduced growth, yield, and even plant death. Plants respond by activating their immune systems to combat the pathogens [59].
- **Allelopathy:** Some plants release chemicals that inhibit the growth of neighbouring plants and their associated microbial communities. This can alter the microbial dynamics in the soil, often leading to competitive exclusion of certain microbes [60].
- **Microbial Warfare:** Within the microbial community, microbes can compete for resources and space, sometimes producing toxins or antibiotics that suppress the growth of competing microbes [60].

### 6.6 Endophyte Interactions

Endophytic microbes, which live inside plant tissues without causing harm, can interact with their host plants in various ways:

- **Growth Promotion:** Endophytes can promote plant growth by producing growth hormones, enhancing nutrient uptake, or helping plants cope with stress [61].
- **Pathogen Resistance:** Some endophytes produce antimicrobial compounds or stimulate plant defences, protecting the plant from pathogens [62].
- **Stress Alleviation:** Endophytes can enhance a plant's tolerance to environmental stresses by modulating stress-related pathways or producing protective compounds [63].

## 7. ENVIRONMENTAL FACTORS INFLUENCING PHYTOBIOMES

Environmental factors play a critical role in shaping the composition, diversity, and functionality of phytobiomes. These factors influence the interactions between plants and microbial communities, ultimately affecting plant health, growth, and resilience. The key environmental factors influencing phytobiomes include:

## 7.1 Soil Composition and Structure

- **Soil Type:** The physical and chemical properties of soil, such as texture (clay, silt, sand), pH, and organic matter content, greatly influence the microbial communities within the phytobiome. Different soil types harbour distinct microbial populations, which in turn affect nutrient availability and plant health [64].
- **Nutrient Availability:** The levels of essential nutrients like nitrogen, phosphorus, and potassium in the soil influence the microbial community structure. For example, nutrient-poor soils may promote the growth of microbes that specialize in nutrient cycling or fixation, which can benefit plants in nutrient-deficient environments [65].
- **Soil Moisture:** The amount of water in the soil affects microbial activity and diversity. Microbes require water for metabolic processes, and soil moisture levels can influence the oxygen availability and the balance between aerobic and anaerobic microorganisms [66].

## 7.2 Climate and Weather Conditions

- **Temperature:** Temperature is a key determinant of microbial metabolism and community composition. Extremes in temperature, whether hot or cold, can stress microbial populations and alter their interactions with plants. Some microbes thrive in specific temperature ranges, influencing their presence and activity in different climatic zones [67].
- **Precipitation and Humidity:** Water availability, through rainfall or irrigation, affects both the plant and the microbial communities in the soil and on plant surfaces. High humidity levels can promote the growth of certain fungi and bacteria, while low humidity may limit microbial activity and alter the composition of the phytobiome [68].
- **Seasonal Variation:** Changes in temperature, light, and moisture across seasons can lead to shifts in microbial community composition and activity. For example, certain microbes may become more active in the growing season, while others dominate in the off-season, affecting plant health throughout the year [69].

## 7.3 Light Availability

- **Photosynthetically Active Radiation (PAR):** Light influences plant growth and the production of root exudates, which in turn affect the microbial communities in the rhizosphere. Different light conditions can lead to changes in the types and amounts of exudates released by plants, thereby shaping the microbial population and its functions [70].
- **Photoperiod:** The length of day and night can impact the growth cycles of plants and their associated microbes. Some microbial processes, such as those involved in nitrogen fixation or decomposition, are influenced by the photoperiod, which affects the timing and intensity of microbial activity [71].

## 7.4 PH Levels

- **Soil pH:** The acidity or alkalinity of the soil is a major factor determining microbial diversity and function. Most microbes have an optimal pH range, and deviations from this range can suppress certain microbial groups while promoting others. For example, acidic soils may favour the growth of fungi, while alkaline soils may support different bacterial populations [72].
- **pH and Nutrient Availability:** Soil pH influences the solubility of nutrients, thereby affecting their availability to plants and microbes. For instance, at low pH, some essential nutrients like phosphorus become less available, influencing the types of microbes that can thrive in such conditions [73].

## 7.5 Oxygen Availability

- **Aeration:** The availability of oxygen in the soil, often related to soil compaction and water content, influences the types of microbes present. Aerobic microbes require oxygen for their metabolic processes, while anaerobic microbes thrive in low-oxygen environments, such as waterlogged soils [74].
- **Waterlogged Conditions:** Excess water can create anaerobic conditions, promoting the growth of microbes that thrive in low-oxygen environments, such as certain bacteria and fungi. These conditions can lead to changes in nutrient



cycling and the production of gases like methane, which can impact plant health [75].

## 7.6 Agricultural Practices

- **Fertilization:** The use of chemical fertilizers can alter the nutrient balance in the soil, influencing microbial diversity and activity. Over-fertilization, in particular, can lead to nutrient imbalances that favour certain microbes over others, potentially disrupting the natural balance of the phytobiome [76].
- **Pesticides and Herbicides:** The application of chemical pesticides and herbicides can have direct and indirect effects on microbial communities. While these chemicals target pests and weeds, they can also harm beneficial microbes or disrupt microbial interactions, leading to reduced plant health [77].
- **Tillage:** Tillage practices influence soil structure, aeration, and microbial habitat. Conventional tillage can disrupt microbial communities by breaking up soil aggregates and exposing deeper soil layers to the surface, while no-till or reduced-till practices help maintain a more stable and diverse microbial community [78].
- **Crop Rotation and Diversity:** The diversity of crops grown in a particular area influences the diversity of microbial communities. Crop rotation and polyculture practices promote a more diverse and resilient phytobiome by providing a range of habitats and nutrient sources for different microbes [79].

## 7.7 Pollution and Contaminants

- **Heavy Metals and Toxins:** Soil contamination with heavy metals, such as cadmium, lead, and mercury, or organic pollutants can negatively affect microbial communities by inhibiting microbial growth and activity. These contaminants can lead to reduced microbial diversity and function, impacting plant health and soil fertility [80].
- **Industrial and Agricultural Runoff:** Runoff containing chemicals from industrial or agricultural sources, such as fertilizers, pesticides, and pollutants, can alter the microbial balance in soils and water bodies, potentially leading to the

proliferation of harmful microbes and a decline in beneficial ones [81].

## 7.8 Biotic Interactions

- **Plant Species and Genotype:** Different plant species and even different genotypes of the same species can host distinct microbial communities. The specific traits of a plant, such as root architecture, exudate composition, and immune response, influence the types of microbes that associate with it [82].
- **Insect and Pest Activity:** The presence of insects and pests can influence the phytobiome by directly interacting with the plant or by affecting microbial communities. For example, pest damage can alter root exudate profiles, leading to changes in the rhizosphere microbiome [83].

## 8. METHODS FOR STUDYING PHYTOBIOMES

Studying phytobiomes requires a multidisciplinary approach that integrates various methods and technologies to understand the complex interactions between plants and their associated microbial communities. The following are some of the key methods used to study phytobiomes:

### 8.1 Culture-Dependent Techniques

- **Isolation and Cultivation:** Microorganisms from phytobiomes can be isolated and cultured on specific media to study their characteristics, such as morphology, growth conditions, and metabolic capabilities. These methods allow for the identification and characterization of cultivable microbes but are limited to those microbes that can grow in artificial conditions [84].
- **Biochemical Tests:** Cultured microbes can be subjected to various biochemical assays to determine their enzymatic activities, nutrient requirements, and potential interactions with plants. This information helps in understanding their role within the phytobiome [85].

### 8.2 Culture-Independent Techniques

- **DNA Sequencing:** High-throughput DNA sequencing technologies, such as next-

generation sequencing (NGS), are widely used to analyse the genetic material of entire microbial communities without the need for cultivation. This includes:

- **16S rRNA Gene Sequencing:** Used to identify and classify bacterial and archaeal communities based on the highly conserved 16S ribosomal RNA gene [86].
- **Internal Transcribed Spacer (ITS) Sequencing:** Used for identifying and studying fungal communities by targeting the ITS regions of fungal rRNA genes.
- **Metagenomics:** Involves sequencing the entire genetic material from a sample, allowing for the study of the functional potential of microbial communities by analysing genes associated with specific metabolic pathways [87].
- **Metatranscriptomics:** This approach involves sequencing RNA transcripts from microbial communities to study active gene expression. It provides insights into the functional activity of microbes within the phytobiome and how they respond to environmental changes or plant signals [88].
- **Metaproteomics:** Metaproteomics involves the analysis of the complete set of proteins produced by microbial communities in the phytobiome. This method helps in understanding the functional roles of microbes by identifying and quantifying proteins involved in various metabolic processes and interactions with plants [89].
- **Metabolomics:** Metabolomics studies the complete set of metabolites produced by the phytobiome, including small molecules like amino acids, sugars, organic acids, and secondary metabolites. It provides insights into the biochemical interactions between plants and microbes, including nutrient exchange and signalling [90].

### 8.3 Microscopy and Imaging Techniques

- **Fluorescence Microscopy:** Fluorescent dyes or genetically encoded fluorescent proteins can be used to visualize specific microbial cells or plant tissues under a fluorescence microscope. This technique allows for the study of spatial relationships between microbes and plant cells [91].
- **Confocal Laser Scanning Microscopy (CLSM):** CLSM provides high-resolution, three-dimensional images of microbial communities and their interactions with

plant tissues. It is particularly useful for studying the localization and distribution of microbes within the rhizosphere, phyllosphere, or endosphere [92].

- **Scanning Electron Microscopy (SEM) and Transmission Electron Microscopy (TEM):** These techniques provide detailed images of microbial morphology and plant-microbe interactions at the ultrastructural level [93].

### 8.4 Stable Isotope Probing (SIP)

Stable isotope probing involves incorporating stable isotopes, such as  $^{13}\text{C}$  or  $^{15}\text{N}$ , into specific substrates (e.g., carbon or nitrogen sources) provided to plants. Microbes that utilize these labelled substrates can be identified and studied by tracking the incorporation of isotopes into their DNA, RNA, or proteins. SIP is a powerful method for linking microbial activity to specific metabolic processes within the phytobiome.

### 8.5 Network Analysis and Bioinformatics

- **Co-occurrence Networks:** Network analysis involves constructing co-occurrence networks based on the presence and abundance of different microbial taxa in phytobiome samples. These networks help identify key microbial interactions and potential keystone species that play central roles in the community [93].
- **Functional Prediction:** Bioinformatics tools can be used to predict the functional capabilities of microbial communities based on metagenomics or 16S rRNA gene data. Tools like PICRUSt (Phylogenetic Investigation of Communities by Reconstruction of Unobserved States) predict the presence of functional genes in microbial communities [94].
- **Machine Learning:** Machine learning algorithms are increasingly used to analyse large and complex phytobiome datasets. These methods can identify patterns, predict microbial functions, and provide insights into the relationships between environmental factors, microbial communities, and plant health [95].

### 8.6 Experimental Manipulation

- **Inoculation Studies:** Experimental inoculation of plants with specific microbes

or microbial consortia allows researchers to study the effects of these microbes on plant growth, health, and resistance to stress or pathogens. These studies can be conducted in controlled environments, such as growth chambers or greenhouses, or in field conditions [96].

- **Gnotobiotic Systems:** Gnotobiotic (germ-free) systems involve growing plants in sterile conditions and then introducing specific microbial communities to study their interactions with the plant. This controlled approach helps isolate the effects of individual microbes or microbial consortia on plant development and health [97].
- **Microcosm and Macrocosm Experiments:** These are controlled,

simplified ecosystems used to study phytobiomes under various environmental conditions. Microcosms are small-scale experimental setups, while mesocosms are larger and more complex, allowing for the study of interactions between plants, microbes, and environmental factors [98].

### 8.7 Omics Integration

Integrating data from multiple "omics" approaches, such as genomics, transcriptomic, proteomics, and metabolomics, provides a comprehensive understanding of the phytobiome. This systems biology approach helps in unravelling the complex interactions and feedback mechanisms between plants and their associated microbial communities [99].

**Table 1. How phytobiomes, through various microbial communities, benefit specific crops**

Crop	Microbial Community	Role in Plant Health	Phytobiome Impact
<b>Wheat</b>	Fungi (Mycorrhizae)	Enhances nutrient uptake, especially phosphorus	Improves drought tolerance and growth in poor soils
<b>Rice</b>	Bacteria (Rhizobia and <i>Azospirillum</i> )	Nitrogen fixation, enhances root development	Increases nitrogen availability, boosting crop yield
<b>Tomato</b>	Fungi ( <i>Trichoderma</i> ) and Bacteria ( <i>Bacillus</i> )	Biocontrol of soil-borne pathogens and disease suppression	Reduces need for chemical fungicides, improves growth
<b>Maize</b>	Bacteria ( <i>Pseudomonas</i> )	Phytostimulation and production of plant growth-promoting substances	Enhances root biomass and nutrient absorption
<b>Soybean</b>	Bacteria ( <i>Bradyrhizobium</i> )	Nitrogen fixation, improves plant nitrogen content	Reduces need for synthetic nitrogen fertilizers
<b>Potato</b>	Fungi (Endophytic fungi)	Helps with disease resistance and drought tolerance	Enhances resistance to pathogens like <i>Phytophthora infestans</i>
<b>Sugarcane</b>	Bacteria ( <i>Gluconacetobacter diazotrophicus</i> )	Fixes atmospheric nitrogen, enhances growth	Reduces nitrogen input, increases biomass production
<b>Barley</b>	Viruses (Plant-associated viruses)	Can confer resistance to certain pathogenic viruses	Provides natural viral immunity
<b>Alfalfa</b>	Fungi (Arbuscular Mycorrhizal Fungi)	Improves phosphorus uptake, helps in drought tolerance	Promotes soil health and plant vigour
<b>Carrot</b>	Nematodes (Beneficial nematodes)	Controls root-feeding pests like root-knot nematodes	Reduces crop damage, decreases need for chemical nematicides

**Table 2. Harnessing Phytobiomes in Agriculture**

Aspect	Description	Agricultural Application	Impact on Plant Health
Microbiome Enhancement	Utilizing beneficial microbes like bacteria, fungi, and archaea.	Application of biofertilizers, mycorrhiza fungi, and probiotics.	Enhances nutrient uptake, promotes growth, and increases disease resistance.
Soil Health Management	Maintaining and improving soil physical, chemical, and biological properties.	Crop rotation, cover cropping, organic amendments.	Supports diverse microbial communities, improves fertility, and suppresses diseases.
Environmental Modulation	Adapting to and modifying the surrounding environment to favour phytobiomes.	Implementing irrigation management, shade systems, and windbreaks.	Mitigates environmental stress, stabilizes microclimates, and enhances microbial activity.
Plant-Microbe Interaction	Promoting beneficial plant-microbe symbioses.	Inoculation with rhizobia for nitrogen fixation, use of biopesticides.	Boosts plant immunity, facilitates nutrient exchange, and controls pests.
Phytobiome Engineering	Designing phytobiomes through genomic and metagenomic technologies.	Developing microbial inoculants, gene editing in microbes.	Tailors specific traits like stress tolerance and pathogen resistance.
Sustainable Farming Practices	Integrating phytobiome knowledge into farming systems.	Reduced chemical inputs, conservation tillage, agroforestry.	Enhances ecosystem services, reduces environmental impact, and improves long-term productivity.
Disease Suppression	Targeting pathogenic microbes within the phytobiome.	Use of biological control agents, disease-resistant cultivars.	Controls outbreaks, reduces reliance on chemical fungicides, and promotes overall plant health.

## 9. PHYTOBIOMES AND PLANT HEALTH

Phytobiomes are the complex systems that include the plant itself, its associated environment, and all the living organisms (microbiomes) that interact with it. This concept emphasizes the interconnectedness of plants with their surrounding environment, including soil, climate, and other organisms like bacteria, fungi, viruses, insects, and animals [100, 101]. Understanding phytobiomes is crucial for improving plant health, productivity, and resilience.

### Key Aspects of Phytobiomes and Plant Health:

1. **Microbiomes:** The microbial communities associated with plants, particularly in the rhizosphere (soil region near roots), phyllosphere (above-ground parts), and endosphere (internal tissues). These

microorganisms can promote plant growth, enhance nutrient uptake, and protect against pathogens.

2. **Soil Health:** Soil is a major component of the phytobiome, influencing plant health through its physical, chemical, and biological properties. Healthy soils support diverse microbial communities that contribute to nutrient cycling, organic matter decomposition, and disease suppression.
3. **Environmental Factors:** Climate, weather patterns, and geographical location affect phytobiomes. Changes in temperature, humidity, and other environmental conditions can alter microbial communities and their interactions with plants, impacting plant health.
4. **Plant-Microbe Interactions:** Beneficial microbes can enhance plant health by improving stress tolerance, increasing nutrient availability, and inducing

systemic resistance to pathogens. Conversely, pathogenic microbes can cause diseases that threaten plant health.

- 5. Phytobiome Management:** Sustainable agricultural practices, such as crop rotation, cover cropping, and the use of bio fertilizers or biopesticides, can promote healthy phytobiomes. Understanding the interactions within phytobiomes allows for better

management practices to enhance plant health and crop yield.

- 6. Research and Innovation:** Advances in genomics, metagenomics, and bioinformatics are enabling deeper insights into phytobiomes. Research is focused on identifying key microbial species and their functions, developing microbial inoculants, and engineering phytobiomes for specific agricultural outcomes.

**Table 3. Successful Applications of Phytobiome Management**

Application	Description	Crop/Region	Impact	Outcome
Mycorrhizal Fungi Inoculation	Introduction of beneficial mycorrhizal fungi to enhance nutrient uptake.	Wheat in North America	Improved phosphorus uptake and drought tolerance.	Increased yield and reduced fertilizer dependency.
Rhizobia Inoculation	Use of nitrogen-fixing bacteria to enhance soil fertility.	Legumes (e.g., soybeans) in Africa	Enhanced nitrogen fixation, leading to better plant growth.	Improved crop yields and reduced need for chemical fertilizers.
Biocontrol Agents	Application of microbes that suppress plant pathogens.	Tomatoes in Europe	Controlled soil-borne diseases like Fusarium wilt.	Reduced crop losses and minimized use of chemical fungicides.
Cover Cropping	Use of cover crops to promote beneficial soil microbes.	Corn in the United States	Enhanced soil health and microbial diversity.	Improved soil structure, increased organic matter, and better water retention.
Compost and Organic Amendments	Application of compost to boost beneficial soil microbiomes.	Vineyards in California	Increased microbial activity and nutrient cycling.	Improved grape quality and reduced synthetic input usage.
Endophyte Inoculation	Introducing beneficial endophytes to promote plant growth.	Rice in Asia	Improved growth under stress conditions (e.g., salinity).	Increased resilience to environmental stresses and higher yields.
Probiotic Applications	Application of beneficial bacteria to the phyllosphere.	Leafy greens in Greenhouses	Enhanced resistance to foliar diseases.	Reduced disease incidence and higher-quality produce.
Biochar Addition	Incorporation of biochar to support microbial communities.	Various crops in degraded soils (Global)	Enhanced soil microbial diversity and stability.	Improved soil health and increased crop productivity.
Agroforestry Practices	Integrating trees with crops to enhance phytobiomes.	Coffee plantations in Central America	Enhanced soil organic matter and microbial interactions.	Improved crop resilience and sustainable land use.
Biostimulant Use	Application of microbial biostimulants to enhance plant growth.	Grapes in Europe	Improved nutrient uptake and stress tolerance.	Enhanced grape yield and quality, reduced chemical input needs.

## 10. CONCLUSION

Phytobiomes represent a powerful tool for enhancing plant health and agricultural sustainability. By harnessing the complex interactions between plants and their associated microbial communities, we can develop innovative strategies for improving crop productivity, reducing the environmental impact of farming, and addressing the challenges of climate change. The success stories and challenges outlined in this article highlight the immense potential of phytobiomes, as well as the need for continued research, innovation, and collaboration. As we move towards a more sustainable agricultural future, phytobiomes will undoubtedly play a central role in shaping the health and resilience of crops, ensuring food security, and protecting the environment. In conclusion, the study and application of phytobiomes offer a promising path forward for modern agriculture. By embracing the power of microbial communities, we can build more resilient and sustainable farming systems that benefit both people and the planet. The future of agriculture lies not only in the genetic makeup of crops but also in the dynamic and diverse microbial communities that support them.

## DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Agus A, Planchais J, Sokol H. Gut microbiota regulation of tryptophan metabolism in health and disease. *Cell Host Microbe*. 2018;23:716–724.
2. Arpan Mukherjee, Bansh Narayan Singh, Simranjit Kaur, Minaxi Sharma, Ademir Sérgio Ferreira de Araújo, Arthur Prudêncio de Araujo Pereira, Raj Morya, Gerardo Puopolo, Vânia Maria Maciel Melo, Jay Prakash Verma, Unearthing the power of microbes as plant microbiome for sustainable agriculture, *Microbiological Research*. 2024;286:127780, ISSN 0944-5013.

- Available: <https://doi.org/10.1016/j.micres.2024.127780>
3. Alabouvette C. Fusarium-wilt suppressive soils from the Châteaurenard region: review of a 10-year study. *Agronomie*. 1986;6:273–284.
  4. Alabouvette C. Fusarium wilt suppressive soils: an example of disease-suppressive soils. *Australas. Plant Pathol.* 1999;28:57–64.
  5. Anthony MA, Crowther TW, Maynard DS, van den Hoogen J, Averill C. Distinct assembly processes and microbial communities constrain soil organic carbon formation. *One Earth*. 2020;2:349–360.
  6. Babalola OO. Beneficial bacteria of agricultural importance. *Biotechnol. Lett.* 2010;32:1559–1570.
  7. Bai Y, Müller DB, Srinivas G, Garrido-Oter R, Potthoff E, Rott M, Dombrowski N, Münch PC, Spaepen S, Remus-Emsermann M. Functional overlap of the *Arabidopsis* leaf and root microbiota. *Nature*. 2015;528:364–369.
  8. Bais HP, Weir TL, Perry LG, Gilroy S, Vivanco JM. The role of root exudates in rhizosphere interactions with plants and other organisms. *Annu. Rev. Plant Biol.* 2006;57:233–266.
  9. Bakker PAHM, Pieterse CMJ, De Jonge R, Berendsen RL. The soil-borne legacy. *Cell*. 2018;172:1178–1180.
  10. Berendsen RL, Vismans G, Yu K, Song Y, De Jonge R, Burgman WP, Burmolle M, Herschend J, Bakker PAHM, Pieterse CMJ. Disease-induced assemblage of a plant-beneficial bacterial consortium. *ISME J*. 2018;12:1496–1507.
  11. Biliouris K, Babson D, Schmidtdannert C, Kaznessis YN. Stochastic simulations of a synthetic bacteria-yeast ecosystem. *BMC Syst. Biol.* 2012;6:58.
  12. Broadbent P, Baker KF. Behaviour of *Phytophthora cinnamomi* in soils suppressive and conducive to root rot. *Crop Pasture Sci.* 1974;25:121–137.
  13. Bulgarelli D, Schlaeppi K, Spaepen S, Ver Loren van Themaat E, Schulze-Lefert P. Structure and functions of the bacterial microbiota of plants. *Annu. Rev. Plant Biol.* 2013;64:807–838.
  14. Cadot S, Guan H, Bigalke M, Walsler JC, Schlaeppi K. Specific and conserved patterns of microbiota-structuring by maize benzoxazinoids in the field. *Microbiome*. 2020;9:103.

15. Carrion VJ, Cordovez V, Tyc O, Etalo DW, De Bruijn I, De Jager V, Medema MH, Eberl L, Raaijmakers JM. Involvement of Burkholderiaceae and sulfurous volatiles in disease-suppressive soils. *ISME J.* 2018;12:2307–2321.
16. Chen Q-L, Hu H-W, He Z-Y, Cui L, Zhu Y-G, He J-Z. Potential of indigenous crop microbiomes for sustainable agriculture. *Nat. Food.* 2021;2:233–240.
17. Chiaramonte JB, Mendes LW, Mendes R. Rhizosphere microbiome and soil-borne diseases. *Rhizosphere Biol. Interactions Between Microbes Plants.* 2021:155–168.
18. Chilosi G, Aleandri MP, Luccioli E, Stazi SR, Marabottini R, Morales-Rodríguez C, Vettrano AM, Vannini A. Suppression of soil-borne plant pathogens in growing media amended with espresso spent coffee grounds as a carrier of *Trichoderma spp.* *Scientia Horticulturae.* 2020;259:108666.
19. Cira N, Pearce M, Quake S. Neutral and selective dynamics in a synthetic microbial community. *Proc. Natl. Acad. Sci.* 2018;115:201808118.
20. Compant S, Samad A, Faist H, Sessitsch A. A review on the plant microbiome: ecology, functions, and emerging trends in microbial application. *J. Adv. Res.* 2019;19:29–37.
21. Cook RJ, Rovira AD. The role of bacteria in the biological control of *Gaeumannomyces graminis* by suppressive soils. *Soil Biol. Biochem.* 1976;8:269–273.
22. Cordovez V, Dini-Andreote F, Carrión V, Raaijmakers JM. Ecology and evolution of plant microbiomes. *Annu. Rev. Microbiol.* 2019;73:69–88.
23. De Corato U, Patruno L, Avella N, Lacolla G, Cucci G. Composts from green sources show an increased suppressiveness to soilborne plant pathogenic fungi: relationships between physicochemical properties, disease suppression, and the microbiome. *Crop Prot.* 2019;124:104870.
24. Du J, Li Y, Yin Z, Wang H, Zhang X, Ding X. High-throughput customization of plant microbiomes for sustainable agriculture. *Front. Plant Sci.* 2020;11:1365.
25. Duan W, Shi Y, Zhao J, Zhang Y, Yu Z. Depth of nitrogen fertiliser placement affects nitrogen accumulation, translocation and nitrate-nitrogen content in soil of rainfed wheat. *Int. J. Plant Prod.* 2015;9:237–256.
26. Eichmann R, Richards L, Schäfer P. Hormones as go-betweens in plant microbiome assembly. *Plant J.* 2021;105:518–541.
27. Fahad S, Hussain S, Bano A, Saud S, Hassan S, Shan D, Khan FA, Khan F, Chen Y, Wu C. Potential role of phytohormones and plant growth-promoting rhizobacteria in abiotic stresses: consequences for changing environment. *Environ. Sci. Pollut. Res.* 2015;22:4907–4921. DOI: 10.1007/s11356-014-3754-2
28. Fiorilli V, Vannini C, Ortolani F, Garcia-Secco D, Chiapello M, Novero M, Domingo G, Terzi V, Morcia C, Bagnaresi P. Omics approaches revealed how arbuscular mycorrhizal symbiosis enhances yield and resistance to leaf pathogen in wheat. *Sci. Rep.* 2018;8:9625. DOI: 10.1038/s41598-018-27622-8.
29. Fitzpatrick CR, Salas-González I, Conway JM, Finkel OM, Gilbert S, Russ D, Teixeira PJL, Dangl JL. The plant microbiome: from ecology to reductionism and beyond. *Annu. Rev. Microbiol.* 2020;74:81–100.
30. Goldford JE, Lu N, Bajic D, Estrela S, Tikhonov M, Sanchez-Gorostiaga A, Segre D, Mehta P, Sanchez A. Emergent simplicity in microbial community assembly. *Science.* 2018;361:469–474.
31. Großkopf T, Soyer OS. Synthetic microbial communities. *Curr. Opin. Microbiol.* 2014;18:72–77.
32. Haas D, Défago G. Biological control of soil-borne pathogens by fluorescent pseudomonads. *Nat. Rev. Microbiol.* 2005;3:307–319.
33. Hacquard S, Garrido-Oter R, González A, Spaepen S, Ackermann G, Lebeis S, McHardy Alice C, Dangl Jeffrey L, Knight R, Ley R, Schulze-Lefert P. Microbiota and host nutrition across plant and animal kingdoms. *Cell Host Microbe.* 2015;17:603–616.
34. Hadar Y, Papadopoulou KK. Suppressive composts: microbial ecology links between abiotic environments and healthy plants. *Annu. Rev. Phytopathol.* 2012;50:133–153.
35. Harbort CJ, Hashimoto M, Inoue H, Niu Y, Guan R, Rombolà AD, Kopriva S, Voges MJEEE, Sattely ES, Garrido-Oter R, Schulze-Lefert P. Root-secreted coumarins and the microbiota interact to improve iron nutrition in Arabidopsis. *Cell Host Microbe.* 2020;28:825–837.

36. Jaiswal A, Elad Y, Paudel I, Graber E, Cytryn E, Frenkel O. Linking the belowground microbial composition, diversity and activity to soilborne disease suppression and growth promotion of tomato amended with biochar. *Sci. Rep.* 2017;7:44382.
37. Jambhulkar PP, Sharma M, Lakshman D, Sharma P. Natural mechanisms of soil suppressiveness against diseases caused by *Fusarium*, *Rhizoctonia*, *Pythium*, and *Phytophthora*. In: Meghvansi MK, Varma A, editors. *Organic Amendments and Soil Suppressiveness in Plant Disease Management*. Springer; 2015;95–124.
38. Kaminsky LM, Trexler RV, Malik RJ, Hockett KL, Bell TH. The inherent conflicts in developing soil microbial inoculants. *Trends Biotechnol.* 2019;37:140–151.
39. Kessler D, Bhattacharya S, Diezel C, Rothe E, Gase K, Schottner M, Baldwin IT. Unpredictability of nectar nicotine promotes outcrossing by hummingbirds in *Nicotiana attenuata*. *Plant J.* 2012;71:529–538.
40. Keswani C, Dilnashin H, Birla H, Singh S. Unravelling efficient applications of agriculturally important microorganisms for alleviation of induced inter-cellular oxidative stress in crops. *Acta agriculturae Slovenica.* 2019;114:121–130.
41. Kong HG, Song GC, Sim H-J, Ryu C-M. Achieving similar root microbiota composition in neighbouring plants through airborne signalling. *ISME J.* 2021;15:397–408.
42. Kuzakov Y, Domanski G. Carbon input by plants into the soil. *Review. J. Plant Nutr. Soil Sci.* 2000;163:421–431.
43. Kwak M-J, Kong HG, Choi K, Kwon S-K, Song JY, Lee J, Lee PA, Choi SY, Seo M, Lee HJ. Rhizosphere microbiome structure alters to enable wilt resistance in tomato. *Nat. Biotechnol.* 2018;36:1100–1109.
44. Ladau J, Elie-Fadrosh EA. Spatial, temporal, and phylogenetic scales of microbial ecology. *Trends Microbiol.* 2019;27:662–669.
45. Latz E, Eisenhauer N, Rall BC, Scheu S, Jousset A. Unravelling linkages between plant community composition and the pathogen-suppressive potential of soils. *Sci. Rep.* 2016;6:23584.
46. Lawson CE, Harcombe WR, Hatzenpichler R, Lindemann SR, Löffler FE, O'Malley MA, García Martín H, Pflieger BF, Raskin L, Venturelli OS. Common principles and best practices for engineering microbiomes. *Nat. Rev. Microbiol.* 2019;17:725–741.
47. Liu H, Chang Q, Feng W, Zhang B, Wu T, Li N, Yao F, Ding X, Chu Z. Domain dissection of AvrRxo1 for suppressor, Avirulence and cytotoxicity functions. *PLoS One.* 2014;9:e113875.
48. Lu C, Liu H, Jiang D, Wang L, Jiang Y, Tang S, Hou X, Han X, Liu Z, Zhang M. *Paecilomyces variotii* extracts (ZNC) enhance plant immunity and promote plant growth. *Plant and Soil.* 2019;441:383–397.
49. Luo LF, Yang L, Yan ZX, Jiang BB, Li S, Huang HC, Liu YX, Zhu SS, Yang M. Ginsenosides in root exudates of *Panax notoginseng* drive the change of soil microbiota through carbon source different utilization. *Plant and Soil.* 2020;455:139–153.
50. Maarastawi SA, Frindte K, Bodelier PLE, Knief C. Rice straw serves as additional carbon source for rhizosphere microorganisms and reduces root exudate consumption. *Soil Biol. Biochem.* 2019;135:235–238.
51. Martins S, De Medeiros FHV, De Souza RM, De Resende MLV, Ribeiro PM. Biological control of bacterial wilt of common bean by plant growth-promoting rhizobacteria. *Biol. Control.* 2013;66:65–71.
52. Marx V. Engineers embrace microbiome messiness. *Nat. Methods.* 2019;16:581–584.
53. Mastan A, Bharadwaj RKB, Kushwaha RK, Vivek Babu CS. Functional fungal endophytes in *coleus forskohlii* regulate labdane diterpene biosynthesis for elevated forskolin accumulation in roots. *Microb. Ecol.* 2019;78:914–926.
54. Mendes R, Kruijt M, De Bruijn I, Dekkers E, Der Voort MV, Schneider J, Piceno YM, Desantis TZ, Andersen GL, Bakker PAHM. Deciphering the rhizosphere microbiome for disease-suppressive bacteria. *Science.* 2011;332:1097–1100.
55. Molina-Romero D, Baez A, Quintero-Hernandez V, Castañeda M, Fuentes-Ramirez L, Bustillos M, Rodríguez-Andrade O, Morales-García Y, Munive A, Muñoz-Rojas J. Compatible bacterial mixture, tolerant to desiccation, improves maize plant growth. *PLoS One.* 2017;12:e0187913.
56. Morais CLM, Paraskevaidi M, Cui L, Fullwood NJ, Isabelle M, Lima KMG,



- Martin-Hirsch PL, Sreedhar H, Trevisan J, Walsh MJ. Standardization of complex biologically derived spectrochemical datasets. *Nat. Protoc.* 2019;14:1546–1577.
57. Nan Q, Wang C, Wang H, Yi Q, Liang B, Xu J, Wu W. Biochar drives microbially-mediated rice production by increasing soil carbon. *J. Hazard. Mater.* 2019;387:121680.
58. Neal AL, Ahmad S, Gordon-Weeks R, Ton J. Benzoxazinoids in root exudates of maize attract *Pseudomonas putida* to the rhizosphere. *PLoS One.* 2012;7:e35498.
59. Neeno-Eckwall E, Kinkel L, Schottel J. Competition and antibiosis in the biological control of potato scab. *Can. J. Microbiol.* 2001;47:332–340.
60. Noble R, Coventry E. Suppression of soil-borne plant diseases with composts: a review. *Biocontrol Sci.* 2005;15:3–20.
61. Pane C, Celano G, Piccolo A, Villecco D, Spaccini R, Palese A, Zaccardelli M. Effects of on-farm composted tomato residues on soil biological activity and yields in a tomato cropping system. *Chem. Biol. Tech. Agric.* 2015;24:2–13.
62. Pang Z, Chen J, Wang T, Gao C, Li Z, Guo L, Xu J, Cheng Y. Linking plant secondary metabolites and plant microbiomes: a review. *Front. Plant Sci.* 2021;12:621276.
63. Peng C, Zhang A, Wang Q, Song Y, Zhang M, Ding X, li Y, Geng Q, Zhu C. Ultrahigh-activity immune inducer from Endophytic Fungi induces tobacco resistance to virus by SA pathway and RNA silencing. *BMC Plant Biol.* 2020;20:169.
64. Pereg L, Mcmillan M. Scoping the potential uses of beneficial microorganisms for increasing productivity in cotton cropping systems. *Soil Biol. Biochem.* 2015;80:349–358.
65. Plichta DR, Graham DB, Subramanian S, Xavier RJ. Therapeutic opportunities in inflammatory bowel disease: mechanistic dissection of host-microbiome relationships. *Cell.* 2019;178:1041–1056.
66. Pollak S, Cordero OX. Rhizobiome shields plants from infection. *Nat. Microbiol.* 2020;5:978–979.
67. Qin Y, Druzhinina IS, Pan X, Yuan Z. Microbially mediated plant salt tolerance and microbiome-based solutions for saline agriculture. *Biotechnol. Adv.* 2016;34:1245–1259.
68. Rastall RA, Maitin V. Prebiotics and synbiotics: towards the next generation. *Curr. Opin. Biotechnol.* 2002;13:490–496.
69. Roberfroid M. Prebiotics: the concept revisited. *J. Nutr.* 2007;137:830S–837S.
70. Rodriguez PA, Rothballer M, Chowdhury SP, Nussbaumer T, Gutjahr C, Falter-Braun P. Systems biology of plant-microbiome interactions. *Mol. Plant.* 2019;12:804–821.
71. Ronda C, Chen SP, Cabral V, Yaung SJ, Wang HH. Metagenomic engineering of the mammalian gut microbiome in situ. *Nat. Methods.* 2019;16:167–170.
72. Rosenzweig N, Tiedje JM, Quensen JF, Meng Q, Hao J. Microbial communities associated with potato common scab-suppressive soil determined by pyrosequencing analyses. *Plant Dis.* 2012;96:718–725.
73. Santiago TR, Grabowski C, Rossato M, Romeiro RDS, Mizubuti E.S.G. Biological control of eucalyptus bacterial wilt with rhizobacteria. *Biol. Control.* 2015;80:14–22.
74. Schlatter DC, Kinkel LL, Thomashow LS, Weller DM, Paulitz TC. Disease suppressive soils: new insights from the soil microbiome. *Phytopathology.* 2017;107:1284–1297.
75. Singh JS, Gupta VK. Soil microbial biomass: a key soil driver in management of ecosystem functioning. *Sci. Total Environ.* 2018;634:497–500.
76. Singh JS, Pandey VC, Singh DP. Efficient soil microorganisms: a new dimension for sustainable agriculture and environmental development. *Agric. Ecosyst. Environ.* 2011;140:339–353.
77. Sintim HY, Bandopadhyay S, English ME, Bary AI, DeBruyn JM, Schaeffer SM, Miles CA, Reganold JP, Flury M. Impacts of biodegradable plastic mulches on soil health. *Agric. Ecosyst. Environ.* 2019;273:36–49.
78. Steffan JJ, Brevik EC, Burgess LC, Cerdà A. The effect of soil on human health: an overview. *Eur. J. Soil Sci.* 2018;69:159–171.
79. Stringlis IA, Yu K, Feussner K, de Jonge R, Van Bentum S, Van Verk MC, Berendsen RL, Bakker PAHM. MYB72-dependent coumarin exudation shapes root microbiome assembly to promote plant health. *Proc. Natl. Acad. Sci. U S A.* 2018;115:E5213–E5222.

80. Tao C, Rong L, Xiong W, Shen Z, Liu S, Wang B, Ruan Y, Geisen S, Shen Q, Kowalchuk G. Bio-organic fertilizers stimulate indigenous soil *Pseudomonas* populations to enhance plant disease suppression. *Microbiome*. 2020;8:137.
81. Thakur MP, Geisen S. Trophic regulations of the soil microbiome. *Trends Microbiol*. 2019;27:771–780.
82. Tian B, Pei Y, Huang W, Ding J, Siemann E. Increasing flavonoid concentrations in root exudates enhance associations between arbuscular mycorrhizal fungi and an invasive plant. *ISME J*. 2021;15:1919–1930.
83. Tian T, Sun B, Shi H, Gao T, He Y, Li Y, Liu Y, Li X, Zhang L, Li S. Sucrose triggers a novel signaling cascade promoting *Bacillus subtilis* rhizosphere colonization. *ISME J*. 2021
84. Trivedi P, Leach J, Tringe S, Sa T, Singh B. Plant–microbiome interactions: from community assembly to plant health. *Nat. Rev. Microbiol*. 2020;18:607–621.
85. Tu C, Wei J, Guan F, Liu Y, Sun Y, Luo Y. Biochar and bacteria inoculated biochar enhanced Cd and Cu immobilization and enzymatic activity in a polluted soil. *Environ. Int*. 2020;137:105576.
86. Wagg C, Bender SF, Widmer F, van der Heijden MGA. Soil biodiversity and soil community composition determine ecosystem multifunctionality. *Proc. Natl. Acad. Sci. U S A*. 2014;111:5266–5270.
87. Wang L, Li X. Steering soil microbiome to enhance soil system resilience. *Crit. Rev. Microbiol*. 2019;45:743–753.
88. Weller D, Raaijmakers J, Gardener B, Thomashow L. Weller DM, Raaijmakers JM, McSpadden Gardener BB, Thomashow LS.. Microbial populations responsible for specific soil suppressiveness to plant pathogens. *Annu. Rev. Phytopathol*. 2002;40:309–348.
89. Wen T, Zhao M, Yuan J, Kowalchuk GA, Shen Q. Root exudates mediate plant defense against foliar pathogens by recruiting beneficial microbes. *Soil Ecol. Lett*. 2021;3:42–51.
90. Xin XF, Nomura K, Ding X, Chen X, Wang K, Aung K, Uribe F, Rosa B, Yao J, Chen J, He S. *Pseudomonas syringae* effector avirulence protein E localizes to the host plasma membrane and down-regulates the expression of the nonrace-specific disease resistance1/harpin-like3 gene required for antibacterial immunity in *Arabidopsis*. *Plant Physiol*. 2015;169:793–802.
91. Xiong W, Rong L, Ren Y, Liu C, Zhao Q, Wu H, Jousset A, Shen Q. Distinct roles for soil fungal and bacterial communities associated with the suppression of vanilla *Fusarium* wilt disease. *Soil Biol. Biochem*. 2017;107:198–207.
92. Xu X, Wu B, Zhao W, Lao F, Wu J. Shifts in autochthonous microbial diversity and volatile metabolites during the fermentation of chili pepper (*Capsicum frutescens* L.) *Food Chem*. 2020;335:127512.
93. Yang C, Dong Y, Friman V-P, Jousset A, Wei Z, Xu Y, Shen Q. Carbon resource richness shapes bacterial competitive interactions by alleviating growth-antibiosis trade-off. *Funct. Ecol*. 2018;33:868–875.
94. Yang W, Dong R, Liu L, Hu Z, Li J, Wang Y, Ding X, Chu Z. A novel mutant allele of SSI2 confers a better balance between disease resistance and plant growth inhibition on *Arabidopsis thaliana*. *BMC Plant Biol*. 2016;16:208.
95. Yang W, Yin Y, Li Y, Cai T, Ni Y, Peng D, Wang Z. Interactions between polyamines and ethylene during grain filling in wheat grown under water deficit conditions. *Plant Growth Regul*. 2014;72:189–201.
96. Yuan J, Zhao J, Wen T, Zhao M, Li R, Goossens P, Huang Q, Bai Y, Vivanco JM, Kowalchuk GA. Root exudates drive the soil-borne legacy of aboveground pathogen infection. *Microbiome*. 2018;6:156.
97. Zaccardelli M, Perrone D, Pane C, Pucci N, Infantino A. Control of corky root of tomato with compost and role of spore-forming bacteria to inhibit *Pyrenochaeta lycopersici*. *Acta Hortic*. 2011;914:393–396.
98. Zengler K, Hofmockel K, Baliga NS, Behie SW, Bernstein HC, Brown JB, Dinneny JR, Flöge SA, Fory SP, Hess M. EcoFABs: advancing microbiome science through standardized fabricated ecosystems. *Nat. Methods*. 2019;16:567–571.
99. Zhainina K, Louie KB, Hao Z, Mansoori N, da Rocha UN, Shi S, Cho H, Karaoz U, Loqué D, Bowen BP. Dynamic root exudate chemistry and microbial substrate preferences drive patterns in rhizosphere microbial community assembly. *Nat. Microbiol*. 2018;3:470–480.

100. Zheng Y, Han X, Zhao D, Wei K, Yuan Y, Li Y, Liu M, Zhang CS. Exploring biocontrol agents from microbial keystone taxa associated to suppressive soil: a new attempt for a biocontrol strategy. *Front. Plant Sci.* 2021;12:655673.
101. Zhuang L, Li Y, Wang Z, Yu Y, Zhang N, Yang C, Zeng Q, Wang Q. Synthetic community with six *Pseudomonas* strains screened from garlic rhizosphere microbiome promotes plant growth. *Microb. Biotechnol.* 2020;14:488–502.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/122866>