



Safety of Weekly Primaquine in Glucose 6 Phosphatase Dehydrogenase (G6PD) Deficient Children

**Salma Shaikh ^a[≡], Muhammad Nadeem Chohan ^b^{⊖*}, Muhammad Touseef ^b[#],
Hafiz Wajid Ali Buriro ^b[†], Mushtaque Ali Shah ^c[‡][#] and Sehrish Muzaffar ^b[¥]**

^a Bilawal Medical College, Liaquat University of Medical and Health Sciences, Jamshoro, Pakistan.

^b Department of Pediatric, Liaquat University of Medical and Health Sciences, Pakistan.

^c Liaquat University of Medical and Health Sciences, Jamshoro, Pakistan.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i57B34080

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/78636>

Original Research Article

Received 06 October 2021

Accepted 12 December 2021

Published 14 December 2021

ABSTRACT

Aim: To assess the Safety of weekly Primaquine in Glucose 6 Phosphatase Dehydrogenase (G6PD) deficient children, for radical treatment of *Plasmodium vivax* malaria

Study Design: cross sectional study

Place and Duration: Pediatrics Out Patient Department, Liaquat University of Medical and Health Sciences Hyderabad from 11 January 2018 to 31st August 2019 (total 20 months' duration)

Methodology: A sample of 40 patients was studied during study period. Male children between 4 years to 12 years of age having confirmed vivax malaria were included in the study. If G6PD result showed decreased level of G6PD level then, they were enrolled for study. MP was checked by thick and thin slide method. 5 ml blood was taken in anticoagulant bottle for G6PD, liver function test, creatinine, complete blood count, and reticulocyte count tests. Haemoglobin < 7 g/dL,

[≡] Professor;

[⊖] Assistant Professor;

[#] Senior Registrar;

[†] Research Medical Officer;

[‡] Dr.;

[¥] Resident;

*Corresponding author: E-mail: nadeem.chohan@lumhs.edu.pk, nadeemchohanpink76@yahoo.com;

reticulocyte count > 4, SGPT > 80, G6PD Level < 60% of normal and creatinine > 1.2 was considered significant. Treatment was given with Artemether and Lumefantrine for 3 days while Primaquine, 0.75 mg base/kg body weights once a week was given for 8 weeks. Patients were followed at OPD initially on 3rd day of therapy then every week for 8 weeks for any hemolysis.

Results: There was no hemolysis during the first week and 8 weeks after therapy. Most common side effect was abdominal pain 4 (10%). Mean hemoglobin was 11.8mg/dl. *Plasmodium vivax* was negative on 3rd day of therapy, it was also negative on 8 week of therapy. Reticulocyte count, Liver function test, creatinine were also normal on 8 weeks of therapy.

Conclusion: Primaquine 0.75mg/kg/week for total eight weeks is highly effective for the radical cure of *Plasmodium vivax* in G6PD deficient children. There is no recurrence of *Plasmodium vivax* after 8 weeks of therapy. We found this regimen safe as there was no hemolysis demonstrated in children.

Keywords: Malaria; *Plasmodium vivax*; G6PD deficiency; hemolysis; radical cure.

1. INTRODUCTION

Malaria is almost eradicated from developed nations, but continues a health problem in a substantial part of the world. Malaria is a major health issue in Asian and African countries. About 50% of the world's population lives in malaria endemic countries [1].

About 1 million deaths occur each year due to malaria and most of them are young children. In malaria endemic countries it may cause 10% of all deaths in children. In Pakistan about 1.6 million cases of malaria occurs per year including 300 000 confirmed cases in public health-sector [2].

The goals of antimalarial treatment in *P. vivax* are to treat the malaria and to prevent the relapse of malaria. This cannot be achieved by a single drug so a combination of antimalarial is required to achieve the goal [3]. In chloroquine (CQ) sensitive regions, the WHO recommends 3 days of CQ therapy or an artemisinin combination therapy plus 2 weeks of PQ (provided the person is not G6PD deficient) [4].

Primaquine induces dose-dependent acute hemolytic anaemia in individuals with G6PD deficiency, a genetically X-linked disorder [5]. This condition is widely prevalent affecting over 400 million people globally, with a prevalence of 3–35% in tropical areas [6]. Global prevalence of G6PD deficiency is 4.5% and 1.8% in Pakistan [7].

Often the facility of G6PD testing is not available at points of care after a diagnosis of *P. vivax* malaria. As a result, Primaquine is usually given without prior G6PD testing, thus exposing vulnerable patients to the risk for hemolytic anemia. On the other hand, where it is not

administered exposes patients to the risk for repeated relapses of *P. vivax* malaria, with consequent morbidity and transmission. For the elimination of *P. vivax* liver-stage infections (radical cure), Primaquine is given with dose, 0.25mg base/kg/ body weight daily (3.5 mg/kg total dose) for 14 days, in addition to the antimalarial medicine that cures the blood-stage infection [8]. In a significant proportion of G6PD-deficient patients, however, the 14-day regimen of Primaquine induces dose-dependent, potentially severe hemolysis [9]. Instead of daily Primaquine they should receive once weekly Primaquine 0.75 mg/kg for 8 weeks [10].

The rationale of this study was to assess whether once a week for total eight week PQ regimen is effective at radical cure without the associated risk of hemolysis in G6PD deficient children. This regimen may be appropriate for poor countries where G6PD testing is unavailable. The objective of this study was to assess the safety of weekly Primaquine given to G6PD deficient persons, for the radical treatment of *Plasmodium vivax* Malaria.

2. METHODOLOGY

Cross sectional study were conducted at Pediatrics Out Door Department, Liaquat University of Medical and Health Sciences Hyderabad from January 2018 to August 2019. Approval was taken from Ethical Review Committee of LUMHS University Jamshoro. Total sample of 40 patients were include this study. With inclusion criteria 4 to 12 years old were living less than 100 kilometers away from the Civil Hospital Hyderabad due to the concern of delayed management in case of hemolysis. All the respondents were registered. Informed consent was taken from the parents.

Malarial parasite diagnose by doing MP test (Thick and thin slide method). Children with fever are classified as clinical suspected malaria according to integrated management of childhood and neonatal illness (IMNCI) protocols are also checked for malarial parasite. Blood is taken with finger prick on a glass slide and is checked for the presence or absence of malarial parasites. Continuously who are positive for *Plasmodium vivax* parasite, 2 ml blood was taken for G6PD level. Blood was also checked for Complete blood count, Reticulocyte count, Liver function test and creatinine. 5 ml blood was taken in anticoagulant bottle for liver function test, creatinine, complete blood count, and reticulocyte count tests. Haemoglobin < 7 g/dL, reticulocyte count > 4, SGPT > 80, G6PD Level < 60% of normal and creatinine > 1.2 was considered significant

Quantitative testing of G6PD activity was checked at LUMHS Research Laboratory by using spectrophotometric assay and functioning cold chain was maintained. If G6PD result showed decreased level (<60% of normal), they were enrolled for study.

After enrolment they were treated with Artemether and Lumefantrine for 3 days while Primaquine, 0.75 mg base/kg body weights was given once a week for 8 weeks. Patients were followed at OPD initially on 3rd day of therapy then every week for 8 weeks. Patients were informed about the risk for acute hemolytic anaemia when taking Primaquine. They were instructed to monitor the colour of their urine and to stop taking Primaquine if their urine becomes dark and give oral hydration and shift the patient to hospital, where patient will be admitted. Patients were taken from the area that is within 100 kilometers area near to Civil Hospital Hyderabad as they can reach the hospital within 1 hour in case of hemolysis at home. Clinical

assessment will be done and blood will be sent to laboratory to check for haemoglobin or hematocrit, serum creatinine or urea (blood urea nitrogen). Blood will then be transfused if Haemoglobin is < 7 g/dL or Haemoglobin is between 7 to 9 g/dL with concurrent hemolysis. If the Haemoglobin is > 9 or 7–9 g/dL and no evidence of concurrent hemolysis, then careful fluid management with monitoring of urine colour will be done. At weekly checkup we checked for fever and jaundice and asked for vomiting, abdominal pain, dizziness, breathlessness and color of urine. Compliance of Primaquine was also ensured. In case of Hemolysis, Primaquine was stopped and Hb level checked and managed accordingly.

The data was analyzed by using SPSS version 22.0. Categorical variables like: gender, complications, and outcome were analyzed by applying Chi Sq. Test and numerical values were measured in mean and frequency.

3. RESULTS

In this study total 40 children having G6PD deficiency were treated with Artemether and Lumefantrine while radical therapy was done with weekly dose of Primaquine for 8 weeks. 22(55%) children were between 9-12 years of age (as shown in Table 1). *Plasmodium vivax* was negative on 3rd day of therapy, it was also negative on 8 week of therapy (as shown in Table 2). Response to therapy and any adverse effect was monitored. There was no hemolysis during the first week and 8 weeks after therapy. Most common side effect was abdominal pain 4 (10%), other side effects are mentioned in Table 3. Mean hemoglobin was 11.8, while the rest of biochemical factors like SGPT, Creatinine and reticulocyte were normal (as mentioned in Table 4).

Table 1. Demographic parameters of study participants (n=40)

Characteristic	Number (n)	Percentage (%)
Age (Years)		
5-8	18	45
9-12	22	55
Weight (Kg)		
15-25	19	47.5
26-40	18	45
>40	3	7.5

Table 2. Mean value of hemoglobin concentration and vivax status in G6PD deficient children after giving primaquine (n=40)

Day of Primaquine	Mean Hb (mg/dl)	<i>Plasmodium vivax</i> +ve
0	11.6	Yes
1	11.6	Yes
3	11.4	No
7	11.1	No
8 weeks	12.0	No

+ve = Positive, Hb = Hemoglobin (mg/dl), P-value = 0.0001

Table 3. Number of subject adversely affected after Primaquine therapy (n=40)

Symptoms	Day 0 n (%)	Day 1 n (%)	Day 3 n (%)	Day 7 n (%)	Week 8 n (%)
Pallor	1 (2.5)	1 (2.5)	1 (2.5)	1 (2.5)	0 (0)
Bleeding	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Edema	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Jaundice	0 (0)	3 (7.5)	3 (7.5)	3 (7.5)	0 (0)
Respiratory Difficulty	0 (0)	1 (2.5)	0 (0)	0 (0)	0 (0)
Abdominal Pain	0 (0)	4 (10)	4 (10)	0 (0)	0 (0)
Hepatomegaly	1 (2.5%)	3 (7.5)	3 (7.5)	3 (7.5)	0 (0)
Splenomegaly	1 (2.5%)	3 (7.5)	3 (7.5)	3 (7.5)	0 (0)
Low Back Pain	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)

n = Number, % = Percentage, P-value = 0.0034

Table 4. Mean of hematological and biochemical parameters in G6PD deficient patients after taking primaquine (n=40)

Parameters	Day 0	Day 3	Day 7
Total Bilirubin (mg/dl)	1.2	1.3	1.5
ALT	35	35	40
AST	30	32	33
Hemoglobin (g/dl)	11.8	11.8	12.00
Hematocrit (%)	35.2	35.2	35.8
Reticulocyte (%)	1.8	1.8	2.1
Creatinine (mg/dl)	1.0	1.1	0.9

n = Number, P-value = 0.005

4. DISCUSSION

The malaria endemic countries where G6PD testing is not available have adopted Primaquine 0.75mg/kg/week (total 8 weeks) protocol for the radical cure of vivax malaria. As this regimen is useful for G6PDd A- variant (less severe form) so there is need of improvement in radical cure policies and practice. In this study 100% case of *Plasmodium vivax* were cured by giving Artemether and Lumefantrine combination therapy. On day three of treatment all cases had negative MP test. There was no hemolysis noted during eight weeks of treatment with Primaquine. Low dose primaquine prevented the relapse of vivax malaria without causing any major side effect. Minor side effects like abdominal and hepatomegaly was noted in few children. There was no significant change in hemoglobin concentration during the study period.

In a recent study PQ \leq 2.5 mg base/kg was associated with 25% chance of recurrence at 4–6 months, compared with 6.7% chance when $>$ 2.5 mg/kg to $<$ 5.0 mg/kg) was given. $>$ 5.0 mg/kg were associated with a recurrence rate of 0% at 1 month [11]. In other two studies there was high effectiveness of $>$ 5mg/kg PQ compared with a control arm [12,13].

There are different policies for the Radical cure of vivax with Primaquine across the Asian region. Some check G6PD level before Primaquine administration, although in practice mostly this is not applicable. In a study from Cambodia Primaquine was given (0.75 mg/kg PQ dose weekly for 8 weeks) to G6PD deficiency persons. There was no significant hemolysis noted during the total eight weeks of treatment [14]. Although G6PD testing is advised by various countries but it is not available most of the time, that's why physician's decision play an important role in decision making [15].

The use of Primaquine in private sector is conflicting, in some countries there is very little use while in others it is the main source of antimalarial treatment [16]. In an Indian study the relapse rate was 0% with the 14-day regimen, 26.7% with the 5-day, and 11.7% when no Primaquine treatment was given [17,18]. According to Chu et al Primaquine 0.5mg/kg (low dose) for 14 days was well tolerated as compared to 1mg/kg (high dose) for 14 days causing hemolysis [19].

According to a study done at Northern Pakistan the prevalence of G6PD deficiency is 2-8% and G6PD Med is the most frequent variant [20]. Primaquine causes serious hemolysis in persons having G6PD Med deficiency [21]. In Pakistan Primaquine administration needs mandatory G6PD testing although this test is not widely available. A study from the southern Pakistan showed that out of 200 participants 6 were G6PD deficient [22]. Recurrent jaundice due to hemolytic anemia can occur in G6PD deficient people. In a local study G6PD deficiency was detected in 1.8% people, 1.07% in Kashmiris, 1.47% in Punjabis, 2.77% in Sindhis, and 3.17% in Pathans. About 5.7% persons had the history of recurrent jaundice [23]. A 14-day course of Primaquine (PQ) can cause severe hemolysis in G6PD deficient persons and the testing for G6PD is seldom available enforcing the need of safe dose of PQ without G6PD testing.

A study from Pakistan concluded that the 8-week PQ course is more effective in preventing relapse and widespread use of this regimen could make an important contribution [24]. A 5-day course of PQ for vivax malaria is used commonly in South Asia to reduce the risk of hemolysis [25]. Studies from Pakistan and India showed that the 5-day PQ regimen is ineffective in reducing relapses [26,27]. The 14-day course of Primaquine is only recommended where the G6PD status of the individual is known [28]. That's why Primaquine is less frequently used as G6PD testing is not available in developing countries. This regimen was ineffective in Pakistan and India, because of frequent relapse rates [29]. As compared to 15 days course of Primaquine, 8 weeks course was found more effective in a local study [30].

In Asian countries there are different policies for Primaquine use. In some countries G6PD testing is required before the administration of Primaquine. Pakistan follows this protocol but mostly G6PD testing is not available. In Myanmar this facility is lacking in villages and they are using weekly Primaquine for total 8 weeks. This regimen is also followed in Iran.

In Pakistan there is reluctance and low prescription for Primaquine by Pediatricians and physicians due to their concern of hemolysis in G6PD deficiency children. A local study from Karachi showed that Primaquine was prescribed as prophylaxis to prevent relapse in only 6.2% of cases infected with *P. vivax* [31].

5. CONCLUSION

Primaquine 0.75mg/kg/week for total eight weeks is highly effective for the radical cure of *Plasmodium vivax* in G6PD deficient children. There is no recurrence of *Plasmodium vivax* after 8 weeks of therapy. We found this regimen safe as there was no hemolysis demonstrated in children. This regimen can be safely used in children who has *Plasmodium vivax* and whose G6PD level cannot be determined.

DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

CONSENT

Informed consent was taken from the parents.

ETHICAL APPROVAL

Ethical Approval was taken from Ethical Review Committee of LUMHS University Jamshoro, Pakistan.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- World Health Organization. WHO Fact Sheet on Malaria. Fact Sheet No 94. World Health Organization; 2007. Available:<http://www.who.int/mediacentre/factsheets/fs094/en/print.html>. Accessed: June 29, 2007
- Pakistan demographic and health survey; 2013. Available:http://www.nips.org.pk/abstract_files/PDHS%20Final%20Report%20as%20of%20Jan%2022-2014.pdf
- World Health Organization. Guidelines for the Treatment of Malaria. 3rd edition. Geneva, Switzerland: WHO; 2015.
- World Health Organization. Guidelines for the Treatment of Malaria. 2nd edition. Geneva, Switzerland: WHO; 2010.
- Deepika Fernando. Chaturaka Rodrigo. Senaka Rajapakse. Primaquine in vivax malaria: an update and review on management issues. *Malar J*. 2011;10:351.
- Sócrates Herrera Valencia. Iván Darío Ocampo. María Isabel Arce-Plata. Glucose-6-phosphate dehydrogenase deficiency prevalence and genetic variants in malaria endemic areas of Colombia. *Malar J*. 2016;15:291.
- Bushra Moiz. A review of G6PD deficiency in Pakistani perspective. *JPMA*. 2013;501-3
- Primaquine: report from CDC expert meeting on malaria chemoprophylaxis I. Hill DR, Baird JK, Parise ME, Lewis LS, Ryan ET, Magill AJ *Am J Trop Med Hyg*. 2006;75(3):402-15.
- Olalekan A. Uthman, Patricia M. Graves, Rachel Saunders. Safety of primaquine given to people with G6PD deficiency: systematic review of prospective studies. *Malar J*. 2017;16:346.
- World Health Organization. Point-of-care G6PD testing to support safe use of primaquine for the treatment of vivax malaria Geneva, Switzerland: World Health Organization; 2015.
- Primaquine radical cure of *Plasmodium vivax*: a critical review of the literature. John GK, Douglas NM, von Seidlein L, Nosten F, Baird JK, White NJ, Price RN *Malar J*. 2012;11:280.
- Resistance to therapies for infection by *Plasmodium vivax*. Baird JK *Clin Microbiol Rev*. 2009; 22(3):508-34.
- Primaquine for preventing relapses in people with *Plasmodium vivax* malaria. Galappaththy GN, Omari AA, Tharyan P *Cochrane Database Syst Rev*. 2007;(1):CD004389.
- Kheng S, Muth S, Taylor WR, Tops N, Kosal K, Sothea K, et al. Tolerability and safety of weekly primaquine against relapse of *Plasmodium vivax* in Cambodians with glucose-6-phosphate dehydrogenase deficiency. *BMC Med*. 2015;13:203.
- PATH. A guide to fluorescent spot testing for G6PD deficiency; 2014. Available:<http://sites.path.org/dx/files/2012/04/FST-Guidebook>.

16. Phanalasy S, Vongviengxay S. The Malaria Testing and Treatment Landscape in the Southern Lao People's Democratic Republic (PDR). In: ACT watch Group, editor, Washington DC. Proceedings of the 65th Annual American Society of Tropical Medicine and Hygiene Meeting. 2016 Nov 13–17; Atlanta; 2016.
17. Gogtay NJ¹, Desai S, Kamtekar KD, Kadam VS, Dalvi SS, Kshirsagar NA. Efficacies of 5- and 14-day primaquine regimens in the prevention of relapses in *Plasmodium vivax* infections. *Ann Trop Med Parasitol*. 1999;93(8):809-12.
18. Avalos S, Mejia RE, Banegas E, Salinas C, Gutierrez L, Fajardo M, et al. G6PD deficiency, primaquine treatment, and risk of haemolysis in malaria-infected patients. *Malar J*. 2018; 17: 415. Published online 2018;8. DOI: 10.1186/s12936-018-2564-2.
19. Chu CS, Bancone G, Moore KA, Win HH, Thitipanawan N, Po C, Chowwiwat N et al. Haemolysis in G6PD Heterozygous Females Treated with Primaquine for *Plasmodium vivax* Malaria: A Nested Cohort in a Trial of Radical Curative Regimens. *PLoS Med*. 2017;14(2):e1002224.
20. Moiz B, Nasir A, Moatter T, Naqvi ZA, Khurshid M. Molecular characterization of glucose-6-phosphate dehydrogenase deficiency in Pakistani population. *Int J Lab Hematol*. 2011;33:570–8.
21. Ashley EA, Recht J, White NJ. Primaquine: the risks and the benefits. *Malar J*. 2014;13:418.
22. Moiz B, Arshad HM, Raheem A, Hayat H, Ghanchi NK, Beg MA. Frequency of G6PD Mediterranean in individuals with and without malaria in Southern Pakistan. *Malaria Journal*. 2017;16(1):1-6.
23. Ali N, Anwar M, Ayyub M, Bhatti FA, Nadeem M, Nadeem A. Deficiency in some ethnic groups of Pakistan. *JCPSP*. 2005;15(3):137-41.
24. Leslie T, Mayan I, Mohammed N, Erasmus P, Kolaczinski J, Whitty CJ, Rowland M. A randomised trial of an eight-week, once weekly primaquine regimen to prevent relapse of *Plasmodium vivax* in Northwest Frontier Province, Pakistan. *PLoS One*. 2008;3(8):e2861.
25. WHO. Practical Chemotherapy of Malaria. Geneva: World Health Organisation. Technical Report Series, no.805; 1990.
26. Rowland M, Durrani N. Randomised controlled trial of 5 and 14 day primaquine therapy against relapses of malaria in an Afghan refugee settlement in Pakistan. *Tran R Soc Trop Med Hyg*.1999;93:642–643.
27. Gogtay NJ, Desai S, Kamtekar KD, Kadam VS, Dalvi SS, et al. Efficacies of 5 and 14 day primaquine regimens in the prevention of relapses in *Plasmodium vivax* infections. *Ann Trop Med Parasitol*. 1999;93:809–812.
28. WHO. Guidelines for the treatment of malaria. World Health Organisation; 2006.
29. Rowland M, Durrani N. Randomised controlled trial of 5 and 14 day primaquine therapy against relapses of malaria in an Afghan refugee settlement in Pakistan. *Tran R Soc Trop Med Hyg*. 1999;93:642–643.
30. Leslie T, Mayan I, Mohammed N, Erasmus P, Kolaczinski J, Whitty CJ, Rowland M. A randomised trial of an eight-week, once weekly primaquine regimen to prevent relapse of *Plasmodium vivax* in Northwest Frontier Province, Pakistan. *PLoS One*. 2008;3(8):e2861.
31. Bouma MJ, Goris M, Akhtar T, Khan N, Khan N, et al. Prevalence and clinical presentation of glucose-6-phosphate dehydrogenase deficiency in Pakistani Pathan and refugee communities in Pakistan; implications for the use of primaquine in regional malaria control programs. *Tran R Soc Trop Med Hyg*. 1995;89:62–64.

© 2021 Shaikh et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
 The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/78636>