

## Cardiac Myosin Heavy Chains

**Muhanad S. Abdelwahab<sup>1\*</sup>, Mukhallad Am. Mohamemed<sup>2</sup>,  
Abdelwahab Hassan Mohamed<sup>3</sup>, Mansour Abdelgader Bellal<sup>4</sup>  
and Mazin S. Abdalla Mohamed<sup>1</sup>**

<sup>1</sup>Department of Physiology, Napata College, Khartoum, Sudan.

<sup>2</sup>Jordan University of Science and Technology, Irbid, Jordan.

<sup>3</sup>National Ribat University, Khartoum, Sudan.

<sup>4</sup>University of Science and Technology, Khartoum, Sudan.

### Authors' contributions

*This work was carried out in collaboration between all authors. All authors read and approved the final manuscript.*

### Article Information

DOI: 10.9734/CA/2019/46305

#### Editor(s):

(1) Dr. Gerez Fernandes Martins, Department of Cardiology, Cardiology Institute of Rio de Janeiro, Brazil.

#### Reviewers:

(1) Aşkın Ender Topal, Dicle University, Turkey.  
(2) Korumilli Ramesh Kumar, General Surgery, S. V. S. Medical college and Hospital, K. N. R. University of Health Sciences, India.

(3) Paulo Roberto Barbosa Evora, Ribeirao Preto School of Medicine, University of Sao Paulo, Brazil.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/46305>

**Received 03 November 2018**

**Accepted 18 January 2019**

**Published 12 February 2019**

**Review Article**

### ABSTRACT

Myosin is defined as a mechano-enzyme molecule which converts the chemical energy stored as adenosine triphosphate (ATP) into mechanical energy (muscle contraction). Moreover, the cardiac muscle has different types of myosin heavy chain when it separated with the one dimensional electrophoresis; in addition to their structural difference cardiac myosin isozymes have different contractile functions.

**Keywords:** *Myosin heavy chains; myosin isozymes; alpha myosin heavy chain; beta myosin heavy chain.*

### 1. INTRODUCTION

Cardiac myosin isozymes are the thick filament of the cardiac muscle proper contractile proteins,

and they are differ in their nature and contractile properties. If cardiac myosin heavy chains ratio changes towards Alpha or Beta myosin heavy chain, it will cause changes in the contractile

\*Corresponding author: E-mail: muhanadsaad08@gmail.com;

properties of the cardiac muscle proper. Since cardiac contractility is the main determinant of the heart effectiveness, it is important to focus on the nature, properties of the cardiac myosin heavy chains, and the factors affecting myosin heavy chain ratio.

## 2. CARDIAC MYOSIN ISOZYMES

Based on electrophoretic mobility, Hoh et al 1978 [1] discovered that there are five distinct components of cardiac myosin isozymes. Two of them present in the atrial muscle named A1 & A2 and the other three are in the ventricular muscle designated V1, V2 & V3. These ventricular myosin chains have been differentiated by their speed of migration and subsequently on their molecular weights into V1 which is the fastest one and the slower one was detected to be V3 myosin isozyme while V2 was intermediate between V1 & V3. The two atrial components A1 & A2 migrate faster than the fastest ventricular myosin isozyme V1 [1].

In comparison with mammals, ventricular myosin of the amphibian hearts has somewhat different electrophoretic properties. Depending on the species. There are only one or two myosin components in the ventricle. Ventricular isozymes of Urodelan amphibians display mobility similar to V2 or V1 of the rat ventricular isozymes and the ventricular myosin of the Anurans migrates faster than V1 [2].

Cardiac myosin molecule is a hexamer (Fig. 1) which is comprised of two heavy chains and two pairs of light chain [3]. The myosin isozymes differ in their heavy chains structure [4], while

their light chains not being different in molecular size or stoichiometry [1], thus the difference only in their heavy chains.

In contrast to skeletal muscle which expresses multiple myosin heavy chain genes, the heart expresses only two myosin heavy chain genes which produce the Alpha & Beta myosin heavy chains [5]. Therefore the structural component of the ventricular myosin isozymes according to their heavy chains is as the following: V1 isozyme is homodimer consisting of two alpha myosin heavy chains, V3 isozyme is also homodimer consisting of two Beta heavy chains while V2 isozyme is heterodimer formed by one alpha & one beta heavy chains [6] see Fig. 1.

Analysis of the Calcium – activated myosin ATPase activity revealed that A1, A2 & V1 isozymes have about the same activity while V3 has the lowest activity and V2 has intermediate activity [1]. The V1 which has a much higher Calcium ATPase activity resembles fast-twitch skeletal muscle myosin while V3 behaves like slow – twitch skeletal muscle myosin [7].

The ventricular myosin isoforms have different distribution according to the age & species. In all species during the fetal life the ventricular myosin isoforms is essentially V3 ( $\beta\beta$ ) while the V1 ( $\alpha\alpha$ ) appears around the time of birth [8]. There are species differences; in adult mice and rats V1 ( $\alpha\alpha$ ) myosin isozyme remain the predominant one but in rabbit and pigs return to V3 ( $\beta\beta$ ) myosin isozyme after three weeks of age see Fig. 2. Adult dog, beef & human ventricular myosins are also formed by V3 ( $\beta\beta$ ) isoform only [8].





Fig. 1. MHC structure [9]



**Fig. 2. Effect of age on the cardiac MHC in rat and rabbit [9]**

Expression of myosin isoforms (V1 & V3) depends on the location of cardiomyocytes in the heart. V1 myosin isoform is present predominantly in the ventricular papillary and epicardium muscle layers while the V3 myosin isoform is distributed predominantly in the ventricular endocardium muscle layer [10]. Also V1 myosin isoform is present predominantly in the ventricular conductive system which is reflected by the higher concentration of the alpha myosin heavy chain in the ventricular conductive system [11].

### 3. CARDIAC MYOSIN HEAVY CHAINS

There are two genes encoding the alpha and beta cardiac myosin heavy chains [5]. Interestingly the complete sequences of alpha & beta cardiac myosin heavy chain nucleotides and amino acids have been determined in 1989 [3]. The complete alpha myosin heavy chain cloned DNA is 5930 base pairs and encodes a protein of 1938 amino acid residues in length with predicted molecular mass of 223,511 Daltons. The complete beta myosin heavy chain cloned DNA is 5925 base pairs and encodes a protein of 1935 amino acid residues in length with predicted molecular mass of 223,172 Daltons. The alpha and beta myosin heavy

chain nucleotide sequences are 92% identical and the amino acid sequences of the alpha and beta myosin heavy chains are 93% identical [3].

The two myosin heavy chains are different in the concentration of two amino acids in their structure. Methionine is found in higher concentration in alpha myosin heavy chain while Arginine amino acid is found in higher concentration in beta myosin heavy chain [4].

Some functional difference has been found to be correlated with these structural differences. The Calcium – activated ATPase activity of the alpha myosin heavy chain is higher than that of the beta myosin heavy chain [1]. In guinea pig hearts, the beta myosin heavy chain is about five times more economical than the alpha myosin heavy chain [12]. Recently, Narolska et al. 2005 observed that human ventricular muscle consists of beta myosin isoform while the human atrial muscle consists of both alpha and beta myosin isoform [13]. Also they have observed that human ventricular muscle which is beta myosin heavy chain is fivefold more economical in force development and nine times slower than atrial muscle [13]. Functionally, it has been recently found that the activity of the two cardiac myosin heavy chains alpha and beta can be increased by the reversible lysine acetylation [14].

### 4. REDISTRIBUTION OF THE CARDIAC MYOSIN HEAVY CHAINS

There are many factors that could cause redistribution of the cardiac MHC isoforms (Table 1): These factors can be classified into physiological [8,14,15], environmental [16], metabolic [17], pathological factors [18,19,20], and the redistribution due to drugs administration. Some of these factors affecting redistribution of the cardiac MHC isoforms are associated with cardiac hypertrophy, while other factors are not [14,15].

The physiological factors affecting cardiac MHC isoforms redistribution are age and routine physical exercise. The effect of age on cardiac MHC isoforms redistribution is not associated with heart hypertrophy [8]. This redistribution depends on species; in mice and rats the amount of alpha MHC isoform increases with age, while the amount of beta MHC isoform decreases with age [8].

Similarly the distribution of MHC due to hypertrophy depends on type of exercise at least in rat. Hypertrophy after swimming exercise shows that MHC is shifted toward alpha while hypertrophy after running exercise is not associated with changes in the cardiac MHCs distribution [8,14, 15].

The effect of adaptation to heat stress has been demonstrated by Horowitz et al 1986 [16]; they found that there is an alteration in cardiac myosin isozyme distribution as an adaptation to chronic environmental heat stress in rats. Chronic environmental heat stress causes down regulation of V1 ( $\alpha\alpha$ ) myosin isozyme and up regulation of V3 ( $\beta\beta$ ) myosin isozyme [16]. The redistribution of the myosin heavy chains due to acclimatization to hot weather is associated with a decrease in thyroid hormone level [16].

The metabolic factors that can cause cardiac MHC redistribution is demonstrated by food restriction & Carbohydrate rich meal replacement. In food restricted rats there were downregulation of alpha MHC isoform and upregulation of beta MHC isoform but this shifting can be prevented by carbohydrate rich meal replacement [17]. The shifting in MHC isoforms is correlated with reduction in heart weight in food restricted rats [17].

There are multiple pathological factors associated with the redistribution of the myosin heavy chains of the heart. Rupp et al 1981 [15], showed in the case of cardiac hypertrophy due to renal hypertension (Goldblatt II) there was a shift on the ventricular myosin isozyme toward V3 ( $\beta\beta$ ) myosin isoform. This was reinforced in 1989 by the observation of Yazaki et al 1989 [18] where they demonstrated that the left ventricular hypertrophy was evident from 3 days after pressure loading and the isozymic transition of myosin heavy chain from the alpha MHC isoform to the beta MHC isoform was detected within 24 hours after aortic constriction. Thus pressure overload led to MHC shift toward beta MHC isoform.

In some cardiac pathological conditions like myocardial infarction there is change in the cardiac MHCs toward V3 ( $\beta\beta$ ) myosin isoform, but this change can be inhibited by administration of Losartan (an angiotensin II blocker) [19]. Similarly atrial fibrillation was accompanied by a significant shift from the alpha

MHC isoform to the beta MHC isoform in the atrial tissue [20]. Also the alpha MHC isoform in the heart decreases in diabetes mellitus to a significant level [21].

The correlation between sex hormone and the distribution of the cardiac MHCs is shown in experiments on the effect of gonadectomy & sex hormone replacement on the cardiac MHCs in the spontaneously hypertensive rats. Lengsfeld et al 1988 [22], showed that the male gonadectomy causes shift in the myosin isozyme pattern toward V3 ( $\beta\beta$ ) myosin isozyme while testosterone hormone replacement leads to myosin isozyme pattern in favor of V1 ( $\alpha\alpha$ ) myosin isoform. This observation was supported recently by Jazbutyte V et al. 2006 [23] who reported that; in spontaneously hypertensive female rat, the ovariectomy decreases the alpha myosin heavy chain in the heart and shift the MHC ratio toward beta MHC accumulation. While estrogen substitution in ovariectomized young spontaneously hypertensive rat results in an increase in alpha MHC isoform.

Administration of thyroid hormone makes the cardiac MHC to redistribute in favor of V1 ( $\alpha\alpha$ ) myosin isozyme which is associated with heart hypertrophy. While decreasing the thyroid hormone level in the blood has the reverse effect, it changes the cardiac MHC ratio toward the beta MHC isoform [24,25]. Similarly dexamethazone (a synthetic glucocorticoid drug) administration has the same effect of thyroid hormone on the cardiac MHCs. This drug induces cardiac hypertrophy and makes a change in the cardiac MHCs toward the fast alpha MHC isoform [26].

Rupp et al. 1991 [27] showed the effect of positive inotropic agents on myosin isozymes in cultured cardiac myocytes. Isoproterenol and Phenylephrine caused an increase in V1 native myosin isozyme. The isoproterenol made a shift in the cardiac MHC toward the alpha MHC [27].

The only drug that causes a shift of cardiac MHC toward beta MHC is Cocaine [28].

## 5. HEART HYPERTROPHY

The word hypertrophy derived from the Greek, hyper means above or more than normal, and Trophe means nutrition [29]. Hypertrophy is defined as the enlargement or overgrowth of an

**Table 1. Factors affecting ventricular MHC distribution**

<b>Factor affecting MHCs distribution</b>	<b>Shifting of MHC</b>	<b>references (Ref)</b>
Physiological factors	Shifting of MHCs	(Ref)
Increase age	In mice increase $\alpha$ & decrease $\beta$ MHCs	[8]
Hypertrophy after swimming exercise	Increase $\alpha$ MHC	[14,15]
Hypertrophy after running exercise	No change	[14,15]
Environmental factor	Shifting of MHCs	Ref
Adaptation to heat stress	Decrease $\alpha$ MHC & increase $\beta$ MHC	[16]
Metabolic factors	Shifting of MHCs	Ref
Food restriction	Decrease $\alpha$ & increase $\beta$	[17]
CHO rich meal	Increase $\alpha$ & decrease $\beta$	[17]
Pathological factors	Shifting of MHCs	Ref
Hypertrophy due to renal hypertension	Increase $\beta$ MHC	[16,18]
Myocardial infarction	Increase $\beta$ MHC	[19]
Atrial fibrillation	Increase $\beta$ MHC	[20]
Gonadectomy	Decrease $\alpha$ and increase $\beta$ MHCs	[22,23]
Hypophsectomy	Increase $\beta$ MHC	[32,24]
Change due to drug administration	Shifting of MHCs	Ref
Sex hormones	Increase $\alpha$ and decrease $\beta$ MHCs	[22,23]
Thyroid hormone	Increase $\alpha$ MHC	[32,24]
Anti thyroid drugs	Increase $\beta$ & decrease $\alpha$ MHCs	[32,24]
Dexamethasone	Increase $\alpha$ MHC	[26]
Positive inotropic drugs like isoproterenol, forskolin, phenylephrine	Increase $\alpha$ MHC	[27]
Cocaine	Increase $\beta$ MHC	[28]

organ or part due to an increase in size of its cells [29]. The heart hypertrophy has two phenotypes named as concentric hypertrophy and eccentric hypertrophy [29]. In both forms of hypertrophy the cardiac dry mass is increased [29].

Concentric hypertrophy is due to pressure overload in which there is an increased thickness of ventricular wall with little or no change in chamber volume, with new sarcomeres added in parallel to existing sarcomeres [29,30]. While the eccentric form of hypertrophy is due to volume overload in which there is an increased chamber volume with ventricular wall thickness increased in proportion to the chamber dimension [29,30]. The increased thickness of the ventricular wall is by adding new sarcomeres in series to existing sarcomeres [30].

## **6. FACTORS PROMOTING LEFT VENTRICULAR HYPERTROPHY**

It is now appreciated that left ventricular hypertrophy is mediated not only by the

mechanical stress of pressure overload, but also by various neurohormonal substances that independently exert trophic effects on myocytes and nonmyocytes in the heart [31]. The trophic factors that promote left ventricular hypertrophy include angiotensin II, aldosterone, noradrenalin, and insulin which directly promote myocyte hypertrophy and matrix deposition independent of their effects on systemic arterial pressure [31].

## **7. CONCLUSION**

The current review concludes that the ventricular myosin isoforms of the mammalian hearts are distinct, and they do possess different contractile properties. However, when it comes to the structural difference between the ventricular myosin isoforms it appears to be only in their heavy chains. Therefore the contractile property of the ventricle is determined by the dominant type of the myosin heavy chain contained within.

Alpha myosin heavy chain has a higher content of  $Ca^{++}$ - myosin activated ATPase activity when

compared to Beta myosin heavy chain. Moreover, Ca<sup>++</sup> myosin activated ATPase activity of the Alpha myosin heavy chain resembles that in the fast-twitch skeletal muscle, while the Ca<sup>++</sup> myosin activated ATPase activity of Beta myosin heavy chain resembles the slow-twitch skeletal muscle. Beta myosin heavy chain is more economical than Alpha myosin heavy chain regarding force development.

Factors which can affect Alpha to Beta myosin heavy chain ratio will produce changes in the velocity of contraction, and consequently the rate of ATP consumption.

### CONSENT

It is not applicable.

### ETHICAL APPROVAL

It is not applicable.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

1. Hoh JF, McGrath PA, Hale PT. Electrophoretic analysis of multiple forms of rat cardiac myosin: Effects of hypophysectomy and thyroxine replacement. *J Mol Cell Cardiol.* 1978;10:1053-76.
2. Karasiński J, Kilarski W. Myosin isoenzymes of amphibian hearts. *Comp Biochem Physiol B.* 1986;83:677-9.
3. McNally EM, Kraft R, Bravo-Zehnder M, Taylor DA, Leinwand LA. Full-length rat alpha and beta cardiac myosin heavy chain sequences. Comparisons suggest a molecular basis for functional differences. *J Mol Biol.* 1989;210:665-71.
4. Hoh JF, Yeoh GP, Thomas MA, Higginbottom L. Structural differences in the heavy chains of rat ventricular myosin isoenzymes. *FEBS Lett.* 1979;97:330-4.
5. Mahdavi V, Periasamy M, Nadal-Ginard B. Molecular characterization of two myosin heavy chain genes expressed in the adult heart. *Nature.* 1982;297:659-64.
6. Samant SA, Courson DS, Sundaresan NR, Pillai VB, Tan M, Zhao Y, Shroff SG, Rock RS, Gupta MP. HDAC3-dependent reversible lysine acetylation of cardiac myosin heavy chain isoforms modulates their enzymatic and motor activity. *J Biol Chem.* 2011;286:5567-77.
7. Pope B, Hoh JF, Weeds A. The ATPase activities of rat cardiac myosin isoenzymes. *FEBS Lett.* 1980;118:205-8.
8. Lompre AM, Mercadier JJ, Wisnewsky C, Bouveret P, Pantaloni C, D'Albis A, Schwartz K. Species- and age-dependent changes in the relative amounts of cardiac myosin isoenzymes in mammals. *Dev Biol.* 1981;84:286-90.
9. Gupta MP. Factors controlling cardiac myosin-isoform shift during hypertrophy and heart failure. *J Mol Cell Cardiol.* 2007; 43:388-403.
10. Litten RZ, Martin BJ, Buchthal RH, Nagai R, Low RB, Alpert NR. Heterogeneity of myosin isozyme content of rabbit heart. *Circ Res.* 1985;57:406-14.
11. Komuro I, Nomoto K, Sugiyama T, Kurabayashi M, Takaku F, Yazaki Y. Isolation and characterization of myosin heavy chain isoforms of the bovine conduction system. *Circ Res.* 1987;61:859-65.
12. Van der Velden J, Moorman AF, Stienen GJ. Age-dependent changes in myosin composition correlate with enhanced economy of contraction in guinea-pig hearts. *J Physiol.* 1998;507:497-510.
13. Narolska NA, van Loon RB, Boontje NM, Zaremba R, Penas SE, Russell J, Spiegelberg SR, Huybregts MA, Visser FC, de Jong JW, van der Velden J, Stienen GJ. Myocardial contraction is 5-fold more economical in ventricular than in atrial human tissue. *Cardiovasc Res.* 2005; 65:221-9.
14. Rupp H. Differential effect of physical exercise routines on ventricular myosin and peripheral catecholamine stores in normotensive and spontaneously hypertensive rats. *Circ Res.* 1989;65:370-7.
15. Rupp H. The adaptive changes in the isoenzyme pattern of myosin from hypertrophied rat myocardium as a result of pressure overload and physical training. *Basic Res Cardiol.* 1981;76:79-88.
16. Horowitz M, Peyser YM, Muhlrad A. Alterations in cardiac myosin isoenzymes distribution as an adaptation to chronic environmental heat stress in the rat. *J Mol Cell Cardiol.* 1986;18:511-5.
17. Morris GS, Herrick RE, Baldwin KM. Dietary carbohydrates modify cardiac myosin isoenzyme profiles of semistarved rats. *Am J Physiol.* 1989;256:R976-81.
18. Yazaki Y, Tsuchimochi H, Kurabayashi M, Komuro I. Molecular adaptation to

- pressure overload in human and rat hearts. *J Mol Cell Cardiol.* 1989;(Suppl 5):91-101.
19. Zhang ML, Elkassem S, Davidoff AW, Saito K, ter Keurs HE. Losartan inhibits myosin isoform shift after myocardial infarction in rats. *Mol Cell Biochem.* 2003; 251:111-7.
  20. Narolska NA, Eiras S, van Loon RB, Boontje NM, Zaremba R, Spiegelen Berg SR, Stooker W, Huybregts MA, Visser FC, van der Velden J, Stienen GJ. Myosin heavy chain composition and the economy of contraction in healthy and diseased human myocardium. *J Muscle Res Cell Motil.* 2005;26:39-48.
  21. Rundell VL, Geenen DL, Buttrick PM, de Tombe PP. Depressed cardiac tension cost in experimental diabetes is due to altered myosin heavy chain isoform expression. *Am J Physiol Heart Circ Physiol.* 2004;287:H408-13.
  22. Lengsfeld M, Morano I, Ganten U, Ganten D, Rüegg JC. Gonadectomy and hormonal replacement changes systolic blood pressure and ventricular myosin isoenzyme pattern of spontaneously hypertensive rats. *Circ Res.* 1988;63:1090-4.
  23. Jazbutyte V, Hu K, Kruchten P, Bey E, Maier SK, Fritzemeier KH, Prella K, Hegele-Hartung C, Hartmann RW, Neyses L, Ertl G, Pelzer T. Aging reduces the efficacy of estrogen substitution to attenuate cardiac hypertrophy in female spontaneously hypertensive rats. *Hypertension.* 2006;48:579-86.
  24. Carter WJ, Kelly WF, Faas FH, Lynch ME, Perry CA. Effect of graded doses of triiodothyronine on ventricular myosin ATPase activity and isomyosin profile in young and old rats. *Biochem J.* 1987;247: 329-34.
  25. Rundell VL, Manaves V, Martin AF, de Tombe PP. Impact of beta-myosin heavy chain isoform expression on cross-bridge cycling kinetics. *Am J Physiol Heart Circ Physiol.* 2005;28:H896-903.
  26. Muangmingsuk S, Ingram P, Gupta MP, Arcilla RA, Gupta M. Dexamethasone induced cardiac hypertrophy in newborn rats is accompanied by changes in myosin heavy chain phenotype and gene transcription. *Mol Cell Biochem.* 2000;209:165-73.
  27. Rupp H, Berger HJ, Pfeifer A, Werdan K. Effect of positive inotropic agents on myosin isozyme population and mechanical activity of cultured rat heart myocytes. *Circ Res.* 1991;68:1164-73.
  28. Henning RJ, Silva J, Reddy V, Kamat S, Morgan MB, Li YX, Chiou S. Cocaine increases beta-myosin heavy-chain protein expression in cardiac myocytes. *J Cardiovasc Pharmacol Ther.* 2000;5:313-22.
  29. Dorn GW 2<sup>nd</sup>, Robbins J, Sugden PH. Phenotyping hypertrophy: Eschew obfuscation. *Circ Res.* 2003;92(11):1171-5.
  30. Muhl C, Dassen WRM, Kuipers H. Cardiac remodelling: Concentric versus eccentric hypertrophy in strength and endurance athletes. *Netherlands Heart Journal.* 2008; 16(4):129-133.
  31. Richard E, Katholi, Daniel M. Couri. Left ventricular hypertrophy: Major risk factor in patients with hypertension: Update and practical clinical applications. *International Journal of Hypertension.* 2011;10. [Article ID 495349, 2011] DOI: 10.4061/2011/495349
  32. Hoh JF, Egerton LJ. Action of triiodothyronine on the synthesis of rat ventricular myosin isoenzymes. *FEBS Lett.* 1979;101:143-8.

© 2019 Abdelwahab et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<http://www.sdiarticle3.com/review-history/46305>