



Forage Potential of Alfalfa with Oats and Barley in Intercropping System

Poloko E. Mosebi^{1*}, Puleng A. Matebesi-Ranthimo², Moeketsi P. Ntakatsane²
and Ratsele Ratsele²

¹Department of Animal Science, Faculty of Agriculture, National University of Lesotho, Maseru, Lesotho.

²Department of Range and Resource Management, Ministry of Forestry, Range and Soil Conservation, Maseru, Lesotho.

Authors' contributions

This work was carried out in collaboration between all authors. Author PEM designed the study, laid and monitored the field trials, collected data, performed the statistical analysis and wrote the first draft of the manuscript. Authors PAM, MPN and RR did proof reading, restructured and approved the manuscript. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJRAF/2018/42731

Editor(s):

(1) Dr. Cengiz Yucedag, Professor, Department of Landscape Architecture, Mehmet Akif Ersoy University, Turkey.

Reviewers:

(1) Branko Cupina, University of Novi Sad, Serbia.

(2) Estella Rosseto Januszkiewicz, State University of Mato Grosso do Sul, Brazil.
Complete Peer review History: <http://www.sciencedomain.org/review-history/26278>

Original Research Article

Received 05 May 2018
Accepted 24 July 2018
Published 19 September 2018

ABSTRACT

Production of cultivated pastures with high energy and protein levels remains a challenge during winter months for livestock. Field trials were conducted to investigate the effects of cropping systems using intercropping (alfalfa + oat and alfalfa + barley) and monocropping (alfalfa, oat and barley) on plant height, yield and quality characteristics. Furthermore, to study the competition experienced by legume and cereals when planted in a mixture. The field trial was carried out in a randomised complete block design with five treatments including sole oat, sole barley, sole alfalfa, alfalfa + oat intercropping and alfalfa + barley intercropping and three replicates. The study was conducted at the Experimental Farm of the National University of Lesotho for two growing seasons (2015 and 2016). The results revealed that intercropping increased yields of forage crops in comparison to monocrops. Intercropping significantly increased protein levels and reduced fibre concentrations. In intercropping competition indices, land equivalent ratios indicated yield

*Corresponding author: E-mail: pe.mosebi@nul.ls, mosebipe@gmail.com;

advantages for intercropping. Relative crowding coefficient and aggressivity values showed significant advantages of cereal over legume. So, the intercropping system was more productive than respective monocrops.

Keywords: Alfalfa forage; yield; quality; competition indices.

ABBREVIATIONS

<i>A</i>	: <i>Aggressivity</i>
<i>ADF</i>	: <i>Acid detergent fibre</i>
<i>ANOVA</i>	: <i>Analysis of variance</i>
<i>CP</i>	: <i>Crude protein</i>
<i>D</i>	: <i>Days after emergence</i>
<i>DM</i>	: <i>Dry matter</i>
<i>K</i>	: <i>Relative crowding coefficient</i>
<i>LER</i>	: <i>Land equivalent ratio</i>
<i>LSD</i>	: <i>Least Significant Difference</i>
<i>N</i>	: <i>Nitrogen</i>
<i>NDF</i>	: <i>Neutral detergent fibre</i>

1. INTRODUCTION

Intercropping is the system including two or more species sown at the same time and growing together on the same field [1]. Intercropping of legumes together with cereals for food or forage is used in many parts of the world and has shown great potential in many ways. Intercrops have potential on crop productivity by reducing weed pressure, sustaining plant health and improving dry matter yields [2]. Besides, it sustains fertility status of soil and the possibility of nitrogen accumulation from the leguminous plants to other plants in the soil. Biological fixation of nitrogen is completed by two nitrogen-fixing bacteria namely free-living bacteria in the soil and symbiotic relationship bacteria which live in the leguminous plant roots enriching the surrounding soil with usable nitrogen for cereal plants [3].

Cereal-legume combinations are often used for forage and as cover crops. On animal husbandry, legume-cereal mixtures may be interesting forage harvested for green forage and maintenance of continuity of feed supply [4]. In other studies, winter grains like rye and wheat are combined with a leguminous plant as a cover crop. Intercrops protect soil from erosion, help to improve soil tilth, and left as dead cover at the soil surface. Very often the legume provides nitrogen, while the cereal produces organic matter [5]. Some research indicates that legume plants like clover fix more nitrogen into plant – usable form when grown with cereals such as grasses than when planted in a pure stand [6].

Intercrops including legumes are known to use natural resources well and enhance forage than cereal sole cropping [7].

However, in Southern African countries, in particular, Lesotho, intercropping of legume and cereals mixture for production of forage is not a common cropping system in the winter season. Monocultures of leguminous plants or cereal crops are rarely used for forage and do not provide satisfactory results for forage production [8]. In particular, production of forage from cereals is usually lower than that required to meet satisfactory nutritional requirements for ruminant animals. The causes of declining livestock productivity in Lesotho are the deteriorating communal rangelands, low use of planted forages and low quality feeds in winter and spring season. Ruminant animals often face green forage scarcity during winter and spring months and thus have to survive on cereal residues of previous plants including maize straw and sorghum stalk which are not a rich source of digestible nutrients [8].

Intercropping systems may contribute significantly to the winter and spring season diet of animals [9]. Other studies reported the benefits of growing leguminous plants together with cereal crops in winter months. Intercrops including alfalfa legume and cereals like oat and barley are the important winter forage for the sustenance of livestock [10,11]. Studies have shown that nutritive value and yield of forage is high when produced in cereal-legume mixtures [10,11]. Cereal crops provide sufficient amount of carbohydrate, while legumes are efficient in increasing protein and mineral content of forage which is necessary for livestock health and productivity.

Alfalfa (*Medicago sativa* L.) is a herbaceous legume that has great growth productivity and good forage recovery after cutting [12]. Alfalfa has the ability to add nitrogen to the soil and store energy in the root crown that helps the buds in a quick re-growth which results in high yield [13]. It is the most important and profitable legume used for the production of forage mainly in intercrop combinations. Mixtures of alfalfa legumes with cereals forage (rye) greatly

improve minerals and decrease the prevalence of pasture toxicity [14]. Alfalfa has the ability to adapt to various environmental conditions and tolerates low temperatures. It was stated that alfalfa and wheat in the intercropping system in winter months increased yield, improved growth, reduced weeds, made a better soil coverage and keep it from erosion [15].

Oat plant (*Avena sativa* L.) is a forage crop planted primarily for grain and forage and often grown with a leguminous plant [16]. Oats form an excellent combination and produce high yields, forage quality and minerals when planted along with other winter season legume crops including vetch (*Vicia sativa*) and senji (*Indian clover*) [17]. Studies have shown that some intercropping systems such as vetch (*Vicia sativa* L.) – oats intercropping have a great potential for improving nutritive value of forage compared to sole cropping [18], whereas other intercropping systems such as berseem (*Trifolium alexandrinum*) – oats intercropping protect soil from erosion, limit weed population and enhance forage productivity [19; 20]. Oats – shaftal (*Persian clover*) intercropping has been shown to reduce diseases, suppress weeds, and improve the nutritive (protein) value of crop compared to oats alone [21].

Barley plant (*Hordeum vulgare* L.) is a leafy forage species and produce valuable forage for livestock. Barley can be conserved in the form of hay or silage and used later when pastures become unproductive [22]. In forage cropping system, barley forage has been planted in mixture with a legume such as berseem (*Trifolium alexandrinum*) and Persian clover (*Trifolium resupinatum*) and improved forage production [23]. Studies have shown that berseem clover and barley mixture produced higher forage productivity and nutritive value than sole barley [24]. Intercropping barley with vetch produced greater yields and nutritive value higher than either cereal or legume crop alone [25].

In this study, the hypothesis was that cropping systems impacted forage production mainly through influencing nutritive value composition

and plant growth attributes. This study is, therefore, conducted to determine forage yield potential and nutritive value of alfalfa with oat and barley at four growth stages in intercropping and sole cropping systems, and to study the influence of intercropping system on the growth rate of cereal-legume species planted in the mixtures.

2. MATERIALS AND METHODS

2.1 Site Description

The study was conducted at the Experimental Farm of the National University of Lesotho (29°45'S; 27°72'E) for two growing seasons during the winter seasons of 2015 and 2016. The field trial was established on a well- drained sandy clay loam soil with medium to coarse texture whose characteristics are presented in Table 1. Weather data during the experimental period for two growing seasons are given in Table 2.

2.2 Forage Management and Experimental Design

The field trial comprised of alfalfa, barley and oats planted in pure stands and intercropping alfalfa with barley and oats. Seedbed was prepared with tractor disc harrow. Animal manure was applied at the rate of 70 kg ha⁻¹ in accordance with soil properties, in particular, N content and incorporated into the soil before plantation to reflect the common practice. Oat, barley and alfalfa crop were sown at the rate of 80, 100 and 25 kg ha⁻¹ respectively. In mixture combinations, the seed of component forage crops was homogenised at 50:50 ratios before planting. The sole crop and intercrop treatments were established in rows spaced 20 cm apart within 10m × 20m plots; the intercrops of alfalfa with oats and barley were planted in alternate rows with the same row spacing done with a single row drill. Alfalfa seeds were inoculated with rhizobia to stimulate biological nitrogen fixation. The field trial was carried out in a randomised complete block design (RCBD) with five treatments including sole oat, sole barley and sole alfalfa, alfalfa + oat intercropping and

Table 1. Pre-plant soil nutrient analysis (0-40 cm) at the experimental area

Year	N	P	K	Ca	Mg	Fe	Mn	pH	Organic matter
									(gkg ⁻¹)
									(mgkg ⁻¹)
2015	82	21	101	1.35	0.32	7.2	3.8	6.8	8.9
2016	84	19	104	1.32	0.43	8.1	3.2	6.5	9.9

Table 2. Total rainfall and average temperature per month during experimental period

Month	Rainfall (mm)		Temperature (°C)		Total rainfall (mm)	Average temperature (°C)
	2015	2016	2015	2016		
January	10.5	19	32.2	44.4	29.5	38.3
February	14.6	9.7	30.5	28.6	24.3	29.6
March	20.9	104	27.1	24.2	125.9	25.7
April	11.3	3.3	20.2	23.1	14.6	21.7
May	8.7	5.8	19.5	22.8	14.5	21.2
June	2.1	7.3	14.6	15	9.4	14.8
July	3.2	0	15.2	17.2	3.2	16.2
August	6.1	13.4	17.6	22.2	19.5	19.9
September	4.2	8.5	21.8	25.4	12.7	23.6
October	11.5	5.33	26.7	27.4	16.8	27.1
November	7.3	20.4	29.4	26.5	27.7	30
December	18.5	3.2	30.2	31.4	21.7	26.6
Total rainfall and average temperature	118.9	199.9	23.8	25.7	319.8	24.6

alfalfa + barley intercropping and three replicates and the whole experiment was conducted in rainfed conditions. Rainfall and temperature recorded are presented in Table 2 during the forage crop growth period.

2.3 Forage Measurements

2.3.1 Plant height

The plant height was measured at the major growth stages (tillering, jointing, flowering and ripening) using disc pasture meter by averaging nine readings recorded inside a 2m x 2m square quadrat.

2.3.2 Dry matter yield

Green forage yield was measured at the major growth stages by harvesting above ground biomass within the 4m² quadrat randomly sampled in each plot using manual shears. Forage samples from each plot were oven dried at 80°C for at least 48 h to measure the dry matter yield.

2.3.3 Intercropping competition indices

Land equivalent ratio (LER), the relative crowding coefficient (K) and the aggressivity (A) were calculated to determine the impact of competition between the legume and cereal in a mixture stands. LER shows the efficiency of intercropping compared to sole cropping for the use of environmental resources. When LER is greater than one, the intercropping improved the productivity of the intercropped species [26]. The

LER was calculated according to the following equation:

$$LER = \left(\frac{Y_{Li}}{Y_L} \right) + \left(\frac{Y_{Ci}}{Y_C} \right)$$

Where, Y is the yield per unit area, Y_L and Y_C are the yields of legume (alfalfa) and cereal (oat or barley), respectively, as monocrops and Y_{Li} and Y_{Ci} are the yields of legume (alfalfa) and cereal (oat or barley), respectively, as intercrops.

The relative crowding coefficient (K) measures the relative dominance of one species in a mixture stands [27]. There is no competition when K is equal to one, the species is less competitive when K is lower than one and the species is more competitive in resource use when K is greater than one [27]. The K was calculated with the following equation:

$$K_L = \frac{Y_{Li}Z_{Ci}}{(Y_L - Y_{Li})Z_{Li}}$$

$$K_C = \frac{Y_{Ci}Z_{Li}}{(Y_C - Y_{Ci})Z_{Ci}}$$

Where, Z_{Li} is the sown proportion of legume (alfalfa) in mixture stands and Z_{Ci} the sown proportion of cereal (oat or barley) in mixture stands.

Aggressivity measures the competition between two different plant species [27]. Both plants are equally competitive if A_c is equal to zero. The cereal is the dominated species if A_c is negative. The cereal species is dominant if A_c is positive

[27]. The aggressivity is derived from the following equation:

$$\text{Aggressivity of Legume (alfalfa) } A_L = \left(\frac{Y_{LI}}{Y_L Z_{LI}} \right) - \left(\frac{Y_{CI}}{Y_C Z_{CI}} \right)$$

$$\text{Aggressivity of Cereal (oat or barley) } A_C = \left(\frac{Y_{CI}}{Y_C Z_{CI}} \right) - \left(\frac{Y_{LI}}{Y_L Z_{LI}} \right)$$

2.3.4 Forage quality

The second set of random samples of biomass for forage species of each plot was taken at each major growth stages to measure the forage quality; crude protein (CP), acid detergent fibre (ADF) and neutral detergent fibre (NDF) in the Department of Animal Science Laboratory. CP based on DM was calculated by multiplying the Nitrogen (N) content by 6.25 using the Kjeldahl method [28]. NDF based on DM was measured by boiling a forage sample using neutral detergent under neutral pH conditions. ADF was measured using acid detergent under low pH condition [29].

2.4 Statistical Methods

Statistical Analysis System proc mixed procedure was used for data analysis [30]. Analysis of variance (ANOVA) was performed to measure the effects of intercropping and sole cropping treatments on the productivity of forage. Treatment means were separated using Fisher's Protected Least Significant Difference (LSD) test and differences were considered significant at $p \leq 0.05$. The mean comparison was conducted using Duncan multiple range test.

3. RESULTS AND DISCUSSION

3.1 Forage Growth and Yield Attributes

3.1.1 Plant height

Plant height recorded at different forage growth stages showed significant differences among cropping system treatments in Table 3. At tillering, maximum plant height was obtained by intercropping treatments alfalfa + oat and alfalfa + barley and monoculture treatments of oat alone, barley alone and the lowest by alfalfa alone during two growing seasons 2015 and 2016. Cropping system effect on plant height started to appear at jointing to ripening stages

where maximum plant height was obtained by intercropping treatments alfalfa + oat and alfalfa + barley compared to monoculture treatments oats alone barley alone and the lowest by alfalfa alone during two growing seasons 2015 and 2016. Plant height is a major parameter of crop productivity that maximises the use of the climatic and surface environmental resources available, especially light, carbon dioxide, water and nutrients, allowing the production of maximum crop yield [31]. From the results, it appeared that the height character in intercropping is mainly greater than monoculture which could be the result of efficient utilisation of weather and environmental resources and minimum competition among the plants of different species; legume and cereal. There is a possibility that the alfalfa legume may have produced and shared biologically fixed nitrogen with its oat and barley cereals. These results are similar to the findings of other studies, who reported the highest plant height in mixture stand than pure stand [32,7,4].

3.1.2 Dry matter yield

Forage dry matter yield recorded at different growth stages of the monoculture and intercropping system is presented in Table 4. There were significant differences among the forage dry matter yield at different plant growth stages for the two consecutive years. The intercropping advantage compared to monoculture started to emerge at jointing and lasted until ripening stage where forage dry matter was significantly higher in alfalfa + oat and alfalfa + barley than sole oat, sole barley and sole alfalfa in 2015 and 2016. Sole oat, sole barley and sole alfalfa had forage dry matter yield decreased from 40.3 tha^{-1} , 46.2 tha^{-1} and 45.7 tha^{-1} in 2015 to 36.1 tha^{-1} , 44.9 tha^{-1} and 42.1 tha^{-1} respectively in 2016 at flowering stage, and from 41.7 tha^{-1} , 48.9 tha^{-1} and 46.2 tha^{-1} in 2015 to 38.5 tha^{-1} , 45.1 tha^{-1} and 42.4 tha^{-1} respectively in 2016 at ripening stage. This could be partly due to their continuous planting on the same area for two consecutive years. Forage intercropping systems showed significant advantages in dry matter yield over monocultures. This effect is likely related to niche differentiation in intercropping in spatial resources use; leaves for light and roots for water, which made intercrops able to utilise natural resources at different times during different growth stages. Intercrops were also effective at suppressing weeds [33]. There is a possibility that the alfalfa legume may have

released the fixed nitrogen to oat and barley cereal counterpart. Leguminous plant in mixtures of cereal + legume usually has direct benefits of nitrogen fixation in root nodules and contributed to soil fertility which was used by a companion as well as subsequent crops [33]. Previous studies reported a similar response of more forage yield produced from cereal- legume mixtures than sole cereal/legume [34,35,36].

3.2 Intercropping Competition Indices

3.2.1 Land Equivalent Ratio (LER)

LER's values were calculated for intercropping treatments in two consecutive growing seasons 2015 and 2016 (Table 8). All the intercrops showed LER greater than one. The maximum LER values were obtained from the alfalfa + oat intercropping system followed by alfalfa + barley. With the values of LER greater than one from the results, shows that intercropping is advantageous. LER ranged from 1.05 to 1.11 in 2015 and 1.16 to 1.24 in 2016. Therefore, 5% to 11% in 2015 and 16% to 24% more land should be used in monocropping in order to obtain the same yield of intercropping, which indicated the

advantage of the intercrops over monocrops in terms of the use of water, nutrients, carbon dioxide and light for plant growth. It was found that LER greater than one was primarily due to the nutrient cycling and increase in nitrogen content [37].

3.2.2 Relative crowding coefficient (K)

Relative crowding coefficient's values were calculated for intercropping treatments in two growing seasons 2015 and 2016 to determine competition experienced by legume (alfalfa) and cereals (oat or barley) when grown in a mixture and presented in Table 9. Relative crowding coefficient values were above one in 2015 and 2016 growing season for alfalfa + oat and alfalfa + barley mixtures. The K values of the cereals were higher than K values of legumes in intercropping system. From the results of this study, it appears that cereal forage was dominant probably due to the efficient utilisation of environmental resources. These findings are in agreement with previous studies where there were low relative crowding coefficient's values of the legume when the cereal was more competitive than the legume [38,39].

Table 3. Plant height of alfalfa, oat and barley monocultures and intercrops of alfalfa with oat and barley at different forage growth stages during two winter seasons 2015 and 2016

Cropping systems	Plant growth stages (cm)							
	Tillering (30d)		Jointing (50d)		Flowering (90d)		Ripening (110d)	
	2015	2016	2015	2016	2015	2016	2015	2016
Sole alfalfa	12.4b	14.5b	22.1b	20.9c	42.8d	39.1c	58.2d	56.8d
Sole oat	15.2ab	17.9a	31.4a	27.2b	71.2b	70.8b	92.3c	90.6c
Sole barley	16.6a	18.1a	30.1a	26.6b	68.5c	68.2b	91.5c	90.8c
Alfalfa + Oat	16.8a	18.3a	31.8a	33.1a	75.6a	79.1a	98.4a	99.7a
Alfalfa + Barley	17.2a	18.8a	32.3a	31.9a	72.8b	73.9b	95.8b	97.9b
Standard error (\pm)	1.07	1.12	0.75	1.53	1.17	1.84	1.97	1.89

d – Days after emergence. Means in the same column followed by different letters differ significantly at $P \leq 0.05$

Table 4. Forage yields of alfalfa, oat and barley monocultures and intercrops of alfalfa with oat and barley at different plant growth stages during two winter seasons 2015 and 2016

Cropping systems	Plant growth stages (tha^{-1})							
	Tillering (30d)		Jointing (50d)		Flowering (90d)		Ripening (110d)	
	2015	2016	2015	2016	2015	2016	2015	2016
Sole alfalfa	20.8a	22.8a	28.5b	28.8b	40.3d	36.1c	41.7c	38.5c
Sole oat	21.2a	23.2a	29.1a	29.9b	46.2c	44.9b	48.9ab	45.1b
Sole barley	20.8a	22.3a	27.2b	28.1b	45.7c	42.1b	46.2b	42.4b
Alfalfa + Oat	21.1a	23.7a	30.7a	32.5a	51.5a	53.7a	53.4a	55.7a
Alfalfa + Barley	21.9a	22.9a	29.6a	31.4a	48.7b	51.6a	50.9b	54.9a
Standard error (\pm)	0.32	0.89	1.44	1.85	0.96	1.07	1.56	1.91

d – Days after emergence. Means in the same column followed by different letters differ significantly at $P \leq 0.05$

Table 5. Land equivalent ratio for intercrops of alfalfa with oat and barley in two consecutive growing seasons (2015, 2016)

Intercropping	Land Equivalent Ratio (LER) 2015			Land Equivalent Ratio (LER) 2016		
	LER _{Legume}	LER _{Cereal}	LER _{Total}	LER _{Legume}	LER _{Cereal}	LER _{Total}
Alfalfa + Oat	0.77	0.34	1.11	0.85	0.39	1.24
Alfalfa + Barley	0.74	0.31	1.05	0.81	0.35	1.16

Table 6. Relative crowding coefficient for intercrops of alfalfa with oat and barley in two consecutive growing seasons (2015, 2016)

Intercropping	Relative Crowding Coefficient (K) 2015		Relative Crowding Coefficient (K) 2016	
	K _{Legume}	K _{Cereal}	K _{Legume}	K _{Cereal}
	Alfalfa + Oat	1.22	1.42	1.45
Alfalfa + Barley	1.29	1.38	1.55	1.67

3.2.3 Aggressivity (A_c)

Aggressivity values for intercropping treatments in two consecutive growing seasons 2015 and 2016 are presented in Table 10. Aggressivity has a similar trend as relative crowding coefficient. Cereal was the dominant species (A_c positive) in the alfalfa + oat and alfalfa + barley mixtures in 2015 and 2016 growing season. Considering all aggressivity values cereal showed significant advantages in aggressivity over legume. Cereal aggressivity values were positive while such values for legume forage were negative. This effect is likely related to spatial resource use; light, water and nutrients which made cereals able to be dominant species as measured by the positive value of aggressivity. Similar results were recorded by other researchers, who reported that in other intercrops (pea and barley, soya bean and palisade grass) the cereal values of the above indices were greater than for legume, indicating that cereal was more competitive than legume [40,41]. Similarly, the greater competitive ability of sorghum and barley to exploit resources in association with chickpea and faba bean has been reported by other studies [42,43].

3.3 Forage Quality

3.3.1 Crude Protein (CP)

Crude protein of forage mixtures and pure stands at four growth stages is presented in Table 5. Forage crude protein data indicated that there was a significant difference among treatments at different plant growth stages for two consecutive years. Forage crude protein was significantly higher in alfalfa + oat, alfalfa + barley, sole alfalfa than in sole oat and sole barley from tillering to

ripening stage in two growing seasons (2015 and 2016). Forage crude protein in both intercropping and monocropping gradually decreased from flowering to ripening stage in 2015 and 2016. Thus, the decrease in crude protein accumulation of forage at a late stage in intercropping and monocropping systems could be mainly because of maturity. Crude protein (CP) is often regarded to be the most important parameter of forage quality [44; 4]. The results showed that alfalfa + oat, alfalfa + barley, sole alfalfa forage produced higher crude protein content than their respective cereal counterparts; sole oats and sole barley, which could be the result of efficient utilisation of light, atmospheric nitrogen, moisture and nutrients. Legumes tend to have higher crude protein levels than cereal crops through biological fixation of nitrogen. Thus an overall improvement in crude protein is to be expected when legumes are intercropped with cereals. Crude protein improvement in legume-cereal intercropping has been reported by several studies, who reported a higher crude protein content relative to that of sole cereals [5, 45,46].

3.3.2 Acid Detergent Fibre (ADF)

Acid detergent fibre (ADF) recorded at different growth stages of two cropping systems; monoculture and intercropping are given in Table 6. ADF data revealed that there was a significant difference among treatments at different plant growth stages for two consecutive years. The highest ADF was obtained from sole barley followed by sole oat and sole alfalfa while lowest ADF was recorded with alfalfa + barley and alfalfa + oats intercropping system from tillering to ripening stage in two growing seasons (2015 and 2016). Acid detergent fibre (ADF) is an

important criteria for evaluating forage quality, represents the digestible energy that means as the ADF level increases, digestible energy levels decrease [46]. From the results of the study legume- cereal intercrops have low ADF values whereas monocultures have high values. This could be partly due to efficient utilisation of natural resources; light, atmospheric nitrogen, carbon dioxide, moisture and nutrients among the plants of different species legumes and cereals during different growth stages. The incorporation of legume with cereal could be of paramount importance to the low ADF of the forage mixture and subsequent soil health. These results are in line with the findings of other studies, who reported that combined cereal - legume forage had lower ADF concentration than sole cereal/legume [47,10,6].

3.3.3 Neutral Detergent Fibre (NDF)

Neutral detergent fibre (NDF) data regarding alfalfa, oat and barley monocultures and intercrops of alfalfa with oat and barley at four growth stages are presented in Table 7. NDF recorded at different plant growth stages for two consecutive years showed significant differences among the cropping system treatments. Maximum NDF was recorded in sole barley, sole oat, sole alfalfa and the lowest by alfalfa + oat and alfalfa + barley from tillering to ripening stage in two growing seasons (2015 and 2016). NDF concentration is a major component of forage quality and negatively correlated with dry matter intake, which means as NDF in the forage increases, animals will consume less forage [36]. From the results, it appears that NDF values are

Table 7. Aggressivity for intercrops of alfalfa with oat and barley in two consecutive growing seasons (2015, 2016)

Intercropping	Aggressivity (A) 2015		Aggressivity (A) 2016	
	A _{Legume}	A _{Cereal}	A _{Legume}	A _{Cereal}
Alfalfa + Oat	- 0.28	0.28	- 0.41	0.41
Alfalfa + Barley	- 0.21	0.21	- 0.35	0.35

Table 8. Crude protein (CP) in forage yield of alfalfa, oat and barley monocultures and intercrops of alfalfa with oat and barley at different plant growth stages over a two year period

Cropping systems	Plant growth stages (gkg ⁻¹)							
	Tillering (30d)		Jointing (50d)		Flowering (90d)		Ripening (110d)	
	2015	2016	2015	2016	2015	2016	2015	2016
Sole alfalfa	12.9a	13.5a	15.9a	16.8a	25.7b	25.5b	23.2ab	22.8bc
Sole oat	6.9b	7.5b	10.5b	11.4b	23.9bc	22.3c	21.1b	20.1c
Sole barley	7.2b	8.4b	9.9b	10.9b	19.1c	19.5d	18.4c	17.8d
Alfalfa + Oat	14.3a	13.7a	17.4a	17.3a	27.1a	29.9a	25.7a	27.8a
Alfalfa + Barley	13.1a	13.3a	16.2a	16.2a	26.5bc	27.2b	24.8ab	26.1a
Standard error (±)	1.75	1.58	1.06	1.77	0.86	1.15	0.97	1.25

d – Days after emergence. Means in the same column with different letters are significantly different ($P \leq 0.05$)

Table 9. Acid detergent fibre (ADF) in forage yield of alfalfa, oat and barley monocultures and intercrops of alfalfa with oat and barley at different plant growth stages over a two year period

Cropping systems	Plant growth stages (gkg ⁻¹)							
	Tillering (30d)		Jointing (50d)		Flowering (90d)		Ripening (110d)	
	2015	2016	2015	2016	2015	2016	2015	2016
Sole alfalfa	18.1ab	20.1b	23.3ab	22.1b	37.3b	39.4b	42.2a	43.2a
Sole oat	19.9ab	21.4b	21.6b	23.9a	39.1a	41.3ab	41.9a	42.8ab
Sole barley	21.7a	23.1a	25.5a	24.6a	38.9a	42.8a	42.2a	43.5a
Alfalfa + Oat	17.8b	18.6c	20.6b	21.8b	34.5c	32.8d	36.8b	37.4c
Alfalfa + Barley	18.5ab	19.2c	22.1b	22.2b	35.8c	35.6c	37.9b	38.7c
Standard error (±)	0.58	0.80	1.46	0.82	1.55	1.84	0.76	1.04

d – Days after emergence. Means in the same column followed by different letters differ significantly at $P \leq 0.05$

Table 10. Neutral detergent fibre (NDF) in forage yield of alfalfa, oat and barley monocultures and intercrops of alfalfa with oat and barley at different plant growth stages over a two year period

Cropping systems	Plant growth stages (gkg ⁻¹)							
	Tillering (30d)		Jointing (50d)		Flowering (90d)		Ripening (110d)	
	2015	2016	2015	2016	2015	2016	2015	2016
Sole alfalfa	35.6ab	33.8b	40.1ab	39.8b	55.2b	57.4a	61.5b	62.9b
Sole oat	39.1a	37.6a	44.8a	45.1a	56.8b	57.2a	63.9a	63.5b
Sole barley	38.7a	37.5a	43.4a	44.7a	59.1a	58.8a	64.2a	65.8a
Alfalfa + Oat	33.7b	36.8ab	39.8ab	40.2ab	51.5c	50.8b	55.4c	56.7c
Alfalfa + Barley	29.2c	36.4ab	40.5ab	41.7ab	52.8c	49.3b	55.2c	55.1c
Standard error (±)	0.52	1.76	1.28	0.95	1.85	1.96	1.65	1.81

d – Days after emergence. Means in the same column with different letters are significantly different ($P \leq 0.05$)

lower in cereal- legume intercrops and higher in sole cropping. The lower NDF of cereal- legume intercropping was probably due to the efficient utilisation of natural resources and minimum competition among the plants of different species during different growth stages. The results of this study are similar with previous studies, which investigated legume cereal mixtures and recorded the highest NDF in sole cropping and lowest in intercropping systems [48,11,2].

4. CONCLUSIONS

On the basis of results obtained in this study, intercropping of legume (alfalfa) with cereals (oat and barley) showed many benefits. Intercropping systems significantly increased plant height and forage yield compared with their respective monocrops from jointing to ripening stage in two growing seasons (2015 and 2016). Results obtained from intercropping competition indices indicated a superior advantage of legume-cereal mixtures because of better land use efficiency expressed as LER. Considering relative crowding coefficient (K) and aggressivity (A_c) values, cereal showed significant advantages over legume for two growing seasons (2015 and 2016). Legume – cereal intercrops improved forage quality in terms of crude protein yield (CP), acid detergent fibre (ADF) and neutral detergent fibre (NDF) concentrations than either forage species grown alone from tillering to ripening stage in two growing seasons (2015 and 2016). The results of the study show that with alfalfa + oat and alfalfa + barley, it is possible to produce greater forage yield and quality. Since there is little information in the literature on forage competition indices regarding intercropping systems, the results from this study may fill this gap. If the primary interest of the farmer is forage production, oat or barley forage

should be grown with alfalfa forage and harvested between jointing and flowering stage.

ACKNOWLEDGEMENTS

The authors gratefully acknowledge the financial support of this study by Research and Conferences Committee (RCC) of the National University of Lesotho (NUL). The valuable assistance of the staff of the Experimental Farm of the National University of Lesotho in this study is gratefully acknowledged.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Vandermeer J. The ecology of intercropping. Cambridge University Press, United Kingdom; 2006.
2. Takin F. Advantages of maize-cowpea intercropping over sole cropping through competition indices. J. Agric. Biod. Res. 2012;1:53-59.
3. Bothe H, Ferguson S, Newton W. Biology of the nitrogen cycle. Botanical Institute, University of Cologne, Koln, Germany; 2006.
4. Li Q, Sun J, Wei X, Christie P, Zhang F, Li L. Nitrogen fertilization in strip intercropping of maize with faba bean, wheat and barley. Plant Soil. 2011; 339:147-161.
5. Carr P, Horsley R, Poland W. Barley, oat, and cereal – pea mixtures as dryland forages in the northern Great Plains. Agron. J. 2004;96:677-684.

6. Strydhorst S, King J, Lopetinsky K, Harker K. Intercropping barley with faba bean, lupin, or field pea. *Agron. J.* 2008; 100:182-190.
7. Papastylianoul. Nitrogen fertilizer on barley and common vetch grown in various crop combinations. *J. Agric. Sci.* 2004; 142:41-48.
8. Seeiso S, McCrindle L. An investigation of the quality of meat sold in Lesotho. *J South African Veter. Assoc.* 2006;80(4):237-242.
9. Jensen E. N fixation and inter-specific for inorganic N in pea – barley intercropping. *J. Plant Soil.* 1996;182:25-38.
10. Canan T, Orak A. The effect of intercropping on yield of common vetch and oat. *J. Agric. Biol. Sci.* 2007;2:14-19.
11. Surve V, Arvadia M. Performance of fodder sorghum (*Sorghum bicolor* L.), maize (*Zea mays* L.) and cowpea (*Vigna Unguiculata* (L.) Walp.) under sole and intercropping systems. *Int. J. Agric. Res. Rev.* 2011;2:28-31.
12. Frame J. *Medicago sativa* L. Grassland Index; 2005.
13. Brink G, Hall M, Shewmaker G, Undersander D, Martin N, Walgenbach R. Lucern yield and quality in alley cropping. *Agron. J.* 2010;102 (4):1274-1282.
14. Radovic J, Sokolovic D, Markovic J. Alfalfa – most important perennial legume in animal husbandry. *Biotech. Animal Husban. J.* 2009;25:465-475.
15. Lauriault L, Kirksey R. Production of irrigated winter cereal grass – legume intercrops. *Agron. J.* 2004;96:352-358.
16. Ross S, King J, O'Donovan J, Spaner D. Yield potential on intercropping triticale and berseem clover. *Agron. J.* 2004;96(4): 1013-1021.
17. Erol A, Kaplan M, Kizilsimsek M. Oats and common vetch cultivated in mixture and pure stand. *Trop. Grassl.* 2009;43:191-196.
18. Johnston J, Wheeler B, McKinlay J. Production of forage from cereal/pea intercropping. Ontario, Min. Agric. Food. 1999.
19. Undersander D. Mixtures of legume and cereals. Univ. Wisc. Forage Team, Focus on Forage. 2003;5(7):1-2.
20. Stevens E, Armstrong K, Bezar H, Griffin W, Hampton J. Fodder oats: an overview (Chapter II). *Plant Prod. Prot. Series No. 33*; 2004.
21. Bagg J, Johnston, P. Summer seeding oats for forage. *Field Crop*; 2013.
22. Ecocrop. Ecocrop database. FAO; 2011.
23. Eskandari H, Ghanbari A, Javanmard A. cereals and legumes for production of fodder. *Not. Sci. Biol.* 2009;1(1):7-13.
24. Ditsch D, Bitzer M. Management of forage for livestock. University of Kentucky, Cooperative Extension Service, College of Agriculture; 2005.
25. Surber L, Abdel-Haleem H, Martin J, Hensleigh P, Cash D, Bowman J, Blake T. Mapping quantitative trait loci controlling variation in forage quality traits in barley. *Molecul. Breed.* 2011;28(2):189-200.
26. Mead R, Willey R. 1980. Yields and land equivalent ratio in plant mixtures. *Exp. Agric.* 1980;16:217-228.
27. Agegnehu G, Ghizaw A, Sinebo W. Forage performance in mixed cropping. *Eur. J. Agron.* 2006;25:202-207.
28. Association of Official Analytical Chemists (AOAC). *Official Methods of Analysis*, 11th ed. AOAC, Washington, DC125; 1980.
29. Goering H, Van Soest P. Analysis for quality of forage Handbook 237. U.S. Government Printing Office, Washington, DC;1970.
30. Statistical analysis system institute. SAS or STAT User's Guide, Version 9.1. SAS Institute, Cary, NC; 2003.
31. Sangoi L, Salvador R. Maize production at high plant densities. Ames: Iowa State Univ. Press. 1997;214-218.
32. Xu B, Li F, Shan L. Biomass of switchgrass and milk vetch under intercropping. *Eur. J. Agron.* 2008;28:485-492.
33. Intkhab H, Ahmad R. Intercropping barely with legumes. *Dawn*, Apr; 2008.
34. Ross S, King J, O'Donovan J, Spaner D. Intercropping forage species. *Agron. J.* 2004b;96:1719-1729.
35. Lithourgidis S, Dhima K, Vasilakoglou I, Dordas C, Yiakoulaki MD. Common vetch and wheat under mixed cropping. *Agron. Sustain. Develop.* 2007;27:95-99.
36. Sadeghpoura A, Jahanzada E, Esmaeilib A, Hosseinib M, Hashemia M. Benefits of barley and annual medic intercropping. *Field Crop Res.* 2013;148:43-48.
37. Singh M, Singh A, Singh S. Compatibility, productivity and economics of medicinal and vegetable crops intercropped with menthol mint. *Trop. Agric.* 2012;89(1):47-50.
38. Javanmard A, Nasab A, Javanshir A, Moghaddam M, Janmohammadi H.

- Intercropping of sorghum with three legumes. J. Food Agric. Environ. 2009; 7:163-166.
39. Dhima K, Lithourgidis A, Vasilakoglou I, Dordas C. Cereal and vetch intercrops in different seeding ratio. Field Crops Res. 2007;100:249-256.
40. Bedoussac L, Justes E. Assessing wheat – winter pea intercrop. Plant Soil. 2010; 330:37-54.
41. Weigelt A, Jolliffe P. Indices of plant competition. J. Ecol. 2003;91:707-720.
42. Lithourgidis A, Dordas C. Intercropping faba bean with different cereals in four seeding ratios. Crop Sci. 2010;50:2148-2158.
43. Mateus G, Crusciol C, Borghi E, Pariz C, Costa C, Silveira J. Effect of non – tillage system on sorghum and grass intercrops. Pesqui. Agropecu. Bras. 2011;46:1161-1169.
44. Lithourgidis A, Dhima K, Vasilakoglou I, Dordas C, Yiakoulaki M. Mixtures of vetch with oat and triticale for forage. Field Crop. Res. 2006;99:106-113.
45. Vasilakoglou I, Dhima K. Competition indices of pea with wheat in mixed cropping. Agron. J. 2008;100:1749-1756.
46. Lithourgidis A, Valchostergios D, Dordas C, Damalas C. Water and nitrogen content in pea – cereal intercrops. Eur. J. Agron. 2011b;34:287-294.
47. Aasen A, Baron V, Clayton G, Dick A, McCartney D. Intercropping spring cereals and field legume. Canadian. J. of Plant Sc. 2004;84(4):1051-1058.
48. Hauggaard-Nielsen H, Ambus P, Jensen E. Pea – barley as double-cropped. Field Crops Res. 2001;70:101-09.

© 2018 Mosebi et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history/26278>