



Assessment of X-ray Irradiation on the Morphological Performance of Sweet Potato (*Ipomoea batatas* [L.] Lam)

Ndem E. Edu ^a, Reuben C. Agu ^{a*}, Augusta A. Nwachukwu ^b,
Chinyere Osondu-Anyanwu ^c, Nkoyo A. Nkang ^c
and Covenant I. Egbaji ^a

^a Department of Genetics and Biotechnology, University of Calabar, Nigeria.

^b Department of Biotechnology, Federal University of Technology Owerri, Nigeria.

^c Department of Science Laboratory Technology, University of Calabar, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. Authors NEE and RCA designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors AAN, COA, NAN and CIE managed the analyses of the study and the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JABB/2023/v26i3624

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/98685>

Original Research Article

Received: 22/02/2023

Accepted: 26/04/2023

Published: 11/05/2023

ABSTRACT

Aim: Sweet potatoes (*Ipomoea batatas* [L.] Lam) were studied to assess the influence of x-ray irradiation on the morphological performance.

Place and Duration of Study: Samples were collected from National Root Crop Research Institute (NRCRI), Umudike, irradiation was done in the x-ray unit of a medical laboratory, planting and data analysis were done at the experimental farm and laboratory of the Department of Genetics and Biotechnology, University of Calabar.

Methodology: Stems of Sweet potatoes were grouped and non-control groups irradiated at different x-ray doses, planted and morphological parameters analyzed.

*Corresponding author: E-mail: chinedureuben@yahoo.com;

Results: The results showed no significant difference in the treatments for parameters such as number of leaves, leaf area, leaf length, plant height, leaf width and days to sprouting when kilo voltage (kV) was constant at 40kV and milli Amperes per second (mAs) varied from 1.6mAs to 3.2mAs. There was also no significant difference in the treatments for parameters such as leaf area, leaf length, leaf width and days to sprouting when mAs was constant at 1.6mAs and kilo voltage (kV) varied from 40kV to 100kV. The results also showed that the low doses of x-ray irradiation did not cause aberrations in the morphological performance of sweet potato.

Conclusion: These findings necessitate the need for adequate irradiation doses in the use of ionizing radiations on crops in order to maintain and improve their varieties.

Keywords: Sweet potato; x- ray; irradiation; treatments.

1. INTRODUCTION

Sweet potato (*Ipomoea batatas* [L.] Lam) is a dicotyledonous plant belonging to the family *convolvulaceae*, it is a field crop of major economic importance worldwide and the fourth most cultivated food crop after wheat, rice and maize [1]. It is a short duration crop that produces a large amount of calories and enriched with carbohydrates, protein, iron, magnesium, potassium and vitamin B & C [2]. It is widely used as source of food, feed and as a dietary supplement [3]. The current focus on energy production from biomass has led to the recognition of the potential of Sweet potato as a biomass species [4].

Irradiation-induced mutation breeding is effective in improving Sweet potato characters such as yield, starch and soluble sugar content, carotenoid content of storage roots and disease resistance [5,6]. Irradiation has also been successfully used for mutation breeding in various crops and ornamental plants [7] and has proven an adept means of encouraging the expression of recessive genes and producing new genetic variations [7,8,9].

Ahloowalia and Maluszynski [10] established the use of ionizing radiation, such as x-rays, gamma rays, neutrons and chemical mutagens for inducing variation. Ahloowalia et al. [11] presented the global impact of mutation-derived varieties on food production and quality enhancement. Al-Safadi et al. [12] established that low doses of gamma irradiation on potato (*Solanum tuberosum* L.) cultivars enhanced tuberization *in vitro* without genetic changes. Das et al. [13] obtained heat tolerant mutants in commercial potato cultivars through *in vitro* mutagenesis of *in vitro* propagated plantlets.

Ikram et al. [14], investigated the exposure of x-rays on leguminous seeds in combination with

Aerva javanica in the management of root rot fungal pathogens and on the growth of crop plants, seed of cowpea & mung bean and established that seeds treated with x- rays for 5 & 10 seconds showed significant enhancement in growth parameters and completely controlled the infection of root rot fungi (*Fusarium spp*, *Rhizoctonia solani* and *Macrophomina phaseolina*).

2. MATERIALS AND METHODS

2.1 Study Site

The research work was carried out at the experimental farm in the department of Genetics & Biotechnology, University of Calabar, Calabar.

2.2 Sample Collection and Processing

Stems of sweet potatoes were obtained from the National Root Crop Research Institute (NRCRI), Umudike and stem irradiation was done in the x-ray unit of a medical laboratory in Calabar metropolis.

2.3 X-ray Irradiation (Treatment) of Stems

Thirty (30) Stems of sweet potatoes (five for each group) were cut into length of 6cm and divided into groups and labeled appropriately. The controls (group 1) were not exposed to irradiation (treatment), while the other groups (2-6) groups were irradiated at different doses as shown in Table 1.

2.4 Planting of Stems

A land measuring 10 x 10 meters was used for the planting of the stems. The land was manually cleared, soil tilled and beds made. The stems were planted at 30cm apart and about 3cm deep in a randomized complete block design (RCBD). Weeding was done when the need arose.

Table 1. Irradiation schedule

Groups	First Trial	Second trial
1	0kV/0mAs	0kV/1.6mAs
2	40kV/1.6mAs	40kV/1.6mAs
3	40kV/2.0mAs	55kV/1.6mAs
4	40kV/2.4mAs	70kV/1.6mAs
5	40kV/2.8mAs	85kV/1.6mAs
6	40kV/3.2mAs	100kV/1.6mAs

*kV was kept constant in the first trial while mAs varies
mAs was kept constant in the second trial while kV varies;
kV = kilo Voltage; mAs = milli Amperes per second

2.5 Data Analysis

Data such as days to sprouting, plant height, leaf length, leaf width, leaf area and number of leaves per plant was analyzed using Analysis of Variance (ANOVA) and means separated with the Least Significant Difference (LSD) at 0.05 probability level.

3. RESULTS AND DISCUSSION

The results showed the morphology of sweet potato when kilo voltage (kV) was constant and milli Amperes per second (mAs) varied (Table 2). The results for the number of leaves showed that there was no significant difference in the treatments and in the block ($P>0.05$). The mean number of leaves ranged between 5.8 ± 0.5 – 7.7 ± 1.1 with treatments from group 2 having the highest mean value. The leaf area showed no significant difference ($P>0.05$) in the treatments and the block. The mean leaf area varied from 192.9 ± 32.06 – 334.9 ± 43.86 with treatments from group 3 having the highest mean value. The leaf length were not significantly different in the block and also in the treatments ($P>0.05$). The leaf length also

varied from 7.1 ± 0.54 – 8.2 ± 0.8 with treatments from group 1 (control) having the highest mean value. The plant height showed significant difference in the block ($P<0.05$) but no significant difference in the treatments ($P>0.05$). The mean for the plant height ranged between 10.02 ± 0.60 – 10.9 ± 0.98 with treatments from group 1 (control) having the highest mean value.

The leaf width showed no significant difference in the block and also in the treatments ($P>0.05$). The leaf width also ranged from 6.1 ± 0.52 – 7.2 ± 0.22 with treatments from group 5 having the highest mean value. The days to sprouting showed no significant difference in the block and also in the treatments. Statistically no significant difference was observed in the days of germination. The parameters for the days of sprouting varied from 16.5 ± 0.89 – 20.5 ± 2.33 with treatments from group 2 having the highest mean value.

The results also showed the morphology of sweet potato when milli Amperes per second (mAs) was constant and kilo voltage (kV) varied (Table 3). The results for the number of leaves showed that the treatments had significant difference ($P<0.05$) and there was no significant difference in the block ($P>0.05$). The mean number of leaves ranged between 8.8 ± 0.54 – 12.0 ± 0.58 with treatments from group 2 and 5 having the highest mean value. The leaf area showed no significant difference ($P>0.05$) in the treatments and the block. The mean leaf area varied from 78.08 ± 3.21 – 92.63 ± 4.78 with the mean leaf area of the control (Group 1) higher than that of the treatments. The leaf length were not significantly different in the block and also in the treatments ($P>0.05$).

Table 2. Mean values for studies on x-ray irradiation on morphological performance of sweet potato at constant kV and variable mAs

Parameters	1	2	3	4	5	6	LSD
No of leaves	7 ± 0.6	7.7 ± 1.1	6.3 ± 0.7	7.3 ± 0.66	5.8 ± 0.5	6.7 ± 0.66	3.02
Leaf area	192.9 ± 32.06	266.6 ± 25.45	334.9 ± 43.86	227.3 ± 31.09	217.29 ± 54.89	285.43 ± 121.86	243.2
Leaf length	8.2 ± 0.8	7.4 ± 0.47	7.3 ± 0.69	7.7 ± 0.36	7.1 ± 0.54	7.3 ± 0.63	2.40
Plant height	10.9 ± 0.98	10.7 ± 0.71	10.52 ± 0.82	10.8 ± 0.74	10.02 ± 0.60	10.08 ± 1.01	2.86
Leaf width	6.1 ± 0.52	6.6 ± 0.58	6.1 ± 0.48	7.0 ± 0.39	5.6 ± 0.20	7.2 ± 0.22	1.82
Days to Sprouting	17.2 ± 1.85	20.5 ± 2.33	18.3 ± 2.47	16.5 ± 0.89	17.8 ± 0.60	19.8 ± 0.79	8.80

*1= 0kV/ 0mAs; 2 = 40kV/ 1.6mAs; 3 = 40kV/ 2.0mAs; 4 = 40kV/ 2.4mAs; 5 = 40kV/ 2.8mAs; 6 = 40kV/ 3.2mAs

Table 3. Mean values for studies on x-ray irradiation on morphological performance of sweet potato at constant mAs and variable kV

Parameters	1	2	3	4	5	6	LSD
No of leaves	8.8±0.54	12.0±0.58	10.83±0.94	9.5±0.43	12.0±0.58	11.0±0.67	2.67
Leaf area	92.63±4.78	84.53±4.30	81.31±5.89	85.53±12.36	92.04±10.87	78.08±3.21	31.66
Leaf length	10.7±0.39	9.6±0.59	9.5±0.60	9.1±1.12	10.2±0.83	8.1±0.52	2.80
Plant height	21.2±1.85	16.7±0.49	15.3±2.08	20.6±1.23	22.7±1.10	25.4±1.11	5.66
Leaf width	8.7±0.47	8.9±0.46	8.6±0.32	9.3±0.32	8.9±0.71	9.7±0.38	1.74
Days to sprouting	21.5±2.30	19.3±0.80	22.7±2.47	20.2±1.48	20.3±1.49	24.5±2.35	8.80±0.82

*1= 0kV/ 0mAs; 2 = 40kV/ 1.6mAs; 3 = 55kV/ 1.6mAs; 4 = 70kV/1.6mAs; 5 = 85kV/1.6mAs; 6 = 100kV/1.6mAs

The leaf length also varied from $8.1 \pm 0.82 - 10.7 \pm 0.39$ with leaf length of the control (Group 1) higher than that of the treatments. The plant height were significantly different in the treatments ($P < 0.05$) and not significantly different in the block ($P > 0.05$). The mean for the plant height ranged between $15.3 \pm 2.08 - 25.4 \pm 1.11$ treatments from group 6 having the highest mean value. The leaf width showed no significant difference in the block and also in the treatments ($P > 0.05$). The mean for the leaf width ranged from $8.6 \pm 0.32 - 9.7 \pm 0.38$ with treatments from group 6 having the highest mean value. The days to sprouting showed no significant difference in the block and also in the treatments ($P > 0.05$). The mean days to sprouting ranged from $19.3 \pm 0.80 - 24.5 \pm 2.35$ with treatments from group 6 having the highest mean value.

The influence of x-ray irradiation on the morphological performance of Sweet potato (*Ipomoea batatas* [L.] Lam) when kilo voltage (kV) was constant at 40kV and milli Amperes per second (mAs) varied from 1.6mAs to 3.2mAs showed no significant difference in the treatments for parameters such as number of leaves, leaf area, leaf length, plant height, leaf width and days to sprouting. This means that the x-ray doses did not cause any aberrations in the morphological performance of sweet potato. This is in line with the study of Kehinde et al. [15] who reported that small doses of x-ray may stimulate cellular activities and growth while higher doses may cause higher aberrations. The study is also in conformity with Andrew et al. [16] who reported that low x-ray irradiation doses could induce variability and enhance vegetative growth and yield of plants using cowpea as a test plant.

The influence of x-ray irradiation on the morphological performance of Sweet potato (*Ipomoea batatas* [L.] Lam) when milli Amperes per second (mAs) was constant at 1.6mAs and

kilo voltage (kV) varied from 40kV to 100kV showed no significant difference in the treatments for parameters such as leaf area, leaf length, leaf width and days to sprouting. This also means that the low x-ray doses did not cause any aberrations in the morphological performance of sweet potato. This is also in line with Akpaniwo et al. [17] who evaluated the field performance and morphological variation of fluted pumpkin treated with various doses of x-ray radiation and observed that high doses of x-ray radiation reduce germination and survival percentage in the M1 generation.

4. CONCLUSION

The study evaluated the morphological performance of Sweet potato (*Ipomoea batatas* [L.] Lam) irradiated with x-ray which is one of the ionizing radiations that have been used in improving the varieties and characters of important crops such as Sweet potato (*Ipomoea batatas* [L.] Lam).

From the study, low doses of x-ray irradiation did not cause any aberration in the morphological performance of sweet potato. Therefore adequate irradiation doses are necessary in the use of ionizing radiations on Sweet potato and other crops in order to maintain or improve their varieties.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Jones DA, Hyman LJ, Tumeseit M, Smith P, Perombelon, MC. Blackleg potential of potato seed: determination of tuber contamination by *Erwinia carotovora* sub species *Atroseptica* by

- immunofluorescence colony staining and stock and tuber sampling. *Annals of Applied Biology*. 1994;124(3):557-568.
2. Nasrin S, Hossain MM, Khatun A, Alam MF, Mondal, MRK. Induction and evaluation of somaclonal variation in potato (*Solanum tuberosum* L.). *Online Journal of Biological Science*. 2003;3(2): 183-190.
 3. Woolfe JA. Sweet Potato: An untapped food resource. New York UK: Cambridge University Press; 1992.
 4. Ziska LH, Runionb GB, Tomeceka M, Priorb SA, Torbetb HA, Sichera R. An evaluation of cassava, sweet potato and field corn as potential carbohydrate sources for bioethanol production in Alabama and Maryland. *Biomass and Bioenergy*. 2009;33(11):1503-1508.
 5. Kukimura H. Mutation breeding in sweet potato and tuber crops. *Gamma Field Symptom*. 1986;25:109-130.
 6. Wang Y, Wang F, Zhai H, Liu Q. Production of a useful mutant by chronic irradiation in sweet potato. *Scientia Horticulturae*. 2007;111(2):173-178.
 7. Song HS, Kang SY. Application of Natural Variation and Induced Mutation in Breeding and Functional Genomics: Papers for International Symposium; Current Status and Future of Plant Mutation Breeding. *Korean Journal of Breeding in Science*. 2003;35(1):24-34.
 8. Schum, A. Mutation breeding in ornamentals: An efficient breeding method. *Acta Horticulturae*. 2003;612(612):47-60.
 9. Yoon KE, Im BG, Park YH. Effect of gamma radiation on seed germination and androgenesis in *Nicotiana tabacum* L. *Korean Journal of Breeding in Science*. 1990;21(4):256-262.
 10. Ahloowalia BS, Maluszynski M. Induced mutations – A new paradigm in plant breeding. *Euphytica*. 2001;118:167–173.
 11. Ahloowalia BS, Maluszynski M, Nichterlein K. Global impact of mutation-derived varieties. *Euphytica*. 2004;135:187-204.
 12. Al-Safadi B, Ayyoubi Z, Jawdat D. The effect of gamma irradiation on potato micro-tuber production *in vitro*. *Plant Cell, Tissue and Organ Culture*. 2000;61:183–187.
 13. Das A, Gosal SS, Sidhu JS, Dhaliwal HS. Induction of mutations for heat tolerance in potato by using *in vitro* culture and radiation. *Euphytica*. 2000;114:205–209.
 14. Ikram N, Dawar S, Imtiaz F. X rays exposure on leguminous seeds in combination with *Aerva Javanica* parts powder for the promotion of growth and Management of root rot fungal pathogens. *European Journal of Botany Plant Sciences & Pathology*. 2015;2(2):1-10.
 15. Kehinde DO, Ogunwenmo KO, Ajeniya B, Ogunowo AA, Onigbinde AO. Effects of x-ray irradiation on growth physiology of *Arachis Hypogaea* (Var. Kampala). *Chemistry International*. 2017;3(3):296-300.
 16. Andrew OV, Egbucha KC, et al. Effect of x-ray Irradiation on the Growth and Yield Parameters of four Cowpea (*Vigna unguiculata* (L.) Walp) Genotypes. *NIPES Journal of Science and Technology Research*. 2021;3(4):58-68.
 17. Akpaniwo EG, Ittah MA, Iwo GA. Field Performance and Morphological Variations of x-ray Induced M2 Mutant Lines of Fluted Pumpkin (*Telfairia occidentalis*, Hook. F.). *International Journal of Research in Agriculture and Forestry*. 2015;2(10):12-16.

© 2023 Edu et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/98685>