



# Gastrointestinal Helminth Parasites of *Malapterurus electricus* from Anambra River, Nigeria

C. C. Nwadike <sup>a\*</sup>, P. C. O. Ilozumba <sup>a</sup>, E. C. Amaechi <sup>b</sup>,  
O. A. Okeke <sup>a</sup>, I. O. Nnatuanya <sup>a</sup> and P. N. Osaji-Nwafili <sup>c</sup>

<sup>a</sup> Department of Zoology, Parasitology Unit, Nnamdi Azikiwe University, Awka, Nigeria.

<sup>b</sup> Department of Zoology, University of Ilorin, Nigeria.

<sup>c</sup> Department of Animal and Environmental Biology, Delta State University, Abraka, Nigeria.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/AJFAR/2023/v22i6586

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/100568>

Original Research Article

Received: 22/03/2023

Accepted: 26/05/2023

Published: 03/06/2023

## ABSTRACT

**Aim:** One of the constraints of fish in the wild for sustainable productivity is parasitic infections. The study was carried out to determine the gastrointestinal helminth parasites associated with *Malapterurus electricus* and their prevalence in relation to sex and size.

**Study Design:** The study was a survey done to establish the type of gastrointestinal helminth parasites found in *M. electricus*.

**Place and Duration of Study:** The specimens were collected from Anambra River between August and December 2018.

**Methodology:** A total of thirty-two (32) fishes were examined through dissection and observation of the alimentary canal under the objective lens of the microscope. The chi-square test was employed to determine the possible association between parasite prevalence, the sexes, and size.

\*Corresponding author: Email: nc.chidiogo@unizik.edu.ng;

**Results:** The overall prevalence of parasites was 56.25%. The parasites recovered were, *Electrotaenia malapteruri* (50.00%) and *Tenuisentis niloticus* (6.25%). The male specimens (18) recorded a prevalence of 66.67% while the female (14) recorded a prevalence of 42.86%. There was no significant difference in the prevalence of gastrointestinal helminths among gender ( $P=0.178$ ,  $P>0.05$ ). The length group, 30 – 39cm had a 100% prevalence. Thus, a significant difference between the fish size and parasite prevalence was observed. The weight groups, 100 - 199g, and 400 - 499g also had a 100% prevalence. However, no significant relationship was observed between fish weight and parasite prevalence. The intestine was infected with parasitic worms of the cestode and acanthocephalan species.

**Conclusion:** This study showed that there is high prevalence of gastrointestinal helminth parasites in *M. electricus* and this may possibly affect the fish quality.

**Keywords:** *Malapterurus electricus*; freshwater fish; intestinal parasites; helminth; Anambra river; Nigeria.

## 1. INTRODUCTION

The increased necessity to raise fish for human consumption as a source of nutrition for the thronging, quickly expanding populations in developing countries has necessitated a stepped-up investigation of the parasite fauna of African freshwater fish. However, one of the challenges confronting aquaculture is parasite [1]. Parasites represent an important source of economic losses for aquaculture in terms of reduced fish growth and increased mortality, and also in terms of investments in the farming practices and chemicals necessary for their prevention [1].

One of the major fishes in aquaculture is *Malapterurus electricus*. *Malapterurus electricus* can be found in Western and Central tropical Africa and along the Nile River. *Malapterurus electricus* comprising three species may also be found in all of Africa's major freshwater systems [1]. Their soft, puffy bodies give them the appearance of a hard sausage driven by ostraciform movements while they swim [2]. *M. electricus* most noticeable feature is its great electrogenic capacity. Its pectoral muscle gave rise to the electric organ. This organ likewise surrounds the fish's body for the majority of its length and may discharge up to 350 volts (in a 500mm fish) [1]. The electric organs of *M. electricus* have been used in studies of neuronal metabolism, axonal transport, and transmitter release, yet information on the parasite fauna of *M. electricus* seems to be scarce when compared to other fish species.

In some parts of Nigeria, parasite documentation on *M. electricus* includes *Electrotaenia malapteruri*, *Tenuisentis niloticus*, and *Nilonema* species [3] in Lagos; *Procamallanus* species in

Benin City; *Procamallanus laevisconchus* [4]; *Henneguya* species (Protozoa), *Clinostomum* species (Trematoda), *Diphyllbothrium latum* (cestode) and *Capillaria* species, *Contracaecum* species, *Eustrongylides* species (Nematoda) [1] in River Benue. [5] at Rivers Niger-Benue Confluence also recovered protozoan ciliate (*Trichodinids*), cestodes (*Monobothrioides woodlandi*, *E. malapterurid*, and *Proteocephalus largoploglotis*) and nematodes (*P. laevisconchus*, *Rhabdochona congolensis*, and *Camallanus* species). However, there is a paucity of information on the parasites of *M. electricus* from Rivers in Anambra State. Therefore, this study aims at providing information on the gastrointestinal helminth parasites of *M. electricus* from the Anambra River, Nigeria because of their importance in the artisanal fisheries of the basin.

## 2. MATERIALS AND METHODS

### 2.1 Study Area

The study was conducted in the Anambra River, Southeastern, Nigeria. It is a major tributary of the River Niger. The water emerges from a somewhat inaccessible point near Ankpa in the Kogi State of Nigeria, crosses the Kogi/Anambra State boundary a bit north of Ogorugu, and then meanders through the Ogorugu station to Otuocha, then flows down to its confluence with the Niger at Onitsha [6]. Anambra River Basin lies between latitudes  $6^{\circ}10'$  and  $7^{\circ}8'$  N and longitudes  $6^{\circ}30'$  and  $7^{\circ}15'E$  (mapcarta.com). The basin has an annual rainfall between 150cm-200cm. The area has a low altitude of under 1000 above sea level. Temperatures are uniformly high with a small annual range of 5-10°C. The study area is one of the richest areas for agricultural and fishery production in Nigeria

[7], thus agriculture and fishing form the dominant occupations of the local people. These two major economic activities are closely geared to the two seasons of the year (wet and dry seasons).

## 2.2 Collection of Fishes

Thirty-two (32) *M. electricus* (18 males and 14 females) were purchased from local fishermen at Otuocha market, Anambra State. The fishes were transported to the Parasitology Laboratory of the Department of Zoology, Nnamdi Azikiwe University. The weights, standard lengths, and total lengths of the fish were recorded. The weights were recorded to the nearest gram using Adams electronic weighing balance; model AQP 1600 while the lengths of the fishes were determined using a measuring board calibrated in centimeters. The sexes of the fish were determined by examination of their papillae and were immediately subjected to parasitological examinations.

## 2.3 Examination of Gastrointestinal Tract for Helminths

Finally, we conducted internal examination of the fish. For this analysis, the body wall was cut from the ventral side, the intestinal parts were opened and examined for gastrointestinal parasite investigation, especially for Cestodes and Nematode worms. The detection and counting of intestinal parasitic eggs, larvae and cysts was performed using a compound microscope (objective 10X) [8]. The recognition of the worms was enhanced by the wriggling movements on emergence.

## 2.4 Identification of Parasites

Parasites seen were picked up with a small paintbrush. Cestode parasites were relaxed in the water and preserved in 70% alcohol. Acanthocephalans were shaken vigorously in cold 4% formaldehyde until they died. They were also preserved in cold 4% formaldehyde. The number of parasites per fish was recorded along with the site/location from which each parasite

was collected. The parasites were identified using keys developed by Yorke and Mapplestone [9], Yamaguti [10], Cheng [11], Soulsby [12], Williams and Jones [13] and Paperna [14].

## 2.5 Statistical Analysis

Infection statistics of Bush et al. [15] were used for the determination of prevalence, mean intensity, and mean abundance. Chi-Square was done to determine the relationship between fish sex, length, weight, and helminth infection at a 95% significant level ( $P < 0.05$ ).

## 3. RESULTS

Out of the 32 *M. electricus* examined, 18 (56.25%) were infected while 14 (43.75%) were uninfected. A total of 112 parasites belonging to one species of cestode (*Electrotaenia malapteruri*) and one species of acanthocephalan (*Tenuisentis niloticus*) were recovered from the fishes. Among the two parasites, *Electrotaenia malapteruri* recorded a prevalence of 50.00% while *T. niloticus* recorded a prevalence of 6.30%. Both parasites were recovered from the intestine (Table 1).

The prevalence of gastrointestinal helminths was higher in males 12(66.67%) than in females 6(42.86%). However, there was no significant difference in the prevalence among gender ( $P = .178, P > 0.05$ ) (Table 2).

The length groups 10 - 19cm had a prevalence of 50.0%, and 20 - 29cm (54.0%), while the highest length groups 30 - 39cm had a 100% prevalence. A significant difference between size and parasite prevalence was observed (0.866;  $P < 0.05$ ) (Table 3).

The weight groups 0 – 99g had a prevalence of 66.67%, 100 – 199g (100%), 200 – 299g (16.67%), and 300 – 399g (75%) while the highest weight groups 400 – 499g recorded 100% prevalence of infection. However, no significant relationship between weight and parasite prevalence was recorded (13.714;  $P > 0.008$ ) (Table 4).

Table 1. Parasite species spectrum in *M. electricus*

Parasite species	Number examined	Number infected	Prevalence (%)	N. P. R	M.I.I	M.A
<i>Electrotaenia malapteruri</i>	32	16	50.00	110	6.87	3.43
<i>Tenuisentis niloticus</i>	32	2	6.30	2	1.00	0.10

\*N.P.R = Number of Parasites Recovered; \*M.I.I = Mean Intensity of Infection; \*M.A = Mean Abundance

**Table 2. Relationship between sex and prevalence of gastrointestinal helminth infection**

	Male	Female	Combined sex
Number examined	18	14	32
Number infected	12	6	18
Prevalence (%)	66.67	42.86	56.25

$\chi^2 = 1.814, df = 1, P\text{-value} = .178$

**Table 3. Relationship between size distribution and gastrointestinal helminth infection**

Body size (cm)	10 – 19	20 – 29	30 -39
Number examined	6	24	2
Number infected	3	13	2
Prevalence	50.00	54.00	100.00
N.P.R	24	51	37
M.I.I	8.00	3.92	18.50
M. A	4.00	2.13	18.50

$\chi^2 = 0.866; P < 0.05$

**Table 4. Relationship between fish weight and gastrointestinal helminth infection**

Body weight (g)	0 - 99	100 - 199	200 - 299	300 - 399	400 - 499
Number examined	6	4	12	8	2
Number infected	4	4	2	6	2
Prevalence (%)	66.67	100.00	16.67	75.00	100.00
Mean intensity	6.00	2.00	1.33	4.50	18.50

$\chi^2 = 13.714, df = 4, P\text{-value} = 0.008$

#### 4. DISCUSSION

The recovery of *E. malapteruri* and *T. niloticus* in this study is not surprising as they have been recorded previously from the fish species by Iyaji and Eyo [5] in Rivers Niger – Benue Confluence, [3] in Lekki Lagoon, and [6] in Anambra River. Surprisingly, [1] in their study of Ecto and intestinal parasites of *M. electricus* from upper river Benue did not recover any of these parasites. The reason may be attributed to the location and sampling.

Of the two parasite species recorded, *E. malapteruri* had the highest prevalence of 50.00%. Numerous fish species have been known to be infected by cestode (tapeworm) parasites, which are widely distributed in all major freshwater systems of Africa and demonstrate a high degree of host specificity [5]. According to Cheng [16], proteocephalid cestode infection of fish hosts involves a complex life cycle of intermediate and definitive hosts. This complex life cycle may be the reason *E. malapteruri* has been characterized as being host-specific. Khalil and Polling [17] and Alain et al. [18] also affirmed the host specificity of *E. malapteruri*.

The recovery of *T. niloticus* agrees with Akinsanya [3] who made the first scientific report on *T. niloticus* in *M. electricus*. This may be due to the omnivorous feeding habits of *M. electricus*. It is also of interest to note that the host specificity of acanthocephalans is variable [3] as they tend to develop via one or more intermediate hosts that fish can serve as.

The relationship between host sex and helminth infection showed that males had a higher prevalence than females although infections were not significant. Similar observations have been made in other studies [1,19-22]. The reason for the difference in the incidence of infection could be due to differential feeding either by quantity or quality of food eaten. It may also be a result of different degrees of resistance and infection [1]. Thus, the infection could be accidental.

The significant relationship recorded between size (length) and helminth infection was also reported by Iyaji and Eyo [5], and Akinsanya et al. [3]. The increase in parasitism with size may be because older fishes have a longer time to pick up parasites than younger ones since they supposedly feed more [5]. This may also be attributed to the fact that older fishes engage in

more activity requiring a high rate of metabolism and so feed more. Although there was a significant difference between size (length) and helminth infection, infections were not significant with the weight of *M. electricus*. However, the heavier fishes had the highest mean intensity of 18.50. This could be attributed to the random selection of the fish.

## 5. CONCLUSION

The finding from the present study showed that there was a high prevalence of helminth parasites in *M. electricus*. The total number of parasites recovered was also high. Humans who eat undercooked parasites infected fishes are at risk of the infection. Further studies are still required to establish factors responsible for the abundance of parasites in fish of the Anambra River particularly *M. electricus*.

## ETHICAL APPROVAL

This study followed guidelines for the care and use of experimental animals established by the Animal Care and Use Committee of Nnamdi Azikiwe University, Awka, Nigeria, for the control and supervision of experiments on animal (RE: NAU/AREC/2023/00025).

## ACKNOWLEDGEMENT

We wish to acknowledge the assistance of the fishermen who made the fishes available and also the Laboratory Technologists in the Department of Zoology, Nnamdi Azikiwe University for their assistance during the research.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Omeji S, Tihamiyi LO, Annune PA, Solomon SG. Ecto and Intestinal parasites of *Malapterurus electricus* from Upper river Benue. Journal of Global Biosciences. 2014;3(6):895-903.
2. Lissmann HW. On the function and evolution of electric organs in fish. J. Exp. Biol. 1958;35:156 – 191.
3. Akinsanya B, Otubanjo OA, Hassan AA. Helminth parasites of *Malapterurus electricus* (Malapteruridae) from Lekki Lagoon, Lagos, Nigeria. Journal of American Science. 2007;3(3):1-6.
4. Obano EE, Odiko AE, Ebeh DO. Helminth parasitic infection of fishes from Okhuaihe river Benin City, Nigeria. Bioscience Research Combinations. 2010;22(3):129-135.
5. Iyaji FO, Eyo JE. Parasites of *Malapterurus electricus* (Gmelin, 1789, Siluriformes, Malaptermridae) at Rivers Niger-Benue Confluence, Nigeria. Journal of Science and Multidisciplinary Research. 2014;2(2): 33-40.
6. Azugo WI. Ecological studies of the helminth parasites of the fish of the Anabra river system. M. Phil. Thesis, University of Nigeria, Nsukka. 1978:100.
7. Mutter NES. (ed.). Report of the lower Niger survey appraisal mission. London, Overseas Development Administration, PROP/14/73; 1978.
8. Obaid M, Hussein NF, Obed TM, Boundenga L. Common carp (*Cyprinus carpio*) parasites diversity and prevalence in Erbil aquacultures: Gills, skin and intestinal infections. Iran J Vet Sci Technol. 2021;13(1):34-41. Available:<https://doi.org/10.22067/ijvst.2021.64304.0> Available:[https://ijvst.um.ac.ir/article\\_39981.html](https://ijvst.um.ac.ir/article_39981.html)
9. Yorke W, Mapplestone PA. The nematode parasites of vertebrates. London, S. & A. Churchill Publishers; 1926.
10. Yamaguti S. Systema helminthum, nematodes of vertebrates. Interscience Publishers Inc. New York, USA. 1961:1261.
11. Cheng T. General parasitology. Academic Press, New York, USA. 1973:965.
12. Soulsby ELS. Helminths, arthropods and protozoans of domesticated animals. 7<sup>th</sup> Edition. Bailliere Tindall, London, UK. 1982:809.
13. Williams H, Jones A. Parasitic worms of fish, Taylor and Francis, Bristol, UK. 1994:593.
14. Paperna I. Parasites, infections, and diseases of fishes in Africa – An update, CIFA Tech Pap. 1996;31:1-220.
15. Bush AO, Lafferty KD, Lots JM, Shostar AW. Parasitology meets Ecology in its term. Journal of Fish Biology. 1997:575 – 583.
16. Cheng TC. General Parasitology. 2nd ed. Harcourt Brace & Company, Asia PTE Ltd 583 Orchard Road. 09-01 Forum. Singapore; 1999.

17. Khalil LF, Polling. Checklist of the helminth parasites of African freshwater fishes. University of the North Republic of South Africa, South Africa. 1997:161.
18. Alain de Chambrier, Scholz CT, Mohammed HI. Redescription of *Electotaenia malapteruri* (Fritsch, 1886) (Cestoda: Proteocephalidae) a parasite of *Malapterums electricus* (Siluriformes: Malapteruridae) from Egypt. Syst. Parasitol. 2004;57(2):97-109.
19. Anosike JC, Omoregie Ofojekwu PC, Nweke IE. A survey of helminth parasites of *Clarias gariepinus* in Plateau state, Nigeria. Journal of Aquatic Sciences. 1992;7:39–43.
20. Oniye SJ, et al. Helminth parasite of *Clarias gariepinus* in Zaria, Nigeria. African Journal of Aquatic Science. 2004;19:71-76.
21. Omeji S, Solomon SG, Uloko C. Comparative Study on the endo-parasitic infestation in *Clarias gariepinus* collected from earthen and concrete ponds in Makurdi, Benue State, Nigeria. IOSR; 2013
22. Emere MC, Egbe NEL. Protozoan parasites of *Synodontis clarias* (A freshwater fish). Best Journal. 2006;3(3): 58-64.

© 2023 Nwadike et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/100568>