



Economic Assessment against Shoot and Fruit Borer (Lepidoptera: Noctuidae) on Okra at Uttar Pradesh, India

**Chitteti Divya^{a*}, Kurru Charitha^a
and Naidu Venkata Girish Reddy^a**

^a Department of Entomology, Faculty of Agriculture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj-211007, Uttar Pradesh, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i82042

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/101057>

Short Research Article

Received: 01/04/2023

Accepted: 01/06/2023

Published: 08/06/2023

ABSTRACT

The field trial was conducted during *Kharif* 2022 to evaluate the cost benefit ratio by using different insecticidal applications viz., NSKE 10% T₁ @ 10gm/L, *Beauveria bassiana* 1.15% WP (1x10⁸ CFU/gm) T₂ @ 5gm/L, Spinosad 45 SC T₃ # 0.5ml/L, *Metarhizium anisopliae* 4% WSP (1x10⁸ CFU/gm) T₄ @ 5ml/L, Imidacloprid 17.8 SL T₅ @ 0.3 ml/L, Nisco sixer plus T₆ @ 1.5ml/L, Neem oil 5% T₇ @ 5ml/L and untreated control T₈ in three replications against shoot and fruit borer, *Earias vittella* (Lepidoptera: Noctuidae), management on okra. Observations were taken on percent infestation of shoot and fruit on day before, 7DAS and 14DAS after two sprays Results revealed that highest yield is recorded in Spinosad 45 SC (101.4 q/ha), followed by nisco sixer plus (82.2 q/ha), Imidacloprid 17.8 SL (72.5 q/ha). Insecticidal treatment with Spinosad 45 SC with 1:5.66 had the highest cost benefit ratio, followed by Nisco sixer plus (1:4.99).

*Corresponding author: E-mail: chittetidivya98@gmail.com;

Keywords: Cost benefit ratio; okra; shoot and fruit borer; yield.

1. INTRODUCTION

Okra, sometimes referred to as ladies' finger, bhindi, bamia, and gumbo, is an annual vegetable that belongs to the Malvaceae family. Its scientific name is *Abelmoschus esculentus* (L.) Moench. The "Queen of Vegetables" is okra. For its tender green fruits, it is valued [1]. With an area of 1148.0 thousand hectares and an annual production of 6346 million tonnes, India leads the globe in okra production, producing 5784.0 thousand tonnes (72% of the world's population). Okra output and productivity in the Uttar Pradesh region are 48.2 thousand ha, 177.26 thousand tonnes, and 8 tons/ha, respectively [2].

One of the most dangerous pests of okra is the shoot and fruit borer, also known as *Earias insulana* and *Earias vittella*. The larva bore into the okra's terminal developing shoots, floral buds, flowers, and fruits, causing the afflicted shoots, sensitive leaves, and floral buds and flowers to shed heavily. The fruit that has been infected develops malformations and is rendered unfit for both human consumption and the collection of seeds. According to reports, the borer damages okra shoots by 24.6 to 26.0 percent and destroys fruits by 40 to 100 percent [3].

The crop is severely harmed by *E. vittella*, okra jassid, cutworm, white fly, aphids, and other pests. Due to leafhopper assault, there is a drop of 49.8% and 45.1% in height and number of

leaves, respectively [4]. The female moth deposits up to 200–400 eggs alone on the fragile leaves, bracts, and flower buds of okra plants during night. Eggs take 3–4 days to incubate, and the caterpillar goes through 6 stages until reaching maturity in 10–16 days. The moth emerges about 8–14 days in the summer and 18–23 days in the winter after pupating on plants or on the ground among fallen leaves. The lifecycle takes 17 to 29 days to complete. In a year, several overlapping generations are finished [5].

2. METHODOLOGY

The field trails were performed at Central Research Farm (CRF), SHUATS, Naini, Prayagraj. During Kharif season maximum growth is seen, plant will grow upto a height of 5mtrs. The okra variety Panchavati was planted in a randomised block design with a spacing of 45 cm x 30cm. Two sprays were done within a time interval of 15 days. Seven treatments viz., NSKE 10%, *Beauveria bassiana* (1×10^8 CFU/gm) 1.15% WP, Spinosad 45 SC, *Metarhizium anisopliae* (1×10^8 CFU/gm) 4% WSP, Imidacloprid 17.8 SL, Nisco sixer plus, Neem oil with one uncontrolled plot were replicated thrice with a plot size of 2m x 2m. All the agronomic practices were followed like weeding and thinning and two sprays were done. Data was collected based on infestation of shoots and fruits from 5 randomly selected plants per plot.

Table 1. Economics of cultivation

S. No	Treatments	Dose	Yield q/ha	Total cost of yield (₹)	Common cost of cultivation (₹)	Cost Of treatment (₹)	Total cost of cultivation (₹)	C:B ratio
T ₁	NSKE 10%	10gm/ L	130.3	195450	42728	3050	45778	1:4.26
T ₂	<i>Beauveria bassiana</i> 1.15%WP(1×10^8 CFU/gm)	5gm/L	110.8	166200	42728	2760	45488	1:3.65
T ₃	Spinosad 45 SC	0.5ml/ L	175.4	263100	42728	3150	45878	1:5.73
T ₄	<i>Metarhizium anisopliae</i> 4% WSP (1×10^8 CFU/gm)	5gm/L	121.7	182550	42728	1550	44278	1:4.12
T ₅	Imidacloprid 17.8 SL	0.3ml/ L	146.5	219750	42728	3800	46528	1:4.72
T ₆	Nisco sixer plus	1.5ml/ L	156.2	234300	42728	2800	45528	1:5.15
T ₇	Neem oil 5%	5ml/L	134.9	202350	42728	2700	45428	1:4.45
T ₈	Control		74	111000	42728	-	42728	1: 2.59

Cost of okra per quintal is 1500₹ (October 2022)

Yield: The fruits were picked from all plants per plot. The average weight of picked fruits was used to calculate the grain yield. Grain yield was calculated by the following formula

$$\text{Grain yield} = \frac{\text{Grain yield per plot}}{\text{Plot size}} \times 100$$

2.1 Benefit Cost Ratio

Gross return was calculated by multiplying total yield with the market price of the produce. Cost benefit ratio by following formula

$$\text{B: C Ratio} = \frac{\text{Gross returns}}{\text{Total Cost of cultivation}}$$

Where,

B:C = Benefit Cost Ratio

3. RESULTS AND DISCUSSION

The yields among the treatments were significant. The highest yield was recorded in Spinosad 45 SC (175.4 q/ha), followed by Nisco sixer plus (156.2 q/ha), imidacloprid 17.8 SL (146.5 q/ha), Neem oil 5% (134.9 q/ha), NSKE 10% (130.3 q/ha), *Metarhizium anisopliae* 4% WSP (1×10^8 CFU/gm) (121.7 q/ha) and *Beauveria bassiana* 1.15%WP (1×10^8 CFU/gm) (110.8 q/ha) as compared to control plot (74 q/ha). These findings are supported by [6,7,8 and 9].

The yields among the treatments were significant. The highest increased yield over control was recorded in Spinosad 45SC (101.4 q/ha), followed by Nisco sixer plus (82.2 q/ha), Imidacloprid 17.8 SL (72.5 q/ha), Neem oil 5% (60.9 q/ha), *Metarhizium anisopliae* 4% WSP (1×10^8 CFU/gm) (47.7 q/ha) and *Beauveria bassiana* 1.15%WP (1×10^8 CFU/gm) (36.8q/ha).

When cost benefit ratio was worked out, interesting results was achieved. Among the treatments studied, the best and most economical treatment was Spinosad 45SC (1:5.66) followed by Nisco sixer plus (1:4.99), Imidacloprid 17.8SL (1:4.07) and *Beauveria bassiana* 1.15% WP (1×10^8 CFU/gm) (1:3.61) as compared to control (1:2.59). These findings are supported by [10,11,12 and 7].

4. CONCLUSION

It concludes that Spinosad is most effective against okra shoot and fruit borer, followed by nisco sixer plus and imidacloprid. As a result, it is proposed that effective insecticides can be altered in conjunction with current integrated pest management programmes in order to minimise difficulties connected with insecticidal resistance, pest recurrence and so on.

ACKNOWLEDGEMENTS

I am very thankful to Dr. Ashwani Kumar, Associate Professor and Head, Department of Entomology, Sam Higginbottom University of Agriculture Technology And Sciences for providing the required facilities for performing the current work.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Reddy SKB, Patel HP, Bharpoda TM. Utilization of plant extracts for managing fruit borers in Okra, [*Abelmoschus esculentus* (L.) Moench]. International journal of Current Microbiology and Applied Sciences. 2018;7(5):2786-2793.
2. NHB. Area and production of horticulture crops for (2nd Advance Estimates); 2021-22. Available: <https://agricoop.nic.in/en/statistics/state-level/20224>
3. Yadav SK, Kumawat KC, Deshwal HL, Kumar S, Manohar SVS. Bioefficacy of newer and biorational insecticides against shoot and fruit borer, *Earias spp.* On okra. Int. J Curr. Microbiol. App. Sci. 2017; 6(7):1035-1044.
4. Rawat RR, Sahu HR. Estimation of losses in growth and yield of okra due to *Empoaca devastans* Distant and *Earias spp* (India). Indian journal of Entomology. 1973;3(5):252-254.
5. Kaveri G, Kumar A. Comparative efficacy of certain chemicals with biopesticides against diamondback moth *Plutella xylostella* (L.) in cabbage, *Brassica oleracea* (L.). Journal of Entomology and Zoology Studies. 2020;8(6):1350-1353.
6. Patra S, Mondal S, Samantha A, Chatterjee ML. Bioefficacy of some new insecticides against the okra shoot and fruit borer, *Earias vittella* (F.). Pest Management and Economic Zoology. 2007;15(1): 53-56.
7. Sarkar S, Patra S, Samanta A. Evaluation of biopesticides against red cotton bug and fruit borer of okra. The Bioscan. 2015; 10(2):601-604.
8. Pachole SH, Thakur S, Simon S. Comparative bioefficacy of selected chemical insecticides and biorationals

- against shoot and fruit borer, *Earias vittella* Fabricius) on okra [*Abelmoschus esculentus* (L.) Moench]. Journal of pharmacognosy and Phytochemistry. 2017;6(5):1493-1495.
9. Panbude CU, Neharkar PS, Hemant P, Raunt AR. Seasonal incidence and biorational management of fruit and shoot borer (*Earias vittella* (Fab.) on okra. Journal of Pharmacognosy and Phytochemistry. 2019;8:1574-1576.
 10. Gosalwad SS, Kawathekar BR. Efficacy of insecticides against okra fruit borer (*Earias vittella*). Journal of Plant Protection Research. 2009;6(2):59-63.
 11. Singh SK, Singh AK, Singh HM. Relative resistance of okra germplasm to shoot and fruit borer, *Earias vittella* Fabr. under field conditions. Journal of Applied Zoological Researches. 2007;18(2):121–3.
 12. Nalini C, Kumar A. Population dynamics and comparative efficacy of certain chemicals and biopesticides against okra shoot and fruit borer (*Earias vittella*). *International Journal of Life Sciences*. 2016;11(3):1589-1592.

© 2023 Divya et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/101057>