



Influence of Nutrient Management Practices on DTPA Soil Micronutrients and Its Relation with Soil pH and Cation Exchange Capacity in Pearlmillet (*Pennisetum glaucum L.*) Cultivated Soils of Western India

Manish Yadav ^{a*}, N. J. Jadav ^a, Dileep Kumar ^b, Drashti Chaudhary ^a and Suwa Lal Yadav ^a

^a Department of Soil Science and Agricultural Chemistry, Anand Agricultural University, Anand, Gujarat-388110, India.

^b Micronutrient Research Scheme (ICAR), Anand Agricultural University, Anand, Gujarat-388110, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2022/v12i230636

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/83798>

Original Research Article

Received 06 December 2021

Accepted 12 February 2022

Published 13 February 2022

ABSTRACT

A field experiment was conducted during kharif, 2019 at Anand Agricultural University, Anand, Gujarat, India. The effect of nutrient management practices on soil pH, cation exchange capacity, DTPA micronutrient and their inter relationship were studied. The result of present study showed that relationship between available micronutrients and soil pH was significantly and negatively correlated for DPTA Fe, Mn and Zn. In case of DTPA Cu and soil pH relationship was positively correlated. The highest R² value for the relationship between DTPA micronutrients and soil pH was for the available Fe (0.79). The relationship between available micronutrients and cation exchange capacity of soil was reverse of soil pH. Available Fe, Mn and Zn were positively correlated with cation exchange capacity and reverse for DTPA Cu. The lowest R² value for relationship between DTPA micronutrients and soil cation exchange capacity was for DTPA Cu (0.008).

Keywords: Cation exchange capacity; correlation; DTPA micronutrients; soil pH.

1. INTRODUCTION

Nutrient availability from manure has been recognized in crop cultivation for ages. Manure was the major source of nutrients for crop production prior to the advent of chemical fertilizers. There has recently been a resurgence of interest in the utilization of farmyard manure. Concerns about sustaining sustainable agricultural output while safeguarding the environment are said to be driving this interest. Integrated nutrient management is the finest technique for greater resource usage and crop production at a lower cost. In this method, all viable sources of plant nutrients are used depending on cost, and the crop's nutritional balance is supplemented with chemical fertilizers. Organic manures such as animal dung manure (791.6 mt), crop residues (603.5 mt), rural compost (148.3 mt), city compost (12.2 mt), green manure (4.50 m ha), and biofertilizer (0.41 mt) are readily available in India (Bhattacharya and Chakraborty, 2005), and these could be a good substitute for chemical fertilizers in maintaining the soil's physico-chemical and The use of organic manures boosts nutritional content and absorption [1]. Soil is the earth's top layer of loose material, which provides the primary basis for natural flora and other living forms on our planet. Soil fertility is one of the most critical elements that affects crop development and output. Micronutrients are essential for sustaining soil fertility and crop yield. Micronutrients are required in lesser amounts than macronutrients; yet, even if the macronutrients are balanced and high yielding varieties are cultivated, a lack of supply will prevent optimal yields [2]. The acidity and alkalinity of a soil solution are measured by its pH. The hydrogen ion concentration in soil is referred to as pH. The pH scale is not linear, but rather logarithmic. Because of its effect on the availability of vital plant nutrients and the concentration of components poisonous to plants, soil pH can have an impact on plant development [3]. Within a pH range of 4 to 6, the majority of micronutrients such as Cu, Fe, Mn, and Zn are more readily accessible. Organic matter, during its decomposition liberates a number of organic acids, lowers the soil pH and increases the intensity of reduction in soils [4].

Plants get the majority of their nutrients from the soil. It is well understood that optimal plant growth and crop yield are influenced not only by the total amount of nutrients present in the soil at any given time, but also by their availability, which is influenced by physico-chemical properties such as soil texture, organic carbon and calcium carbonate, cation exchange capacity, pH, and electrical conductivity [5]. Gupta (1968) found that soil pH, specific surface area, and clay organic carbon content all affected water soluble boron in soil. Therefore, understanding the chemistry of interrelated soil properties is required for sustainable soil management.

2. MATERIALS AND METHODS

A field experiment was conducted during *kharif* season 2019 at the Collage Agronomy Farm of Anand Agricultural University, Anand, Gujarat, India. The experimental site is located at 22° 35' N and 72° 55' E with an elevation of 45.1 meters above mean sea level. Soils in the experimental field belong to goradu soil series typical of the Inceptisols (Typic Ustochrept). The soil properties and treatment details of the experimental site are given in Table 1 and Table 2, respectively.

The recommended dose of fertilizer was 120-60-0 kg ha⁻¹ N, P₂O₅, K₂O. The application of Bio NP consortia (liquid bio-fertilizer contains *Azotobacter* + PSB) was 5 ml of inoculant in 1 liter of water for 15 minutes as seedling treatment before transplanting of pearl millet and one liter ha⁻¹ at 30 days after transplanting of pearl millet as soil drenching.

2.1 Laboratory Analysis

The collected samples were air-dried and ground to pass through a 2 mm sieve. The samples were stored in polythene lined cotton bags for further analysis in the laboratory for the determination of different properties at ICAR-Micronutrient Research Scheme, Anand Agricultural University, Anand, India. The methods adopted for chemical determinations are listed in the Table 3.

Table 1. Initial soil properties of the experimental site

Soil characteristics	Content
Texture	Loamy sand
pH	8.16
EC	0.23 (dSm ⁻¹)
Cation exchange capacity	11.25 [C mol (P ⁺) kg ⁻¹]
Organic carbon	0.28%
Available nitrogen	188 kg ha ⁻¹
Available phosphorus	53.7 kg ha ⁻¹
Available potassium	537.6 kg ha ⁻¹
DTPA Fe	5.20 (mg kg ⁻¹)
DTPA Mn	6.39 (mg kg ⁻¹)
DTPA Zn	1.20 (mg kg ⁻¹)
DTPA Cu	1.59 (mg kg ⁻¹)

Table 2. Treatment details of experiment

Treatments	Treatment details
T1	100% RDF (control)
T2	100 % RDF + 10 t FYM ha ⁻¹
T3	100 % RDF + 10 t FYM ha ⁻¹ + Bio NP Consortia
T4	100 % RDF + 15 t FYM ha ⁻¹
T5	100 % RDF + 15 t FYM ha ⁻¹ + Bio NP Consortia
T6	75% RDF + 10 t FYM ha ⁻¹
T7	75% RDF + 10 t FYM ha ⁻¹ + Bio NP Consortia
T8	75% RDF + 15 t FYM ha ⁻¹
T9	75% RDF + 15 t FYM ha ⁻¹ + Bio NP Consortia
T10	FYM ha ⁻¹ + 5.0 t Vermicompost ha ⁻¹ + Bio NP Consortia

Table 3. Methods used for laboratory analysis

Parameters	Methods
Soil pH	Potentiometric 1:2 (Jackson, 1973)
CEC	1N NH ₄ OAc pH 7.0 Method (Jackson, 1973)
Micronutrients	
Available Fe	DTPA (Lindsay and Norvell, 1978)
Available Mn	
Available Zn	
Available Cu	

3. RESULT AND DISCUSSION

3.1 Effect of Nutrient Management Practices on DTPA Micronutrients in Soil

The DTPA extractable micronutrient cations (Fe, Mn, Zn, and Cu) in soil as influenced by direct effect of organic manures, inorganic fertilizers along with bio-fertilizers are presented in Fig. 1. The results revealed that addition of FYM, vermicompost and bio-fertilizer in combination of chemical fertilizer or alone significantly affected the DTPA extractable Fe, Mn, Zn and Cu in the soil. It was observed that available Fe (8.58 mg

kg⁻¹), Mn (8.53 mg kg⁻¹) and Zn (1.83 mg kg⁻¹) were found significantly higher due application of 15 t FYM ha⁻¹ + 5.0 t Vermicompost ha⁻¹ + Bio NP Consortia (T10) whereas DTPA extractable Cu noticed significantly higher under T9. The DTPA extractable Fe was higher than the critical limit of 4.5 mg kg⁻¹ in all the treatments. Further, the data indicated that application of 15 t FYM ha⁻¹ + 5.0 t Vermicompost ha⁻¹ + Bio NP Consortia (8.58 mg kg⁻¹) remained at par with T5 (7.90 mg kg⁻¹). In case of DTPA Mn T10 (8.53 mg kg⁻¹) was at par with T3 (8.10 mg kg⁻¹), T4 (8.02 mg kg⁻¹), T5 (8.07 mg kg⁻¹) and T9 (7.98 mg kg⁻¹). DTPA extractable Mn was also higher than the critical limit (2.0 mg kg⁻¹) in all the

treatments. The DTPA-Zn content of soil was significantly higher under T10 (1.83 mg kg^{-1}) over control and it was found at par with T4 (1.54 mg kg^{-1}), T5 (1.79 mg kg^{-1}), T7 (1.77 mg kg^{-1}), T8 (1.76 mg kg^{-1}) and T9 (1.80 mg kg^{-1}). The available Zn status in the soil remained above the critical value of 0.5 mg kg^{-1} in all the treatments. The DTPA-Cu was also significantly influenced due to T9 (1.90 mg kg^{-1}) which was at par with T4 (1.66 mg kg^{-1}), T5 (1.79 mg kg^{-1}), T7 (1.65 mg kg^{-1}) and T8 (1.67 mg kg^{-1}). Many cultivated soils are abundant in Fe, on an average, having a total concentration of 20-40 g kg^{-1} [6]. Fe is frequently present in ferrous (Fe^{2+}) form in primary minerals and few phyllosilicates while its oxidation to the ferric form (Fe^{3+}) showed significant pedogenetic variations [7,8,9] and a sequence of conjugate bases formation where Fe was found to be linked with water and hydroxyls [10,11,12]. Goethite ($\alpha\text{-FeOOH}$) and hematite ($\alpha\text{-Fe}_2\text{O}_3$) are the most abundant minerals among pedogenic forms of crystalline Fe (hydro) oxides in well-drained soil. Occurrence of other Fe oxides could be seen in poorly drained soil as crystalline minerals (magnetite, lepidocrocite and maghemite) or short-range ordered crystalline minerals (ferroxite and ferrihydrite) or non-crystalline precipitates Schwertmann, 1985, [3]. Walia et al. [13] reported that the slight increase in the Cu content ($1.35\text{-}1.66 \text{ mg kg}^{-1}$) was notably observed in plots treated with organic manures over the control plots.

The results indicated that DTPA-micronutrients increased with application of organic matter alone or when it was combined with bio-fertilizers. As vermicompost is rich in micronutrients cations, it might have contributed more to soil. Well decomposed vermicompost and FYM might have involved in formation of organic chelates with organic ligands which have lowered susceptibility to adsorption, fixation and precipitation of micronutrient in the soil and also it was attributed to rapid mineralization of organic manures and consequent release of micronutrients in the soil. Verma et al. [14] reported that higher availability of these micronutrient cations in soil due to application of organic manures and bio-fertilizer was ascribed to the reduction in fixation, increase in mineralization of organic manure and complexing properties of these manures with micronutrients. The release of phenols, aliphatic acids, phenolic acids and stable fraction of

humus such as humic and fulvic acids help in chelation of micronutrients and making it more available to plants and Wadile et al. [15].

3.2 Relationship Between DTPA Micronutrients with Soil pH and CEC

The result regarding correlation was showed that the available Fe ($R^2 = 0.79$) were significant and negative correlation with soil pH (Fig. 2). Among the micronutrients available Fe was most negatively correlated with soil pH. In addition to this, with the increase on soil pH by one unit, available iron decreases by 3.99 times and vice-versa.

The R^2 value of correlation between available Mn, Zn, and Cu were 0.57, 0.49, and 0.16. The affinity of copper with organic matter is fairly high so due this available Cu showed the relatively positive correlation to soil pH because added organic manures of reduced the soil pH of soil. The decreased in soil pH may enhance the availability of above micronutrients, this can be noticed in the experimental results that under T10 micronutrients availability was higher because of added organic matter and subsequently decreased in soil pH. Yadav [2] who suggested that the reduced Fe-availability with increasing pH might be attributed to the conversion of Fe^{2+} to Fe^{3+} ions. The ferric ion (Fe^{3+}) compounds have low solubility in solution and so are less bio-available [16]. The determined result is accordance with the result obtained by Sharma et al. [17]; Mathur et al. [18]; Yadav [19]; Yadav and Meena [20] and Sidhu and Sharma [21]. Correspondingly increasing pH, available manganese decreases gradually and vice-versa. Addition to this, with the increase on soil pH by one unit, available copper increased by 0.45 unit and vice-versa. In addition to this, with the increase on soil pH by one unit, available manganese decreases by 2.66 units and vice-versa.

In contrast to this, the various researcher Sharma et al. [17,22]; Mathur et al. [18,23,24]; Yadav [19,25]; Yadav and Meena [20] and Sidhu and Sharma [21,26] found significant and negative correlation between soil pH and available zinc. The result of correlation study showed the positive relation between DTPA Fe, Mn, Zn and cation exchange capacity of soil whereas available Cu showed the negative relationship with CEC (Fig. 3). The

available Fe ($R^2 = 0.65$) was significant and positive correlation with cation exchange capacity of soil after harvest of pearl millet. The

R^2 values of available Mn and Zn was 0.68 and 0.79, respectively.




Fig. 1. DTPA-micronutrients increased with application of organic matter




Fig. 2. Relationship between pH and DTPA micronutrient under different nutrient management practices after harvest of pearl millet under pearl millet-mungbean system

Table 4. Descriptive statistics of soil properties

	pH	CEC	Fe	Mn	Zn	Cu
Mean	8.35	13.46	6.71	7.46	1.58	1.59
Standard Deviation	0.22	1.27	0.99	0.76	0.27	0.25
Standard Error	0.07	0.40	0.31	0.24	0.08	0.08
Minimum	7.88	11.21	5.32	6.25	1.05	0.98
Maximum	8.62	15.32	8.58	8.53	1.83	1.90




Fig. 3 Relationship between CEC and DTPA micronutrient under different nutrient management practices after harvest of pearl millet under pearl millet-mungbean system

In addition to this, with the increase on cation exchange capacity by one unit, available iron increased by 0.63 times and vice-versa. The available Cu showed negative relationship with CEC, though R^2 value (0.008) was near to zero and indicating almost neutral relation.

4. CONCLUSION

The result of present study showed that relationship between available micronutrients and soil pH was significantly and negatively correlated for DTPA Fe, Mn and Zn. In case of DTPA Cu and soil pH relationship was positively correlated. The highest R^2 value for the relationship between DTPA micronutrients and soil pH was for the available Fe (0.79). The relationship between available micronutrients and cation exchange capacity of soil was reverse of soil pH. Available Fe, Mn and Zn were positively correlated with cation exchange capacity and reverse for DTPA Cu. The lowest R^2 value for relationship between DTPA micronutrients and soil cation exchange capacity was for DTPA Cu (0.008).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Vidyavathi V, Dasog GS, Babalad HB, Hebsur NS, Gali SK, Patil SG, Alagawadi

AR. Nutrient status of soil under different nutrient and crop management practices. *Karnataka Journal of Agricultural Sciences*. 2012;25(2).

2. Yadav BK. Micronutrient status of soils under legume crops in arid region of western Rajasthan, India. *Academic Journal of Plant Sciences*, 2011;4(3):94-97.
3. Brady NC, Weil RR. The nature and properties of soil 13th edition. *Agroforest. Syst.* 2002;54(3):249.
4. Dhaliwal SS, Naresh RK, Mandal A, Singh R, Dhaliwal MK. Dynamics and transformations of micronutrients in agricultural soils as influenced by organic matter build-up: A review. *Environmental and Sustainability Indicators*. 2019;1:100007.
5. Bell RW, Dell B. Micronutrients for sustainable food, feed, fibre and bioenergy production. *International Fertilizer Industry Association (IFA)*; 2008.
6. Cornell RM, Schwertmann U. The iron oxides: Structure, properties, reactions, occurrences, and uses. *Weinheim: Wiley-vch*. 2003;2:71.
7. Stucki JW, Lee K, Zhang L, Larson RA. Effects of iron oxidation states on the surface and structural properties of smectites. *Pure and Applied Chemistry*. 2002;74(11):2145-2158.
8. Adriano DC. Trace elements in terrestrial environments: Biogeochemistry,

9. bioavailability, and risks of metals . New York: Springer. 2001;860.
10. Torrent J, Cabedo A. Sources of iron oxides in reddish brown soil profiles from calcarenites in Southern Spain. *Geoderma*. 1986;37(1):57-66.
11. Stumm W. The dissolution of oxides and aluminum silicates: Examples of surface-coordination-controlled kinetics. *Aquatic surface chemistry*; 1987.
12. Sposito G. *The Chemistry of Soils* Oxford Univ. Press, New York, USA; 1989.
13. Cornell RM, Giovanoli R, Schneider W. Review of the hydrolysis of iron (III) and the crystallization of amorphous iron (III) hydroxide hydrate. *Journal of Chemical Technology & Biotechnology*. 1989;46(2):115-134.
14. Walia SS, Kler DS. Effect of organic and inorganic sources of nutrition on growth, macro and micro nutrient uptake in maize under maize-wheat sequence. *Indian Journal of Ecology*. 2010;37(1):27-29.
15. Verma G, Mathur AK, Bhandari SC, Kanthaliya PC. Long-term effect of integrated nutrient management on properties of a Typic Haplustep under maize-wheat cropping system. *Journal of the Indian Society of Soil Science*. 2010;58(3):299-302.
16. Wadile SC, Girase PP, Sonawane PD, Chitodkar SS. Effect of different fertilizer levels on pearl millet (*Pennisetum glaucum* L.) under rainfed conditions. *Bioinfolet*. 2009;6(4):334-335.
17. Njukeng JN, Ehabe E E, Nkeng GE, Kratz S, Schick J, Schnug E. Investigations on the nutritional status of *Hevea brasiliensis* plantations in the humid forest zone of Cameroon Part 1: Micronutrient status and distribution in soils. *Journal of Cultivated Plants*. 2013;65(10):369-375.
18. Sharma RÁ, Singh M, Sharma JÁ. Correlation studies on DTPA-extractable micronutrients vis-à-vis soil properties in some soils of Nagaur district in semi-arid region of Rajasthan; 2003.
19. Mathur GM, Deo R, Yadav BS Status of zinc in irrigated north-west plain soils of Rajasthan. *Journal of the Indian Society of Soil Science*. 2006;54(3):359-361.
20. Yadav KK. Micronutrient status in soils of Udaipur district of Rajasthan. *Hydrology Journal*. 2008;31:3-4.
21. Yadav RL, Meena MC. Available micronutrients status and their relationship with soil properties of Degana soil series of Rajasthan. *Journal of the Indian Society of Soil Science*. 2009;57(1):90-92.
22. Sidhu GS, Sharma BD. Diethylenetriaminepentaacetic acid-extractable micronutrients status in soil under a rice-wheat system and their relationship with soil properties in different agroclimatic zones of indo-gangetic plains of India. *Communications in Soil Science and Plant Analysis*. 2010;41(1):29-51.
23. Bhattacharya R, Prakash V, Kundu S, Srivastava AK, Gupta HS. Effect of long – term manuring on soil organic carbon, bulk density and water retention characteristics under soybean – wheat cropping sequence in north – western Himalaya. *J. Indian Soc. Soil. Sci.* 2004;52:238 -242.
24. Jackson ML. *Soil Chemical Analysis*, Wisconsin Prentice Hall of India Pvt. Ltd., New Delhi. 1967;46.
25. Lindsay WL, Norvell W. Development of a DTPA soil test for zinc, iron, manganese, and copper. *Soil Science Society of America Journal*. 1978;42(3):421-428.
26. Schwertmann U. The effect of pedogenic environments on iron oxide minerals. In *Advances in Soil Science*. Springer, New York, NY. 1958;171-200.

© 2022 Yadav et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/83798>