



# Oxovanadium (IV) Complexes of $\alpha$ -Amino Acid Schiff Bases and Polypyridyl Ligands: Synthesis, Characterization and Antimicrobial Activity

Muhammad Atique Hasan <sup>a</sup>, Hafizur Rahman <sup>a</sup>,  
Md. Masuqul Haque <sup>a\*</sup> and Md. Nazrul Islam <sup>a\*</sup>

<sup>a</sup> Department of Chemistry, University of Rajshahi, Rajshahi-6206, Bangladesh.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/AJOCS/2024/v14i2290

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc. are available here: <https://www.sdiarticle5.com/review-history/113318>

Original Research Article


Received: 15/12/2023  
Accepted: 20/02/2024  
Published: 05/03/2024

## ABSTRACT

Oxovanadium(IV) complexes of the type [VO(L)(bpy)] (V-1 to V-5) have been synthesized and characterized by FTIR and UV-Vis spectra, molar conductance, melting points, and magnetic susceptibilities measurements, where L= N-salicylidene- $\beta$ -alanine (sal-ala), N-salicylidene-glycine (sal-gly), N-salicylidene-DL- $\beta$ -phenylalanine (sal-pheala), N-salicylidene-leucine (sal-leu), and N-salicylidene-DL-methionine (sal-met), and bpy is 2,2'-bipyridine. The infrared spectral data reveals that the tridentate nature of the amino acid-based Schiff base ligand and the coordination of the ligand through azomethine nitrogen, phenolic oxygen and carboxylate oxygen with vanadyl (VO<sup>2+</sup>) ion. All of these complexes were determined to be non-electrolyte in nature, according to conductivity measurements. The magnetic moment measurements have been attributed that these complexes are paramagnetic and have d<sup>1</sup> configuration of vanadium (IV) ion. In Virto antimicrobial activity of the synthesized complexes was evaluated against two gram-positive (*Bacillus subtilis* and

\*Corresponding authors: E-mail: [mnislam@ru.ac.bd](mailto:mnislam@ru.ac.bd); [masuqul2003@ru.ac.bd](mailto:masuqul2003@ru.ac.bd);

*Staphylococcus aureus*) and two gram-negative (*Escherichia coli*, *Proteus vulgaris*) bacterial strains.



### GRAPHICAL ABSTRACT

**Keywords:**  $\alpha$ -Amino Acid; oxovanadium (IV) complexes; Schiff base; polypyridyl ligands.

## 1. INTRODUCTION

Vanadium coordination chemistry is a new area of interest since it was found to be an important trace element for some organisms [1,2] and a cofactor in some haloperoxidases [3] and nitrogenases [4]. Several studies have been conducted on vanadium complexes with oxidation states (IV) and (V), and several of these compounds have been found to be powerful anti-tumors [5], anti-leukemics [6], and insulin-mimics [7,8].

“Transition metal complexes are now being studied for their potential applications as DNA-dependent electron transfer, DNA structural probes and site-specific nucleic acid cleavage in the development of novel therapeutic and diagnostic agents” [9]. DNA cleavage activity of transition metal complexes has been the subject of investigation [10-12], with the focus on the mid to late transition elements and comparatively little attention paid to the earlier members of the series. Among the latter, vanadium, with its three biologically accessible oxidation states (III, IV, and V) has begun to be recognized as having important biological role [3,13–15]. The bleomycin-vanadyl (IV) complex [16] and  $[\text{VO}(\text{phen})(\text{H}_2\text{O})]^{2+}$ , [17] both have been reported to induce DNA cleavage activity in the presence of  $\text{H}_2\text{O}_2$ . “The dioxovanadium (V) complexes

with 2,2-bipyridine and 1,10-phenanthroline as ancillary ligands have been shown to cleave DNA on photoirradiation” [18–20].

The investigation of oxovanadium (IV) bioinorganic chemistry is motivated by its discernible antibacterial [21-23], anti-leukemia [24], anticancer [25], spermicidal [26], and insulin-mimetic attributes [27,28]. The activity of vanadyl (IV) complexes with triazole ligands is substantial when tested against one or more bacterial or fungal strains [29].

Oxovanadium Schiff base complexes have been recognized for their distinctive bioactive characteristics, notably displaying effectiveness in insulin mimetic functions [30], enzyme inhibition [31], and anti-amoebic activities [32]. These complexes may also have pharmacological [33] and catalytic applications [34], as they exhibit anti-inflammatory [35], antibacterial [36], and anticancer properties [37].

Our interest in developing novel oxovanadium (IV) complexes that potentially exhibit antimicrobial properties and possess unique physicochemical characteristics motivated the current research.

## 2. EXPERIMENTAL METHODS

All chemicals and solvents were reagent grade and were used as received without further

purification. The amino acid-based Schiff base tridentate ligands were synthesized according to published literature. The polypyridyl ligands 2,2'-bipyridine are commercially available. These complexes were synthesized by the template method.

Infrared spectra were recorded on a FTIR-8400, SHIMADZU, Japan using a KBr disc, at the Central Science Lab of Rajshahi University, UV-visible spectra of complexes were recorded on a SHIMADZU DOUBLE BEAM spectrophotometer (model UV-1200) at the Department of Chemistry, Rajshahi University. The melting points or decomposition temperature of all the prepared metal complexes were observed with an electrothermal melting point apparatus. It was, however, not possible to measure the melting points beyond 300°C. The conductance measurements were made at room temperature using a WPACM35 conductivity meter and a dip-cell with a platinized electrode. The SHERWOOD SCIENTIFIC magnetic susceptibility balance was used to probe the magnetic nature of the complexes.

## 2.1 Synthesis and Characterization of Oxovanadium (IV) Complexes of $\alpha$ -Amino Acid Schiff Bases and 2,2'-Bipyridine Ligands

### 2.1.1 Preparation of [VO(sal-ala)(bpy)], (V-1)

The process used for producing oxovanadium (IV) complexes involved mixing  $\alpha$ -amino acids,  $\beta$ -alanine (ala) (0.267 g, 3mmol), and NaOH (0.100 g, 2.25 mmol) in 10 mL methanol with a methanolic solution of salicylaldehyde (sal) (0.3 mL, 3 mmol) in a round-bottom flask. Following an hour of refluxing, a methanolic solution of vanadyl sulphate (0.489 g, 3 mmol) was added to the mixing mixture. After refluxing the mixture for an hour, a light blue precipitate formed. A solution of 2,2'-bipyridine (bpy) (0.469 g, 3 mmol) in 10 mL methanol was added to the mixture [38]. After refluxing the mixture for an additional hour, a dark brown precipitate formed. Subsequently, the precipitate underwent filtration through a Buchner funnel, methanol washing, and subsequent drying in a vacuum desiccator over anhydrous CaCl<sub>2</sub>.

**Yield:** 0.740 g (59%),  $\Lambda_M = 10.0 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$  in DMF at 31°C. IR (KBr phase, cm<sup>-1</sup>): 3471br,

1619vs, 1544m (C=N), 1316m, 959s (V=O), 614m, 456m (br, broad; vs, very strong; s, strong; m, medium; w, weak). UV-Vis (DMSO),  $\lambda/\text{nm}$  ( $\epsilon/\text{M}^{-1} \text{ cm}^{-1}$ ): 268–301 (3436–3462), 364 (2572), 377sh (2312) (sh, shoulder).  $\mu_{\text{eff}} = 1.57$  B.M. at 303 K.

Complexes V-2 to V-5 were prepared by the procedure as described for complex V-1 using DL- $\beta$ -phenylalanine (phyala) (0.495 g, 3 mmol), leucine (leu) (0.393 g, 3 mmol), glycine (gly) (0.225 g, 3 mmol) and DL-methionine (met) (0.447 g, 3 mmol) respectively instead of  $\beta$ -alanine (ala).

### 2.2 Preparation of [VO(sal-phyala)(bpy)], (V-2)

**Yield:** 0.851 g (57%),  $\Lambda_M = 19.3 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$  in DMF at 31°C. IR (KBr phase, cm<sup>-1</sup>): 3435br, 1621vs, 1540s (C=N), 1310m, 942s (V=O), 614m, 445m. UV-Vis (DMSO),  $\lambda/\text{nm}$  ( $\epsilon/\text{M}^{-1} \text{ cm}^{-1}$ ): 267–302 (3675–3524), 364 (2181), 376sh (2161).  $\mu_{\text{eff}} = 1.60$  B.M. at 303 K.

### 2.3 Preparation of [VO(sal-leu)(bpy)], (V-3)


**Yield:** 0.870 g (63%),  $\Lambda_M = 16.1 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$  in DMF at 31°C. IR (KBr phase, cm<sup>-1</sup>): 3453w, 1620w, 1530w, 1310w, 961s (V=O), 618m, 460w. UV-Vis (DMSO),  $\lambda/\text{nm}$  ( $\epsilon/\text{M}^{-1} \text{ cm}^{-1}$ ): 268–301 (3662–3325), 364 (2027), 374sh (2008).  $\mu_{\text{eff}} = 1.55$  B.M. at 303 K.

### 2.4 Preparation of [VO(sal-gly)(bpy)], (V-4)

**Yield:** 0.693 g (57%),  $\Lambda_M = 16.1 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$  in DMF at 31°C. IR (KBr phase, cm<sup>-1</sup>): 3414br, 1654s, 1535s (C=N), 1312m, 963s (V=O), 803s, 617s, 463m. UV-Vis (DMSO),  $\lambda/\text{nm}$  ( $\epsilon/\text{M}^{-1} \text{ cm}^{-1}$ ): 270–303 (3462–3311), 364 (2459), 375sh (2364).  $\mu_{\text{eff}} = 1.54$  B.M. at 303K.

### 2.5 Preparation of [VO(sal-met)(bpy)], (V-5)

**Yield:** 0.862 g (60%),  $\Lambda_M = 7.9 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$  in DMF at 31°C. IR (KBr phase, cm<sup>-1</sup>): 3412br, 1618s, 1535s (C=N), 1312m, 959vs (V=O), 616m, 451m. UV-Vis (DMSO),  $\lambda/\text{nm}$  ( $\epsilon/\text{M}^{-1} \text{ cm}^{-1}$ ): 269–307 (3263–3325), 363 (3175), 388sh (2709).  $\mu_{\text{eff}} = 1.80$  B.M. at 303 K.



**Scheme 1. One pot synthesis of Oxovanadium (IV) Complexes**

### 3. RESULTS AND DISCUSSION

The complexes were prepared in a high yield according to the template method. All the complexes of oxovanadium (IV) are soluble in DMF and DMSO but insoluble in common organic solvents such as methanol, ethanol, benzene, chloroform. The molar conductance of the complexes, as indicated in Table 1, varies between 7.9 and 19.3  $\text{ohm}^{-1} \text{cm}^2 \text{mol}^{-1}$  when measured in DMF at a concentration of  $10^{-3} \text{M}$ . The readings are lower than someone would expect for an electrolyte. As predicted, molar conductance measurements show that the complexes are non-electrolytes in nature [39,40]. "Magnetic moments of oxovanadium (IV) complexes were measured at room temperature and the values are given in Table 1. The magnetic moments of complexes were in the range 1.54–1.80 B.M., which correspond to a single electron of the  $d^1$  system of oxovanadium

(IV) center and paramagnetic in nature" [39,40]. Melting point gives an approximate idea about the nature of the complexes and can suggest whether it is covalent or ionic. The melting point of the complexes prepared for this study is given in Table 1.

#### 3.1 IR Spectral Studies

"The IR spectral data of oxovanadium (IV) complexes show a broad band in the 3413–3472  $\text{cm}^{-1}$  region which is possibly due to the hydrated water molecule in the complexes" [41]. "The complexes exhibit  $\nu(\text{C}=\text{O})$  bands at 1619–1654  $\text{cm}^{-1}$  and  $\nu(\text{C}-\text{O})$  bands at 1310–1316  $\text{cm}^{-1}$  which are significantly lower than the values for respective bands of uncoordinated amino acids. Further, the appearance of  $\nu(\text{V}-\text{O})$  modes at around 618  $\text{cm}^{-1}$  confirms the coordination of carboxylate ion to the central metal ion" [42]. "The  $\nu(\text{O}-\text{H})$  band usually appeared at 3600  $\text{cm}^{-1}$  for

the phenolic –OH is absent in the present complexes which supports the coordination of phenolic oxygen to vanadyl ion. The bands appeared at around 1540  $\text{cm}^{-1}$  may be assigned to  $\nu(\text{C}=\text{N})$  stretching frequency suggesting the coordination of the azomethine nitrogen and heterocyclic nitrogen to the  $\text{VO}^{2+}$  moiety. The coordination of nitrogen of azomethine and heterocyclic nitrogen is further evident by the appearance of  $\nu(\text{V}-\text{N})$  modes at 446–463  $\text{cm}^{-1}$  region” [43]. “The present oxovanadium(IV) complexes exhibit the  $\nu(\text{V}=\text{O})$  stretching frequency in the 942–964  $\text{cm}^{-1}$  region characteristic of metal-oxygen multiple bond, thus ruling out the possibility of polymeric nature of the complexes since the polymeric oxovanadium(IV) complexes exhibit one or more broad absorption bands below 900  $\text{cm}^{-1}$  due to bridging vanadyl group,  $-\text{V}-\text{O}-\text{V}-$ ” [44]. “The present complexes exhibit medium intense band in the region  $\sim 960 \text{ cm}^{-1}$  indicating the monomeric nature of the complexes” [45].

The IR spectra of oxovanadium (IV) complexes (V-1 to V-5) are shown in the Figs. 1–5.

### 3.2 UV-Visible Spectral Studies

The absorption spectra of the complexes were recorded in DMSO in the wavelength of 200–800 nm range. Important UV-Visible spectra of complexes (V-1 to V-5) are tabulated in Table 3.

“Complexes (V-1 to V-5) exhibit a shoulder at  $\sim 375 \text{ nm}$  due to ligand-to-metal charge-transfer (LMCT,  $\text{PhO}^- \rightarrow \text{V}$ ) transition, and the remaining bands appearing in the UV region are assignable to the intra ligand transitions” [46]. “All complexes display bands at 267–307 nm which are assignable to the  $\pi \rightarrow \pi^*$  transition” [47]. The absence of low-intensity bands for the d-d transition at around 500 nm [48] in the current complexes could be the result of not optimizing sample concentration during UV data collection. UV-Visible spectra of complexes (V-1 to V-5) are given in the Figs. 6–10.

**Table 1. Physical Properties of the prepared oxovanadium (IV) complexes**

Complex Symbol	Complex	Color	Melting point/De $^{\circ}\text{C}$	Solubility		Molar conductance $\text{ohm}^{-1} \text{cm}^2 \text{mol}^{-1}$	$\mu \text{ eff}$ in B.M
				DMF	DMSO		
V-1	[VO(sal-ala)(bpy)]	Dark brown	214–217(De)	+ve	+ve	10.0	1.57
V-2	[VO(sal-pheala)(bpy)]	Orange	182–186(De)	+ve	+ve	19.3	1.60
V-3	[VO(sal-leu)(bpy)]	Ash	225–228(De)	+ve	+ve	16.1	1.55
V-4	[VO(sal-gly)(bpy)]	Blackish red	252–255(De)	+ve	+ve	16.1	1.54
V-5	[VO(sal-met) (bpy)]	Brown	212–215(De)	+ve	+ve	7.9	1.80

**Table 2. Important IR frequencies of complexes (V-1 to V-5)**

Complex symbol	Complex	$\nu(\text{OH}) \text{ cm}^{-1}$	$\nu(\text{C}=\text{O}) \text{ cm}^{-1}$	$\nu(\text{C}-\text{O}) \text{ cm}^{-1}$	$\nu(\text{C}=\text{N}) \text{ cm}^{-1}$	$\nu(\text{V}-\text{N}) \text{ cm}^{-1}$	$\nu(\text{V}-\text{O}) \text{ cm}^{-1}$	$\nu(\text{V}=\text{O}) \text{ cm}^{-1}$
V-1	[VO(sal-ala)(bpy)]	3472	1619	1316	1544	457	614	959
V-2	[VO(sal-pheala)(bpy)]	3435	1621	1310	1540	446	619	942
V-3	[VO(sal-leu)(bpy)]	3453	1621	1310	1530	460	618	961
V-4	[VO(sal-gly)(bpy)]	3414	1654	1312	1536	463	618	964
V-5	[VO(sal-met)(bpy)]	3413	1643	1312	1536	452	616	960




Fig. 1. IR spectrum of [VO(sal-ala)(bpy)] complex, V-1




Fig. 2. IR spectrum of [VO(sal-pheala)(bpy)] complex, V-2




Fig. 3. IR spectrum of [VO(sal-leu)(bpy)] complex, V-3




Fig. 4. IR spectrum of [VO(sal-gly)(bpy)] complex, V-4




Fig. 5. IR spectrum of [VO(sal-met)(bpy)] complex, V-5

Table 3. Important UV-Visible spectra of complexes (V-1 to V-5)

Complex symbol	Complex	$\lambda$ , nm ( $\epsilon$ , M <sup>-1</sup> cm <sup>-1</sup> )		
V-1	[VO(sal-ala)(bpy)]	268–301 (3436–3462)	364 (2572)	377sh (2312)
V-2	[VO(sal-pheala)(bpy)]	267–302 (3676–3524)	364 (2181)	376sh (2161)
V-3	[VO(sal-leu)(bpy)]	268–301 (3662–3325)	364 (2027)	374sh (2008)
V-4	[VO(sal-gly)(bpy)]	270–303 (3462–3311)	364 (2459)	375sh (2364)
V-5	[VO(sal-met)(bpy)]	269–307 (3263–3325)	363 (3175)	388sh (2709)




Fig. 6. UV-Visible spectrum of [VO(sal-ala)(bpy)] complex, V-1






Fig. 7. UV-Visible spectrum of [VO(sal-pheala)(bpy)] complex, V-2




Fig. 8. UV-Visible spectrum of [VO(sal-leu)(bpy)] complex, V-3






Fig. 9. UV-Visible spectrum of [VO(sal-gly)(bpy)] complex, V-4



**Fig. 10. UV-Visible spectrum of [VO(sal-met)(bpy)] complex, V-5**

On the basis of the above physical and spectroscopic data, the structure of the complexes may be proposed as distorted octahedral geometry with  $VO_3N_3$  coordination environment. The proposed structure of the oxovanadium (IV) complexes is given below.



Oxovanadium (IV) complexes (V-1 to V-5)

### 3.3 Antimicrobial Screening Result


The antibacterial activities of the five oxovanadium (IV) complexes were screened at the concentration of 10  $\mu\text{g}/\text{disc}$  against four pathogenic bacteria viz. *Escherichia coli*, *Proteus vulgaris*, *Bacillus subtilis* and *Staphylococcus aureus*. The results obtained were compared with the inhibition of the standard antibiotic, streptomycin (10  $\mu\text{g}/\text{disc}$ ). The results are shown in the Table 4. The complexes (V-1 to V-5) were found to be active against all the test bacteria, with the complexes V-2 and V-3 being

even more potent than the standard against all the bacteria except for *Escherichia coli*. The activity of the complexes V-2 against *Escherichia coli* and the activity of the complex V-5 against all pathogens are comparable with the standard.

The antimicrobial activity of the complexes may be described on the basis of their effective interaction with the microbes which cause discrete and distinct types of injuries to microbial cells as a result of oxidative stress, protein dysfunction or membrane damage.

**Table 4. Antibacterial activities of the oxovanadium (IV) complexes and streptomycin**

Bacterial strains		Zone of inhibition, diameter in mm					Streptomycin 10 µg/disc
		V-1 10 µg/disc	V-2 10 µg/disc	V-3 10 µg/disc	V-4 10 µg/disc	V-5 10 µg/disc	
Gram positive	<i>Bacillus subtilis</i>	10	24	25	5	19	19
	<i>Staphylococcus aureus</i>	13	26	33	9	15	19
Gram negative	<i>Escherichia coli</i>	8	12	16	6	12	19
	<i>Proteus vulgaris</i>	12	23	27	6	18	19



**Fig. 11. Graphical representation of antibacterial activity of oxovanadium (IV) complexes**

#### 4. CONCLUSION

The synthesis and characterization of  $VO^{2+}$  complexes of  $\alpha$ -amino acid Schiff bases and 2,2'-bipyridine with O,N,O-donor properties have been conducted. The analytical data indicates that the complexes exhibit non-electrolytic behavior and possess paramagnetic properties. The magnetic moment values of the complexes correspond with the  $d^1$  electronic configuration of the  $V^{IV}O^{2+}$  moiety. The IR spectrum data demonstrates the binding of tridentate amino acid Schiff base ligands to the vanadyl ( $VO^{2+}$ ) ion. Based on physical and spectroscopic evidence, the structure of the complexes can be described as having a distorted octahedral

geometry with a coordination environment of  $VO_3N_3$ . Additional investigation is necessary to consider the V-3 complex as a significant and effective drug in the realm of medicinal chemistry.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- Ran X, Wang L, Cao D, Lin Y, Hao J. Synthesis, characterization and in vitro

- biological activity of cobalt (II), copper (II) and zinc (II) Schiff base complexes derived from salicylaldehyde and D, L-selenomethionine. *Applied Organometallic Chemistry*. 2011;25(1):9-15.
2. Roy M, Saha S, Patra AK, Nethaji M, Chakravarty AR. Ternary iron (III) complex showing photocleavage of DNA in the photodynamic therapy window. *Inorganic Chemistry*. 2007;46(11): 4368-4370.
  3. Bhunia A, Manna S, Mistri S, Paul A, Manne RK, Santra MK, Manna SC. Synthesis, characterization, TDDFT calculation and biological activity of tetradentate ligand based square pyramidal Cu (II) complexes. *RSC Advances*. 2015;5(83):67727-67737.
  4. Obaleye JA, Adediji JF, Adebayo MA. Synthesis and biological activities on metal complexes of 2, 5-diamino-1, 3, 4-thiadiazole derived from semicarbazide hydrochloride. *Molecules*. 2011;16(7): 5861-5874.
  5. Tai X, Yin X, Chen Q, Tan M. Synthesis of some transition metal complexes of a novel Schiff base ligand derived from 2, 2'-bis (p-methoxyphenylamine) and salicylaldehyde. *Molecules*. 2003;8(5): 439-443.
  6. Saladi N, Srivastava AK. Synthesis and crystal structure of di- $\mu$ -methoxy-bis (oxido {N'-[3-(oxidoimino) butan-2-ylidene] benzohydrazonato- $\kappa^3$  O, N, O'} vanadium (V)). *Acta Crystallographica Section E*. 2022;78(12):1213-1216.
  7. Das U, Pattanayak P, Santra MK, Chattopadhyay S. Synthesis of new oxido-vanadium complexes: catalytic properties and cytotoxicity. *Journal of Chemical Research*. 2018;42(1):57-62.
  8. Messerschmidt A, Wever R. X-ray structure of a vanadium-containing enzyme: chloroperoxidase from the fungus *Curvularia inaequalis*. *Proceedings of the National Academy of Sciences*. 1996;93(1):392-396.
  9. Crans DC, Smee JJ, Gaidamauskas E, Yang L. The chemistry and biochemistry of vanadium and the biological activities exerted by vanadium compounds. *Chem. Rev*. 2004;104:849-902
  10. Sasmal PK, Saha S, Majumdar R, De S, Dighe RR, Chakravarty AR. Oxovanadium (IV) complexes of phenanthroline bases: The dipyridophenazine complex as a near-IR photocytotoxic agent. *Dalton Transactions*. 2010;39(8):2147-2158.
  11. Prasad P, Sasmal PK, Majumdar R, Dighe RR, Chakravarty AR. Photocytotoxicity and near-IR light DNA cleavage activity of oxovanadium (IV) Schiff base complexes having phenanthroline bases. *Inorganica Chimica Acta*. 2010;363(12):2743-2751.
  12. Sasmal PK, Saha S, Majumdar R, Dighe RR, Chakravarty AR. Photocytotoxic oxovanadium (IV) complexes showing light-induced DNA and protein cleavage activity. *Inorganic Chemistry*. 2010;49(3):849-859.
  13. Holder AA. *Inorganic pharmaceuticals. Annual Reports Section "A"(Inorganic Chemistry)*. 2011;107:359-378.
  14. Storr T, Thompson KH, Orvig C. Design of targeting ligands in medicinal inorganic chemistry. *Chemical Society Reviews*. 2006;35(6):534-544.
  15. Dash SP, Pasayat S, Dash HR, Das S, Butcher RJ, Dinda R. Oxovanadium (V) complexes incorporating tridentate aroylhydrazonoximes: synthesis, characterizations and antibacterial activity. *Polyhedron*. 2012;31(1):524-529.
  16. Cao Y, Liu H, Yuan Z, Wei G. Two new oxovanadium (iv) compounds containing amino acid schiff base and 1, 10-bathophenanthroline ligands: syntheses, crystal structures, and in vitro evaluation of the anticancer activities. *Australian Journal of Chemistry*. 2017;70(5):608-613.
  17. Sakurai H, Tamura H, Okatani K. Mechanism for a new antitumor vanadium complex: hydroxyl radical-dependent DNA cleavage by 1, 10-phenanthroline vanadyl complex in the presence of hydrogen peroxide. *Biochemical and Biophysical Research Communications*. 1995;206(1):133-137.
  18. Sam M, Hwang JH, Chanfreau G, Abu-Omar MM. Hydroxyl radical is the active species in photochemical DNA strand scission by bis (peroxo) vanadium (V) phenanthroline. *Inorganic Chemistry*. 2004;43(26):8447-8455.
  19. Kwong DW, Chan OY, Wong RN, Musser SM, Vaca L, Chan SI. DNA-photocleavage activities of vanadium(V)-peroxo complexes. *Inorganic Chemistry*. 1997;36(7):1276-1277.
  20. Hiort C, Goodisman J, Dabrowiak JC. Cleavage of DNA by the insulin-mimetic compound,  $\text{NH}_4 [\text{VO}(\text{O}_2)_2(\text{phen})]$ . *Biochemistry*. 1996;35(38):12354-12362.

21. Butler A, Walker JV. Marine haloperoxidases. *Chemical Reviews*. 1993;93(5):1937-1944.
22. Mishra AP, Soni M. Synthesis, structural, and biological studies of some Schiff bases and their metal complexes. *Metal-Based Drugs*; 2008.
23. Muhammad N, Ali S, Shahzadi S, Khan AN. Oxovanadium (IV) complexes of non-steroidal anti-inflammatory drugs: Synthesis, spectroscopy, and antimicrobial activity. *Russian Journal of Coordination Chemistry*. 2008;34:448-453.
24. Dong Y, Narla RK, Sudbeck E, Uckun FM. Synthesis, X-ray structure, and anti-leukemic activity of oxovanadium (IV) complexes. *Journal of Inorganic Biochemistry*. 2000;78(4):321-330.
25. Noblía P, Vieites M, Parajón-Costa BS, Baran EJ, Cerecetto H, Draper P, Gambino D. Vanadium (V) complexes with salicylaldehyde semicarbazone derivatives bearing in vitro anti-tumor activity toward kidney tumor cells (TK-10): crystal structure of [VVO<sub>2</sub> (5-bromosalicylaldehyde semicarbazone)]. *Journal of Inorganic Biochemistry*. 2005;99(2):443-451.
26. D'Cruz OJ, Dong Y, Uckun FM. Spermicidal activity of oxovanadium (IV) complexes of 1, 10-phenanthroline, 2, 2'-bipyridyl, 5'-bromo-2'-hydroxyacetophenone and derivatives in humans. *Biology of Reproduction*. 1999;60(2):435-444.
27. Sakurai H, Kojima Y, Yoshikawa Y, Kawabe K, Yasui H. Antidiabetic vanadium (IV) and zinc (II) complexes. *Coordination Chemistry Reviews*. 2002;226(1-2):187-198.
28. Butler A, Carrano CJ. Coordination chemistry of vanadium in biological systems. *Coordination Chemistry Reviews*. 1991;109(1):61-105.
29. Chohan ZH, Sumrra SH, Youssoufi MH, Hadda TB. Design and synthesis of triazole Schiff bases and their oxovanadium (IV) complexes as antimicrobial agents. *Journal of Coordination Chemistry*. 2010;63(22):3981-3998.
30. Melchior M, Thompson KH, Jong JM, Rettig SJ, Shuter ED, Yuen VG, Orvig C. Vanadium complexes as insulin mimetic agents: coordination chemistry and in vivo studies of oxovanadium (IV) and dioxovanadate (V) complexes formed from naturally occurring chelating oxazolate, thiazolate, or picolate units. *Inorganic Chemistry*. 1999;38(10):2288-2293.
31. Mahroof-Tahir M, Brezina D, Fatima N, Choudhary MI. Synthesis and characterization of mononuclear oxovanadium (IV) complexes and their enzyme inhibition studies with a carbohydrate metabolic enzyme, phosphodiesterase I. *Journal of Inorganic Biochemistry*. 2005;99(2):589-599.
32. Maurya MR, Khurana S, Shailendra Azam A, Zhang W, Rehder D. Synthesis, characterisation and antiamoebic studies of dioxovanadium (v) complexes containing ons donor ligands derived from s-benzylidithiocarbazate. *European Journal of Inorganic Chemistry*. 2003(10);1966-1973.
33. Pawar V, Joshi S, Uma V, Prakash B, Joshi SC. Template synthesis and pharmacological properties of oxovanadium (IV) complexes of macrocyclic Schiff bases. *Int. J. Pharm. Sci. Rev. Res*. 2013;19: 21-26.
34. Maurya MR, Khurana S, Shailendra Azam A, Zhang W, Rehder D. Synthesis, Characterisation and Antiamoebic Studies of Dioxovanadium (v) Complexes Containing ONS Donor Ligands Derived from S-Benzylidithiocarbazate. *European Journal of Inorganic Chemistry*. 2003(10):1966-1973.
35. Shukla S, Mishra AP. Metal complexes used as anti-inflammatory agents: Synthesis, characterization and anti-inflammatory action of VO (II)-complexes. *Arabian Journal of Chemistry*. 2019;12(7): 1715-1721.
36. Mahalakshmi N, Rajavel R. Synthesis, spectroscopic, DNA cleavage and antibacterial activity of binuclear Schiff base complexes. *Arabian Journal of Chemistry*. 2014;7(4):509-517.
37. Abdel-Rahman LH, Abu-Dief AM, Moustafa H, Abdel-Mawgoud AAH. Design and nonlinear optical properties (NLO) using DFT approach of new Cr (III), VO (II), and Ni (II) chelates incorporating tri-dentate imine ligand for DNA interaction, antimicrobial, anticancer activities and molecular docking studies. *Arabian Journal of Chemistry*. 2020;13(1):649-670.
38. Ghosh T, Bandyopadhyay C, Bhattacharya S, Mukherjee G. Synthesis, spectral and electrochemical studies of mixed-ligand oxovanadium (IV) and oxovanadium (V)

- complexes incorporating the tridentate ONO donor schiff base derived from acetylacetone and benzoylhydrazine. Transition metal chemistry. 2004;29(4):444-450.
39. Bitu NA, Hossain S, Kader A, Islam MS, Haque MM, Hossen F, Asraf A. SBDTC Derived Schiff Base Metal Complexation: Synthesis, Physicochemical, and Spectral Studies with Antibacterial Activity. Russian Journal of General Chemistry. 2020;90:1529-1532.
40. Latif MA, Ahmed T, Hossain MS, Chaki BM, Abdou A, Kudrat-E-Zahan M. Synthesis, Spectroscopic Characterization, DFT Calculations, Antibacterial Activity, and Molecular Docking Analysis of Ni (II), Zn (II), Sb (III), and U (VI) Metal Complexes Derived from a Nitrogen-Sulfur Schiff Base. Russian Journal of General Chemistry. 2023;93(2):389-397.
41. Maurya MR, Khurana S, Schulzke C, Rehder D. Dioxo-and oxovanadium (v) complexes of biomimetic hydrazone ono donor ligands: Synthesis, Characterisation, and Reactivity. European Journal of Inorganic Chemistry. 2001;2001(3):779-788.
42. Bitu MNA, Hossain MS, Asraf MA, Hossen MF. Synthesis, characterization and biological activity of oxovanadium (IV) complexes containing  $\alpha$ -amino acid Schiff bases and 5, 6-dimethyl-1, 10-phenanthroline ligands. Croatica Chemica Acta. 2020;93(3):231-236.
43. Heravi MM, Amiri Z, Kafshdarzadeh K, Zadsirjan V. Synthesis of indole derivatives as prevalent moieties present in selected alkaloids. RSC advances. 2021;11(53):33540-33612.
44. Bitu MNA, Hossain MS, Asraf MA, Hossen MF, Kudrat-E-Zahan M. Mixed Ligand Oxovanadium (IV) Complexes of 5,6-Dimethyl-1,10-phenanthroline and Amino Acid Schiff Bases Containing Salicylaldehyde: Synthesis and Characterization with Antibacterial Properties. African Journal of Pharmaceutical Sciences; 2021.
45. Sathyanarayana DN, Patel CC. Ligand series from VO stretching frequency of oxovanadium (IV) complexes. Journal of Inorganic and Nuclear Chemistry. 1968;30(1):207-212.
46. Dutta SK, Kumar SB, Bhattacharyya S, Tiekink ER, Chaudhury M. Intramolecular electron transfer in (BzImH)[(LOV)2O](H2L= S-methyl 3-((2-Hydroxyphenyl)methyl) dithiocarbamate): A novel  $\mu$ -Oxo dinuclear oxovanadium (IV/IV) compound with a trapped-valence (V2O3) 3+ core. Inorganic Chemistry. 1997;36(22):4954-4960.
47. Bermejo-Barrera P, Beceiro-Gonzalez E, Bermejo-Barrera A, Bermejo-Martinez F. Determination of vanadium in water by electrothermal atomisation atomic absorption spectrometry after extraction with 8-hydroxyquinoline in isobutyl methyl ketone. Analyst. 1990;115(5):545-547.
48. Ahmed TT, Alajrawy OI. Oxovanadium (IV) complexes with bidentate ligands synthesis, characterization, and comparison between experimental and theoretical. Materials Today: Proceedings. 2023;80:3823-3836.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/113318>