



Profitability of Annual Chrysanthemum (*Chrysanthemum coronium* L.) Flower Production as Influenced by Application of Mycorrhiza and Vermicompost

Ali Haidar Shah^{a*}, Nomita Laishram^a and Arvinder Singh^a

^a *Division of Vegetable Science and Floriculture, Sher-e-Kashmir University of Agricultural Sciences and Technology of Jammu, Main Campus, Chatha, Jammu - 180009, India.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2022/v12i1030846

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/87439>

Original Research Article

Received 12 March 2022

Accepted 23 May 2022

Published 25 May 2022

ABSTRACT

To check the effect of mycorrhiza on growth and flowering of chrysanthemum, an open field experiment was conducted the effect of vermicompost and mycorrhizal treatments on profitability of *Chrysanthemum coronarium* L. Treatments comprises of 04 mycorrhiza treatments (No application, *Glomus mosseae*, *Acaulospora laevis*, *Gigaspora margarita*) and 04 vermicompost doses (0 g/ m², 500 g/ m², 750 g/ m², 1 kg/ m²). The experiment was laid out in factorial randomized block design with three replications. Chrysanthemum plants showed improved growth and flowering with the application of vermicompost and mycorrhiza. In the present investigations, economic analysis of different treatments reveals that application of treatment M₃V₂ (*Gigaspora margarita* + Vermicompost @ 750 g/m²) resulted in highest benefit cost ratio (5.83) closely followed by benefit cost ratio of the treatments; M₂ V₂ (*Acaulospora laevis* + Vermicompost @ 750 g/m²) and M₁ V₂ (*Glomus mosseae* + Vermicompost @ 750 g/m²).

Keywords: *Chrysanthemum*; profitability; mycorrhiza; vermicompost; gross income; B:C ratio.

1. INTRODUCTION

Annual Chrysanthemum (*Chrysanthemum coronarium* L.) belongs to the family Asteraceae. It is one of the most widely cultivated garden flowers. The flower comes in yellow and white colour and is highly suitable for garland making, pot culture and bedding purposes. The production constraints in annual chrysanthemum cultivation are poor soil fertility, traditional system of crop management etc. The quality of flowers is greatly influenced by the quantity of nutrients and source of nutrients [1,2].

Current development in sustainability involves a rational exploitation of soil microbial activities and the use of less expensive source of plant nutrients which may be made available to the plants by microbiologically mediated process. Arbuscular mycorrhizal fungi have been found to increase plant growth, increase chlorophyll content, phosphorus content, increase resistance to cultural and environmental stresses [3-6].

Addition of organic amendments to soil has been reported to enhance plant biomass, mycorrhizal infectivity and proliferation of AM fungal hyphae in soil. Decomposition of many organic materials by earthworms to vermicompost has been known as a cheaper and environment friendly process. It is a rich source of different essential nutrients which improve overall soil condition and promote yield and growth of plant. Vermicompost contain different types of soil beneficial microbes that can improve plant growth through release of vitamins and hormones [7-9].

The uses of chemical fertilizers cause environmental hazards [10], whereas use of soil microbes and organic medium such as manure and compost has been suggested as a useful method of fertilization and possess a great potential in sustainable agriculture systems [11,12,13,14]. Vermicompost when added to soil loosens the soil and improves the physical and biological properties of the soil including structure of the soil, aeration and water-holding capacity of

the soil [15,16]. Organic manures and biofertilizers improve plant growth, overcome rivalry between vegetative and reproductive stage and increasing the yield potential [17].

Keeping in view the importance of the crop, the present investigation was carried out with the view to increase the profitability of annual chrysanthemum flower production by application of different mycorrhizal strains and vermicompost.

2. MATERIALS AND METHODS

The investigation was carried out in 2019 at the Experimental Farm, Division of Vegetable Science & Floriculture, Sher-e-Kashmir University of Agricultural Sciences and Technology, Main Campus, Chatha, Jammu (J&K).

The experimental field was prepared to a fine tilth and beds of the required dimension were made according to the lay out plan. Healthy seedlings were transplanted on 25/10/2019 in the experimental plots at a spacing of 30 cm x 30 cm thereby accommodating 20 seedlings per bed size of 1.5 m x 1.2 m. At the time of planting, vermicompost at different doses were incorporated into the beds according to the treatment requirements. Vesicular arbuscular mycorrhiza (*Glomus mosseae*, *Acaulospora laevis* and *Gigaspora margarita*) were applied @ 2 g/plant and were incorporated in the planting pits at the time of planting. All other intercultural operations were carried out Irrigations were given as and when required during the crop growth. No disease incidence was recorded during the experiment. In each treatment, five plants were randomly selected and tagged for recording data on flowering parameters. The chlorophyll content of leaf was recorded by using SPAD - 502 chlorophyll meter and expressed in percentage. The partitioning coefficient of root and shoot was measured at peak flowering stage. It was calculated by using the formula.

List 1. Experimental treatments and notations

Experimental treatments and notations					
Factor A (Mycorrhiza treatments)			Factor B (Vermicompost Doses)		
1.	No application	(M ₀)	1.	0 g/ m ²	(V ₀)
2.	<i>Glomus mosseae</i>	(M ₁)	2.	500 g/ m ²	(V ₁)
3.	<i>Acaulospora laevis</i>	(M ₂)	3.	750 g/ m ²	(V ₂)
4.	<i>Gigaspora margarita</i>	(M ₃)	4.	1 kg/ m ²	(V ₃)

The data relating to each parameter were statistically analyzed by applying the technique of analysis of variance using Factorial Randomized Block Design (Gomez and Gomez 1985). The level of significance for f-test and t-test were kept at 5% ($P=0.05$).

3. RESULTS AND DISCUSSION

Results indicated that irrespective of the mycorrhizal species used, inoculated plants produced highest vegetative and floral parameters of commercial importance than non-inoculated plants.

3.1 Flowering and Yield Parameters

Maximum number of flowers per plant (103.58), maximum flower diameter (7.30 cm) and maximum flower duration (49.33 days) was recorded with the application of *Glomus mosseae*. Maximum flower weight (7.84 g), flower yield per plant (767.69 g) and highest shelf life (7.75 days) was recorded with the application of *Gigaspora margarita*. However, the effect of mycorrhiza on days to 50% flowering was found to be non-significant. Asrar and Elhindi [18] also reported that under well-watered conditions, mycorrhizal fungi significantly increased flower diameter and flower weight of marigold plants compared to non-mycorrhiza plants.

The promotion of flowering by mycorrhizal inoculation might be the result of improved plant nutrient concentrations like potassium and a possible hormonal effect by fungal colonization [19] and [20]. Mycorrhizal root systems influence the source to sink balance by utilizing recent photosynthate supplied by photosynthesis in leaves and a considerable proportion of the assimilated carbon [21,22]. The enhanced flowering of plants associated with *G. mosseae* may be the consequence of higher carbohydrate production, especially at the beginning of flower production, and/or more efficient carbohydrate use of these plants during the reproductive phase. This confirms the earlier findings of Dufault et al. [23] that mycorrhizal inoculation in gerbera improves the phosphorus and potassium uptake which results in improved flower quality.

The enhancement of flowering in plants inoculated with *G. mosseae* was independent of foliar nutrient concentration [24] similar theory was also explained by Aboul Nasr [25] in flower production of *Tagetes* and *Zinnia* after their inoculation with *Gigaspora etunicatum*.

Salehi et al. [26] and Shamshiri et al. [27] reported that mycorrhizal symbiosis causes flowering acceleration in cumin and kinnow due to better plant nourishment and uptake of less mobile nutrients viz-a-viz augmenting water uptake [28].

Naeni et al. [29] reported highest number of capitule with the conjoint application of 75% vermicompost and *Glomus mosseae* in milk thistle (*Silybum marianum*). Garmendia and Mangas [24] reported relatively higher flower production of rose plants with the inoculation of *Glomus mosseae* than non-mycorrhizal control. They also found that plants associated with *Glomus mosseae* inoculum exhibited early flowering and increased number of cut flowers of rose. Long et al. [30] also reported increased flower number per plant with application of *Glomus mosseae* in zinnia.

Among the vermicompost doses, earliest 50% flowering (148.83 days) was recorded with the application of vermicompost @ 1kg/m². Maximum number of flowers per plant (102.88), flower weight (7.78 g), flower diameter (7.4 cm), flowering duration (51.92 days), flower yield per plant (799.90 g) and maximum shelf life (7.89 days) was recorded with the treatment of vermicompost @1kg/m².

The earliness of flowering might also be attributed to the supply of macro and micro nutrients, enzymes and growth hormones by vermicompost. These results are in line with the findings of Gayathri et al. [31] in limonium. Advanced flowering due to VAM have also been reported by Gaur et al. [32] in *Petunia hybrida*, *Callistephus chinensis* and *Impatiens balsamina*.

Further, vermicompost continues to decompose after application resulting in an increase in temperature in rhizosphere, which is responsible for acceleration of bloom date. Vermicompost also contains humic acid which is known to increase nutrient accumulation in conditions of limited nutrient availability and when additional nutrients were supplied [33].

More photosynthesis enhanced food accumulation which might have resulted in better plant growth and subsequently higher number of flowers per plant and hence more flower yield [34-36]. Vermicompost, being the source of macro and micro nutrients like; Fe and Zn, enzymes, growth hormones and beneficial microflora might have played a secondary role in increasing the flower yield.

Table 1. Economic profitability of *Chrysanthemum coronarium* L. flower production as influenced by mycorrhiza and vermicompost

Treatments			Flower yield (Kg/ha)	Total expenditure (Rs.)	Gross returns (Rs.)	Net returns (Rs.)	Benefit: Cost ratio
T ₁	M ₀ V ₀	No Mycorrhiza + Vermicompost (0 g/m ²)	23780.50	1,76,500	713415	5,36,915	3.04
T ₂	M ₀ V ₁	No Mycorrhiza + Vermicompost (500 g/m ²)	27246.50	1,80,500	817395	6,36,895	3.53
T ₃	M ₀ V ₂	No Mycorrhiza + Vermicompost (750 g/m ²)	32378.00	1,82,500	971340	7,88,840	4.32
T ₄	M ₀ V ₃	No Mycorrhiza + Vermicompost (1 kg/m ²)	34896.50	2,96,500	1046895	7,50,395	2.53
T ₅	M ₁ V ₀	<i>Glomus mosseae</i> + Vermicompost (0 g/m ²)	31234.50	1,79,000	937035	7,58,035	4.23
T ₆	M ₁ V ₁	<i>Glomus mosseae</i> + Vermicompost (500 g/m ²)	33200.00	1,83,000	996000	8,13,000	4.44
T ₇	M ₁ V ₂	<i>Glomus mosseae</i> + Vermicompost (750 g/m ²)	37700.50	1,85,000	1131015	9,46,015	5.11
T ₈	M ₁ V ₃	<i>Glomus mosseae</i> + Vermicompost (1 kg/m ²)	38243.50	2,99,000	1147305	8,48,305	2.84
T ₉	M ₂ V ₀	<i>Acaulospora laevis</i> + Vermicompost (0 g/m ²)	32983.50	1,79,000	989505	8,10,505	4.53
T ₁₀	M ₂ V ₁	<i>Acaulospora laevis</i> + Vermicompost (500 g/m ²)	35056.50	1,83,000	1051695	8,68,695	4.75
T ₁₁	M ₂ V ₂	<i>Acaulospora laevis</i> + Vermicompost (750 g/m ²)	38938.50	1,85,000	1168155	9,83,155	5.31
T ₁₂	M ₂ V ₃	<i>Acaulospora laevis</i> + Vermicompost (1 kg/m ²)	41722.00	2,99,000	1251660	9,52,660	3.19
T ₁₃	M ₃ V ₀	<i>Gigaspora margarita</i> + Vermicompost (0 g/m ²)	29047.00	1,79,000	871410	6,92,410	3.87
T ₁₄	M ₃ V ₁	<i>Gigaspora margarita</i> + Vermicompost (500 g/m ²)	37231.50	1,83,000	1116945	9,33,945	5.10
T ₁₅	M ₃ V ₂	<i>Gigaspora margarita</i> + Vermicompost (750 g/m ²)	42142.00	1,85,000	1264260	10,79,260	5.83
T ₁₆	M ₃ V ₃	<i>Gigaspora margarita</i> + Vermicompost (1 kg/m ²)	45118.00	2,99,000	1353540	10,54,540	3.53

The higher flower yield due to application of vermicompost has also been reported in marigold [37] and golden rod [38]. Patil et al. [39] confirmed that using organic, inorganic and *in situ* vermiculture in *Jasminum sambac* increased the number of flowers per plant by increasing the leaf area and chlorophyll content.

Interaction effect revealed least number of days for 50% flowering (146.67 days) with the application of M_0V_3 (no Mycorrhiza + vermicompost @ 1kg/m² days). Maximum number of flowers per plant (107.88), flower diameter and flowering duration (57.00 days) was recorded with M_1V_3 (*Glomus mosseae* + vermicompost @ 1 kg/m²). Maximum flower weight (8.64 g), maximum flower yield (902.36 g) and shelf life (8.40 days) was recorded with M_3V_3 (*Gigaspora margarita* + vermicompost @ 1 kg/m²).

Longer shelf life might also be attributed to the better overall food and nutrient status of the flower under these treatments.

3.2 Economic Ratio

The economic value of a crop is determined by its yield and quality. If growing conditions provide required microclimate and nutrition, plants exhibit full expression of genetic potential, yield and quality for long period. The acceptance of any package by farmers depends largely on the comparative economics of a practice and also feasibility of adoption and the effect on yield and quality as well. In the present investigations, economic analysis of different treatments reveals that application of treatment M_3V_2 (*Gigaspora margarita* + Vermicompost @ 750 g/m²) resulted in highest benefit cost ratio (5.83) closely followed by benefit cost ratio of the treatments; M_2V_2 (*Acaulospora laevis* + Vermicompost @ 750 g/m²) and M_1V_2 (*Glomus mosseae* + Vermicompost @ 750 g/m²). This increase in monetary return may be attributed to higher yield and improved quality of flowers which fetches more prices in the market.

4. CONCLUSION

From the results of present investigation, the following conclusions have been drawn which may be beneficial for cultivation of *Chrysanthemum coronarium* L. under the Jammu agro climatic conditions.

Among the various mycorrhizal treatments tested, *Glomus mosseae* (M1) and *Gigaspora margarita* (M3) performed better in terms of better vegetative as well as floral parameters of economic importance. Vermicompost @ 1 kg/ m² recorded highest number of flowers per plant, flower weight, flowering duration and flower yield per plant.

The interaction effect revealed highest number of flowers per plant, flower duration, shelf life with M_1V_3 (*Glomus mosseae* + vermicompost @ 1 kg/ m²). However, highest benefit cost ratio (5.83) was recorded with the treatment M_3V_2 (*Gigaspora margarita* + Vermicompost @ 750 g/m²).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Janowska B, Andrzejak R. Effect of mycorrhizal inoculation on development and flowering of *Tagetes patula* L. 'Yellow Boy' and *Salvia splendens* 'Saluti Red'. Acta Agrobot. 2017;70(2):1703.
2. Kapoor RB, Giri B, Mukherji KG. Improved growth and essential oil yield and quality in *Foeniculum vulgare* Mill. on mycorrhizal inoculation supplemented with P-fertilizer. Biresource Technology. 2004;93:307-311.
3. Anwar M, Patra DD, Chand S, Alpesh K, Naqvi AA, Khjanuja SPS. Effect of organic manures and inorganic fertilizer on growth, herb and oil yield, nutrient accumulation and oil quality of French basil. Communications in Soil Science and Plant Analysis. 2005;36:1737-1746.
4. Asghari MM, Yousefi R, Zavarian AM. Organic fertilizers compost and vermicompost effects on quantitative and qualitative traits of *Lippi citriodora*. Journal of Medicinal Plants. 2016;58:63-71.
5. Gholamhoseini M, Ghalavand A, Dolatabadian A, Jamshidi E, Khodaei Joghian A. Effects of arbuscular mycorrhizal inoculation on growth, yield, nutrient uptake and irrigation water productivity of sunflowers grown under drought stress. Agricultural Water Management. 2013;117:106-114.
6. Hussain S, Sharif M, Khan S, Wahid F, Nihar H, Ahmad W, Khan I, Haider N, Yaseen T. Vermicompost and mycorrhiza

- effect on yield and phosphorus uptake of wheat crop. *Sarhad Journal of Agriculture*. 2016;32(4):372-381.
7. Adhikary. Vermicompost: The story of organic gold- A review. *Agricultural Science*. 2012;3(7):905-917.
 8. Akhzari D, Attaeian B, Arami A, Mahmoodi F, Aslani F. Effects of vermicompost and Arbuscular Mycorrhizal fungi on soil properties and growth of *Medicago polymorpha*. *Compost Science Utilization*. 2015;23:142-153.
 9. Alizadeh O, Alizadeh A. Consideration use of mycorrhiza and vermicompost to optimizing of chemical fertilizer application in corn cultivation. *Advances in Environmental Biology*. 2011;5(6):1279-1284.
 10. Akhzaria D, Kalantari N, Mahdavi Sh. Studying the effects of mycorrhiza and vermicompost fertilizers on the growth and physiological traits of Vetiver Grass (*Chrysopogon zizanioides* L.). *Desert*. 2018;23(1):57-62.
 11. Kumar A, Bhatti SK, Aggarwal A. Field evaluation of vermicompost and selective bio inoculants for the improvement of health status of tomato plants. *Biological Forum – An International Journal*. 2012; 4(2):45-51.
 12. Pezeshkpour P, Ardakani MR, Paknejad F, Vazan S. Effects of vermicompost, mycorrhizal symbiosis and biophosphate solubilizing bacteria on some characteristics related to chickpea root growth under autumn in the dryland condition. *Bulletin of Environment, Pharmacology and Life Sciences*. 2014; 3(2):19-25.
 13. Bhattacharjee P, Chakraborty B, Chakraborty U. Mycorrhiza Modulates Morphology, Color and Duration of Flowers in Hyacinth. *Journal of Biology and Earth Sciences*. 2015;5(1):25-33.
 14. Naeeni FN, Moghadam ARL, Moradi P, Rezaei M, Abdoosi V. Effect of vermicompost and mycorrhiza fungi on yield and growth of milk thistle and antioxidant system activity. *Iranian Journal of Plant Physiology*. 2017;7(3):2063-2074.
 15. Jain MC, Sharma MK, Bhatnagar P, Meena M, Yadav RK. Effect of mycorrhiza and vermicompost on properties of vertisol soil and leaf NPK content of Nagpur Mandarin (*Citrus reticulata* Blanco). *The Asian Journal of Horticulture*. 2012;7(2): 528-532.
 16. Dhayal M, Aravindkshan K. Vermicompost, mycorrhiza and micronutrients mixture improve okra yield. *International Journal of Chemical Studies*. 2018;6(3):1795-1797.
 17. Xie MM, Wu QS. Influence of different potting media on growth and flowering of pot chrysanthemum var. Ajinapurple. *International Journal of Chemical Studies*. 2017;5(4):1667-1666.
 18. Asrar AWA, Elhindi KM. Alleviation of drought stress of marigold (*Tagetes erecta*) plants by using arbuscular mycorrhizal fungi. *Saudi Journal of Biological Science*. 2011;18:93–98.
 19. Perner H, Schwarz D, Bruns C, Maider P, George E. Effect of arbuscular mycorrhizal colonization and two levels of compost supply on nutrient uptake and flowering of *Pelargonium* plants. *Mycorrhiza*. 2007;17: 469-474.
 20. Meir D, Pivonia S, Levita R, Dori I, Ganot L. Application of mycorrhizae to ornamental horticultural crops: Lisianthus (*Eustoma gradiflorum*) as a test case. *Spanish Journal of Agricultural Res*. 2010; 8(1):5-10.
 21. Douds DD, Pfeffer PE, Shachar Hill Y. Carbon partitioning, cost, and metabolism of arbuscular mycorrhizas. In: *Arbuscular mycorrhizas: physiology and function* (Kapulnik Y, Douds DD, eds). Kluwer Academic Publication, Boston, USA. 2000; 107-129.
 22. Smith SE, Read DJ. *Mycorrhizal Symbiosis*. Academic Press, London; 2008.
 23. Dufault RJ, Philips T, Kelly JW. Nitrogen and potassium fertility and plant production influence field production of gerbera. *HortScience*. 1990;25(12):1599-1602.
 24. Garmendia I, Mangas VJ. Application of arbuscular mycorrhizal fungi on the production of cut flower roses under commercial-like conditions. *Spanish Journal of Agricultural Research*. 2012; 10(1):166-174.
 25. Aboul Nasr A. Effect of vesicular arbuscular mycorrhiza on *Tagetes erecta* and *Zinnia elegans*. *Mycorrhiza*. 1996;6: 61-64.
 26. Salehi A, Ganjeh SG, Dehnavi MM, Khajeianr, Gholamhoseini. How vermicompost rates and mycorrhizal treatments affect quantity and quality yield of cumin (*Cuminum cyminum* L.). *Indian Journal of Fundamental and Applied Life Science*. 2015;5(3):127-137.

27. Shamshiri MH, Usha K, Singh B. Growth and nutrient uptake responses of Kinnow to vesicular arbuscular mycorrhizae. International Scholarly Research Network. 2012;7.
28. Ortas I, Kaya Z, Cakmak I. Influence of mycorrhiza inoculation on growth of maize and green pepper plants in phosphorus and zinc deficient soils. In: Plant nutrition-Food security and sustainability of agroecosystems. Horst WJ, et al. eds. Kluwer Academic Publication. 2001;632-633.
29. Naeni NF, Moghadam ARL, Moradi P, Rezaei M, Abdoosi V. Effect of vermicompost and mycorrhiza fungi on yield and growth of milk thistle and antioxidant system activity. Iranian Journal of Plant Physiology. 2017;7(3):2063-2074.
30. Long LK, Yao Q, Huang YH, Yang RH, Gou J, Zhu HH. Effect of Arbuscular Mycorrhizal fungi on Zinnia and the different colonization between *Gigaspora* and *Glomus*. World Journal of Microbiology and Biotechnology. 2010;26:1527-1531.
31. Gayathri HN, Jayaprasad KV, Narayanaswamy P. Response of biofertilizers and their combined application with different levels of inorganic fertilizers in statice (*Limonium caspia*). Journal of Ornamental Horticulture. 2004; 7(1):70-74.
32. Gaur AC. Handbook of Organic Farming and Biofertilizers. Ambica Book Agency, Jaipur, India. 2000;667.
33. David PP, Nelson PV, Sanders DC. A humic acid improves growth of tomato seedling in solution culture. Journal of Plant Nutrition. 1994;17(1):173-184.
34. Khalighi J. The effects of mycorrhizal fungi *Glomus intraradices* on root growth and shoot and total protein content in wheat plants under cadmium toxicity. In: First conference of Agricultural Development Specialist, Astan Hay, North West Iran, Meshkinshar. PNU. 2011;1-11.
35. Tinker DA, Brosnan JT, Herzberg GR. The role of microorganisms in mediating and facilitating the uptake of plant nutrients from soil. Biochemistry and Physiology. 1978;77:115-118.
36. Wu QS, Xia RX. Arbuscular mycorrhiza fungi influence growth, osmotic adjustment and photosynthesis of citrus under well-watered and water stress conditions. Journal of Plant Production. 2006;8:47-55.
37. Mashaldi A. Effect of organic and inorganic fertilizers on growth, yield and post harvest life of marigold (*Tagetes erecta* L.) cv. Double Orange. M.Sc. (Agriculture) Thesis, University of Agricultural Sciences, Bangalore; 2000.
38. Kusuma G. Effect of organic and inorganic fertilizers on growth, yield and quality of golden rod. M.Sc. (Horticulture) Thesis, University of Agricultural Sciences, Bangalore, India; 2001.
39. Patil SR, Reddy BS, Prasanth JM. Effect of organic, inorganic and *in situ* vermiculture on chlorophyll content and flower yield of *Jasminum sambac* Ait. Journal of Ornamental Horticulture. 2004;7(3-4):164-167.

© 2022 Shah et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/87439>