



Nutritional Properties of Fortified Rice (PDS) Distributed at KGBV Hostels of Telangana State, India

Siradala Shiny Pavithra ^{a++*}, T. Sucharita Devi ^{b#},
Aparna Kuna ^{c†} and R. Neela Rani ^{d‡}

^a Post Graduate & Research Centre, Professor Jayashankar Telangana State Agricultural University, Hyderabad, India.

^b Department of Foods and Nutrition, College of Community Science, PJTSAU, Saifabad, India.

^c Department of Foods and Nutrition, MFPI, Quality Control Laboratory, PJTSAU, Rajendranagar, India.

^d ACRIP on WIA, Post Graduate and Research Centre, PJTSAU, Rajendranagar, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ejns/2024/v16i81513>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/121631>

Original Research Article

Received: 08/06/2024
Accepted: 10/08/2024
Published: 12/08/2024

ABSTRACT

Rice is a vital staple crop that serves as the primary source of calories for over half of the global population, particularly in Asia. Despite its importance, rice is inherently deficient in several essential micronutrients, and the nutritional value is further diminished through processing. This

⁺⁺ Research Scholar;

[#] Professor and University Head;

[†] Principal Scientist & Head;

[‡] Principal Scientist (Extension) & Unit Coordinator;

*Corresponding author: Email: shinydavithra20@gmail.com;

Cite as: Pavithra, Siradala Shiny, T. Sucharita Devi, Aparna Kuna, and R. Neela Rani. 2024. "Nutritional Properties of Fortified Rice (PDS) Distributed at KGBV Hostels of Telangana State, India". *European Journal of Nutrition & Food Safety* 16 (8):262-67. <https://doi.org/10.9734/ejns/2024/v16i81513>.

study examines the nutritional composition of fortified rice provided through the Public Distribution System (PDS) in KGBV hostels, emphasizing its potential to address nutritional deficiencies. The analysis focused on key parameters including moisture, ash, fat, crude fiber, protein, carbohydrates, energy, and iron content. Results revealed that fortified rice contains 11.8% moisture, 1.57% ash, 0.78% fat, and 31.05 mg of iron, crude fiber content was 0.3%, protein levels were 7.7%, carbohydrate content was 77.85g, and energy content was 349.22 kcal. These findings highlight the superior nutritional profile of fortified rice, particularly in terms of its ash, energy, and iron content, making it a highly beneficial option for addressing nutritional deficiencies and improving public health outcomes.

Keywords: Rice; micronutrients; fortified rice (PDS); nutritional composition.

1. INTRODUCTION

Rice is an important grain crop, next to wheat in terms of production. Half of the global population and 90% of Asians consider it as a staple food. Rice is a staple meal for about half of the world's population [1] with over 440 million Metric Tons consumed globally each year. Rice makes up at least one-third of daily calorie intake, equivalent to 70 kg per capita per year or 200 g per day in 17 Asian and Pacific countries, 6 Sub-Saharan African countries, and 1 Latin American and Caribbean country [2,3]. It is a significant source of calories for many households. Rice is inherently low in some important micronutrients required for human health [4]. Rice has several macro and micronutrients in its un-milled form. Milling removes the fat and micronutrient-rich bran layers, resulting in the starch-rich white rice that is widely consumed. Rice accounts for 30% of calories on average, but can reach up to 70% in low-income nations [2]. Rice has the capacity to provide micronutrients to a wide population and address micronutrient deficits. Fortifying rice with micronutrients can boost daily consumption of important vitamins and minerals, addressing prevalent nutritional shortages [5] and can effectively treat micronutrient deficits in nations where rice is a primary food. Improved new technologies may create grain-like structures similar to rice kernels with vitamins and minerals in terms of shine, taste, texture, and color [3]. Rice can be fortified by adding micronutrient powders, spraying with vitamins and minerals, or extruding into grain-like structures. These methods can impart iron, folic acid, B vitamins, vitamin A, and zinc. Rice fortification with vitamin A can improve iron and vitamin A levels, while folic acid fortification boosts folate status. A staple food, consumed regularly provides significant calories and nutrients, with preparation varying by location and available foods [6]. In India government provides subsidy rice to the low-income groups through public

distributive system. Earlier the rice was used to provide, to combat malnutrition government started to provide fortified rice because rice-based micronutrient delivery is only effective if fortified rice is affordable for low-income individuals [7].

2. MATERIALS AND METHODS

Fortified rice (PDS) was procured from the KGBV Hostels in Telangana state. The present study was conducted during 2023-2024 at the Department of Foods and Nutrition, Post Graduate Research Centre (PG&RC), Professor Jayashankar Telangana State Agricultural university, Rajendranagar, Hyderabad.

Nutritional properties of rice:

Moisture: To determine the moisture content of sample an empty petri dish with its lid was weighed. A 2.0 g sample of fortified rice was placed in the dish and spread evenly. The sample was dried in a hot air oven at 105°C for 2 hours with the lid open. After drying, the dish and sample were cooled in a desiccator, and the final weight was recorded. The following formula was used to determine the moisture content:

$$\text{Moisture (\%)} = (W2 - W1) (g) / W (g) \times 100. [8]$$

Ash: Fortified rice samples are heated to temperature at 550 – 600°C where the water and other volatile constituents evaporated. The organic constituents were burned in the presence of oxygen to carbon dioxide and oxides of nitrogen that were eliminated together with hydrogen as water. The inorganic residue as ash contains oxides, sulphates, phosphates and chlorides and it constitutes the minerals in food product [8]. The ash content was evaluated by incineration in a muffle furnace. Samples were burnt on a hot plate in a pre-weighed silica crucible. The crucible was then heated to 550°C

in a muffle furnace for four hours. After cooling in the desiccator, the crucible was weighed. The ash content of the sample was determined using the following formula:

$$\text{Ash (\%)} = \frac{\text{Weight of the ash}}{\text{weight of the sample taken}} \times 100.$$

Fat: Fat was estimated as crude ether extract of the dry material using automatic Soxhlet extraction unit [8]. The Soxhlet extraction process was used to determine total fat content. Weighed samples were extracted in a thimble with petroleum ether in a Soxhlet apparatus. The fat extract was then transferred to a previously weighed beaker using a funnel. After evaporating the petroleum ether, the amount of fat was determined by weighing the beaker. The fat content of the sample was determined using the following formula:

$$\text{Fat (\%)} = \frac{\text{weight of the beaker (g)} - \text{weight of the fat (g)}}{\text{weight of the sample}} \times 100.$$

Crude fiber: The crude fiber content of samples was determined by boiling the sample with 1.25% dilute H₂SO₄, washing with water, followed by boiling with 1.25% dilute NaOH and again washing with water. The remaining residue after digestion was taken as crude fiber [8]. To determine crude fiber, 1.0 g powdered sample was placed in fiber bags with a glass spacer, then loaded into a sample carousel in a glass container. The container was filled with 500 ml of 1.25% H₂SO₄ and heated for 30 minutes, followed by boiling in 500 ml of distilled water for 30 minutes. This was succeeded by heating in 500 ml of 1.25% NaOH for another 30 minutes, then boiling again in 500 ml of distilled water for 30 minutes. The residue was transferred to a crucible, weighed (W1), dried at 100°C for 4 hours, cooled in a desiccator, and reweighed (W2). Finally, the crucible was incinerated at 600°C for 3 hours, cooled in a desiccator, and weighed again (W3). The crude fiber content of the sample was determined by using the following formula:

$$\text{Crude fiber (\%)} = \frac{\text{weight of the sample at } 100^{\circ}\text{C} - \text{weight of the sample at } 600^{\circ}\text{C}}{\text{Weight of the sample}} \times 100$$

Protein: Micro kjeldahl method [9] was used to assess the nitrogen content of the sample. Crude protein was computed with a conversion factor 6.25. The nitrogen in protein or any other organic material is converted to ammonium sulphate by Sulphuric acid during digestion. This during steam distillation liberates ammonia which was

collected in boric acid solution and titrated against standard acid. The nitrogen content present in the sample was multiplied by the conversion factor 6.25, expressed as the protein percent. For digestion, 0.5g of fortified rice was mixed with 1g of digestion mixture and 10mL of concentrated sulfuric acid (H₂SO₄) in digestion tubes, then heated until it is clear. A blank was run simultaneously. After cooling, the samples were subjected to distillation in an auto distillation unit with 40% NaOH, 4% boric acid, and distilled water, with 3-4 drops of mixed indicator in the collection flask. The process lasted for 9 minutes, trapping ammonia in the boric acid, which turned green. The device was rinsed with distilled water for 2.5 minutes. For titration, the green boric acid solution was titrated against 0.1N HCl until it turned pink, with a blank executed simultaneously. The titre value determined the percentage of nitrogen (N) in the sample, which was multiplied by 6.25 to obtain find the protein content.

Carbohydrate: The following formula is used to calculate the total carbohydrate content by difference method:

$$\text{Total carbohydrate (g/100g)} = 100 - [\text{Protein(g)} + \text{Fat(g)} + \text{Ash(g)} + \text{Moisture(g)}] [10].$$

Energy: For all the samples, the energy was calculated as follows:

$$\text{Energy (Kcal)} = [\text{Protein(g)} \times 4] + [\text{Carbohydrate(g)} \times 4] + [\text{Fat (g)} \times 9] [11].$$

Mineral Iron (Fe) by AAS: To determine the iron content for fortified rice (PDS) an aliquot of 500 ml from the 1000ppm iron standard was taken in 50 ml volumetric flask and volume was made up to 50 ml with ultrapure water and 10ppm stock solution was obtained. From this stock solution aliquots of 1.0ml, 2.5ml, 5.0ml, 7.5ml, 10.0ml and 25ml were taken into standard 50ml volumetric flask and volume was made up to 50ml with 0.5M nitric acid to get 0.2ppm, 0.5ppm, 1.5ppm, 2.0ppm, and 5.0ppm standard respectively [12].

Statistical analysis: Statistical analysis was represented in mean and median using the MS Excel.

3. RESULTS AND DISCUSSION

The present research was carried out to study the chemical properties of fortified (PDS) rice.

Fortified rice which is prepared by adding micronutrients addresses the malnutrition in humans. Consuming fortified rice reduces anemia in several cases. The nutritional properties of fortified (PDS) rice are presented in below.

The percentage and mean value of moisture, ash, fat, crude fiber, carbohydrate, energy and iron of fortified rice (PDS) compared to Normal rice were presented in the Table 1. The analysis of the nutritional composition of fortified rice revealed several key findings that highlight its potential benefits for dietary supplementation, particularly in addressing nutritional deficiencies.

Moisture: The moisture content of fortified rice was 11.8% and similar findings was reported [13] in normal rice approximately between the range 1.25 to 15.3%. There was significant difference between fortified rice and normal rice. A study reported normal rice at different moisture contents in the range of 10-14% [14]. A Study conducted [15] showed that moisture levels of all normal rice accessions varied between 8.90%–13.57%. Another study shows the moisture content of rice varied between 11.77-14.67% respectively [16]. Moisture content above 14 percent often leads to instability at room temperature, making rice more prone to spoilage and reducing its shelf life [17].

Ash: Ash content of fortified rice (PDS) was 1.57% and in comparison, with normal rice similar findings were stated [18] which is 0.33%. Normal rice varieties had ash content ranging from 0.88 to 1.46% [13]. Nutritional analysis of normal rice had determined an ash content of 1.3% [19]. Significant difference had been observed between fortified rice and normal rice. Study showed ash content of rice was 0.84 to 1.08% respectively [20] and study revealed that ash content varies in between 2.39-6.70% respectively [16]. Ash content of rice was found to be in the range of 1.4-1.9% [21] and 1.13 to 1.70 % [22].

Fat: Fat content of fortified rice, analyzed was 0.78% and that of normal rice reported [18] was 0.24% respectively. Rice varieties had fat content ranging from 0.68 to 1.45% [13]. There was a significant difference between fortified rice and normal rice. Study shows [20] shows crude fat was ranged between 1.32 to 1.73 (%) and also reported fat ranges between 1.23-1.97% in rice

[16] and another study reports crude fat which varies from 1.68 to 2.22 % [22].

Crude fibre: Results had shown that fibre content of fortified rice (PDS) was 0.3% and the fiber content in normal rice was around 0.5% to 1.0% described [15]. Significant difference was not found between normal rice and fortified rice (PDS). A study conducted showed crude fibre content was between 0.58 to 2.06% [20] and 0.00 to 0.03 % [20,22].

Protein: Protein content of fortified rice (PDS) was 7.70% and according to FAO, (1993) rice has protein content that contains 7.2% [23]. There was no significant difference between fortified rice (PDS) and normal rice. Another study reported [20] protein content was between 7.58 to 10.29 (%) and a study revealed that 6.51-9.13% of protein was present in rice [16] and another study reported protein, which varies from 4.79 to 6.11 % [23].

Carbohydrates: Results had shown that fortified rice (PDS) has carbohydrate content of 77.85g/100g and normal rice contains 77-89g which was reported [13] respectively. There was a significant difference in CHO content in both fortified rice and normal rice, the normal rice showed higher carbohydrate content than fortified rice (PDS). Carbohydrates have an essential part in the human body. They serve as an energy source, regulate blood glucose and insulin metabolism, contribute in cholesterol and triglyceride metabolism, and aid in fermentation [24]. Study reported that CHO was ranged between 79.11 to 81.34% (20) and also 62.84-71.06% of CHO is reported in rice (21). According to the study, [22] carbs vary from 69.76 to 73.07%.

Energy: Energy content of fortified rice (PDS) was 349.22 kcal, and study reported [13] normal rice contains 349-373 kcal [13]. There was a significant difference between fortified rice (PDS) which shows more energy value as compared to normal rice.

Iron: Fortified rice which is prepared by adding micronutrients contains about 31.05 mg of iron, whereas, in normal rice the iron content was in between 0.2-2.8mg [13] respectively. Fortified rice Kernels (FRKs) which contain micronutrient iron was (28-42 mg/kg) [25]. There was a significant difference between fortified rice (PDS) and normal rice in which iron content is rich in fortified rice (PDS) i.e., of about 31.05mg.

Table 1. Nutritional composition of Fortified rice (PDS)

S. No	Parameters	Fortified rice (PDS)	Normal rice
1.	Moisture (%)	11.8 ± 0.68	1.25 to 15.3. ¹²
2.	Ash (%)	1.57 ± 0.07	0.33 ¹⁵
3.	Fat (%)	0.78 ± 0.05	0.24 ¹⁵
4.	Crude fiber (%)	0.3 ± 0.05	0.5 to 1.0 ¹⁴
5.	Protein (%)	7.7 ± 0.1	7.2 ¹⁷
6.	Carbohydrate (g/100)	77.85 ± 0.65	77-89 ¹²
7.	Energy (kcal)	349.22 ± 0.55	349-373 ¹²
8.	Iron (mg/kg)	31.05 ± 0.05	0.2-2.8 ¹²

Note: The above values are expressed as mean ± standard deviation of three determinations in fortified rice and normal rice was expressed in percentages

4. CONCLUSION

The nutritional analysis of fortified rice (PDS) demonstrates its superior nutritional profile compared to normal rice, particularly in addressing dietary deficiencies. There was no significant difference in crude fiber, and protein levels in both fortified and normal rice, but exhibits significant difference in moisture, ash, fat, energy, carbohydrate, and iron levels which are essential for overall health. Specifically, the iron content in fortified rice is significantly elevated, making it a potent option for combating iron deficiency anemia. Although fortified rice has lower carbohydrate content, its enhanced nutritional attributes, especially in minerals and energy, highlight its potential as a valuable dietary supplement. Zinc, folic acid and other micronutrients can be recommended for further nutritional analysis in fortified rice in which these nutrients help in combating anemia.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declares that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

ACKNOWLEDGEMENTS

The author wishes to express gratitude to the chairperson, committee members and the Department of Foods and Nutrition, family and friends for their invaluable support during the entire work.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Timmer CP. Reflections on food crises past. *Food Policy*. 2010;35:1–11.
2. Food and Agriculture Organization. FAOSTAT. Rome: FAO. Food Supply Database—Crops Primary Equivalent; 2009. Available: <http://faostat.fao.org/site/609/>
3. Piccoli NB, Grede N, Pee SD, Singhkumarwong A, Roks E, Moench-Pfanner R, Bloem MW. Normal rice fortification: Its potential for improving micronutrient intake and steps required for implementation at scale. *Food and Nutrition Bulletin*. 2012;33(4):361-372.
4. Akinbode A, Matthew F. Improving the quality Attributes of Micronutrient Fortified Extruded Normal rice. AACC International. 2014; 1.
5. Kennedy G, Burlingame B, Nguyen N. Nutrient impact assessment of normal rice in major normal rice-consuming countries. *International Normal rice Commission Newsletter*. 2002;51:33-42.
6. WHO, Fortification of Normal rice. 2023. Fortification of normal rice (who.int)
7. Steiger G, Fischer NM, Cori H, Petit BC. Fortification of normal rice: technologies and nutrients. *Ann. N.Y. Acad. Sci*. 2014;1324:29-39.
8. AOAC. Official methods of Analysis. Association of Official Analysis Chemists International. 2005;18th Edition. Arlington VA, USA.
9. AOAC. Official methods of Analysis. Association of Official Analytical Chemists. 2010;18th Edition. Washington, DC.
10. AOAC. Official methods of Analysis. Association of Official Analytical Chemists. 2006;18th Edition. Washington, DC.

11. AOAC. Official methods of Analysis. Association of Official Analytical Chemists.1980. 13th Edition. Washington, DC.
12. AOAC. Official methods of Analysis. Association of Official Analysis Chemists International. 2012;19th Edition. Gaithersburg, Maryland.
13. Chaudhari PR, Tamrakar N, Singh L, Tandon A, Sharma D. Normal rice nutritional and medicinal properties: A review article. Journal of Pharmacognosy and Phytochemistry. 2018;7(2):150-156.
14. Kibar H, Öztürk T, Esen B. The effect of moisture content on physical and mechanical properties of normal rice (*Oryza sativa* L.). Spanish Journal of Agricultural Research. 2010;8(3):741-749.
15. Verma KD, Srivasthav PP. Proximate Composition, Mineral Content and Fatty Acids Analyses of Aromatic and Non-Aromatic Indian Normal rice; 2017.
16. Rauf R, Sagar A, Rahman T, Hossain MA, Kabir H, Hossain AKMZ. Seed Quality Status of Rice Varieties Based on Physical Properties, Seed Health and Proximate Composition. International Journal of Plant & Soil Science. 2020;32(20):14-24.
17. Mugalavai VK, Aduol KO, Onkware AO. Nutritional Characteristics of Rice (*Oryza sativa* L.) Composite Flours Obtained by Food Fortification. European Journal of Agriculture and Food Sciences. 2021;3(1):79-83.
18. Zubair MA, Rahman SM, Islam SM, Abedin MZ, Sikder AM. A Comparative Study of The Proximate Composition of Selected Normal rice Varieties in Tangail, Bangladesh. J. Environ. Sci. & Natural Resources. 2015;8(2):97-102.
19. Yankah N, Intiful FD, Tette EMA. Comparative study of the nutritional composition of local brown normal rice, maize (obaatanpa), and millet—A baseline research for varietal complementary feeding. Food Sci Nutr. 2020;8:2692–2698.
20. Paul C. Determination of the Proximate Composition and Amylose Content of New Rice for Africa (NERICA) Flour. Turkish Journal of Agricultural Engineering Research. 2020;1(1):131-140.
21. Ibrahim A, Bashir M, Garba, ZM. Proximate and Anti-Nutritional Estimation of some Local and Imported Rice (*Oryza Sativa*): A Comparative Approach. Dutse Journal of Pure and Applied Sciences.2021;7(2b):67-75.
22. Preecharram S, Posoongnoen S, Thummawonsa T, Sripakdee T, Tawil S, Poomsuk N, Jandaruang J. Proximate Composition and Antioxidant Activity of Young Flattened Rice (Khao-Mao). Trends in Sciences. 2023;20(4):6575.
23. FAO. Normal rice in human nutrition; 1993. Available:https://www.fao.org/4/t0567e/T0567E00.htm
24. Holesh JE, Aslam S, Martin A. Physiology, Carbohydrates. In StatPearls [Internet]; 2023.
25. NIN. Formative Research to Understand the Rollout of Fortified rice through Public Distributive System in India. 2024. FFR Consolidated report_Final-01-02-2024.pdf (nin.res.in)

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/121631>