



# Kinetics of Chlorophyll Degradation in Malabar Spinach (*Basellae alba*) and Waterleaf (*Talinum triangulare*) during Hydrothermal Processing

Moses Ayodele Ojo <sup>a\*</sup>, Ajeigbe Olajire <sup>b</sup>,  
A. Olusegun Adeoye <sup>a</sup>, J. Akinkunmi Adejuyitan <sup>a</sup>  
and I. Folaji Odedeji <sup>a</sup>

<sup>a</sup> Department of Food Science, Ladoke Akintola University of Technology, Ogbomoso, Nigeria.

<sup>b</sup> Department of Food Engineering, Ladoke Akintola University of Technology, Ogbomoso, Nigeria.

## Authors' contributions

This work was carried out in collaboration among all authors. Author MAO did conception and design of the study, analysis of the samples, writing of the first draft of the manuscript. Author AO kinetic analysis of data. Author AOA writing and review of the manuscript. Author JAA did writing and review of the manuscript. Author IFO analysis of the samples and statistical analysis. manage the literature searches. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ijbcrr/2024/v33i6911>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/123199>

Original Research Article

Received: 17/07/2024  
Accepted: 19/09/2024  
Published: 07/10/2024

## ABSTRACT

Hydrothermal processing of vegetables often results in a number of changes including degradation of chlorophyll, the green plant pigment that is considered to be of therapeutic benefits. This study was aimed at investigating the kinetic degradation of chlorophyll in Malabar spinach (*Basella alba*)

\*Corresponding author: E-mail: [maojo@lautech.edu.ng](mailto:maojo@lautech.edu.ng);

**Cite as:** Ojo, Moses Ayodele, Ajeigbe Olajire, A. Olusegun Adeoye, J. Akinkunmi Adejuyitan, and I. Folaji Odedeji. 2024. "Kinetics of Chlorophyll Degradation in Malabar Spinach (*Basellae Alba*) and Waterleaf (*Talinum Triangulare*) During Hydrothermal Processing". *International Journal of Biochemistry Research & Review* 33 (6):281-89. <https://doi.org/10.9734/ijbcrr/2024/v33i6911>.

and waterleaf (*Talinum triangulare*) during hydrothermal processing. Freshly harvested leafy vegetables-- Malabar spinach and waterleaf, were obtained and subjected to various hydrothermal conditions (60, 65, 70, 75 and 80°C at 0, 5, 10, 15 and 20 minutes) simulating typical cooking conditions. The chlorophyll of the crushed leaf was extracted using acetone. Changes in chlorophyll was determined using spectrophotometric analysis and the kinetics of degradation for each of the vegetables were determined at varying temperatures and times. Kinetics analysis revealed that the degradation of chlorophyll in Malabar spinach and waterleaf followed the first order reaction. The rate constant obtained from the kinetic analysis provided insights into the reaction rates and allowed for the prediction of chlorophyll degradation during hydrothermal processing. The results showed that the lowest percentage decrease in chlorophyll concentration during the blanching process was obtained at 65°C for 5 minutes for Malabar spinach and for waterleaf at 60°C for 5 minutes. Blanching temperature and time have significant effects on degradation of chlorophyll of Malabar spinach and waterleaf which could consequently determine the economic value and overall degree of acceptability of the green leafy vegetables. The findings provide baseline against which future work can be compared and importance of maintaining quality of the vegetables during processing.

**Keywords:** Malabar spinach; waterleaf; kinetics; chlorophyll degradation; hydrothermal processing.

## 1. INTRODUCTION

Consumption of vegetables in many regions of the world is on the increase because they are often associated with a number of health benefits. Vegetables are highly consumed because they are good sources of vitamins, essential mineral elements and dietary fibre [1,2,3]. Addition of vegetables to daily food intake helps to boost mental health, reduce incidences of cancer and life threatening diseases [3,4,5].

Green plants have the ability to make their own food. They do this through a process called photosynthesis which uses a green pigment called chlorophyll. A pigment is a molecule that has a particular color and can absorb light at different wavelengths. Naturally occurring pigments in plants are many and varied. Chlorophyll is the pigment that gives plants their green colour. Chlorophyll is found in the thylakoids membrane of the chloroplast, a specialized structure naturally occurring in photosynthetic plants. Natural pigments, including chlorophyll, absorb and reflect various wavelengths of light. A light absorbing pigment, chlorophyll is called a photoreceptor. Chlorophyll a absorbs light in the blue-violet region and reflects green light while chlorophyll b absorbs red light and reflects the green light. A chelating agent with a central magnesium (Mg) ion bonded to a larger organic molecule, chlorophyll a and b have the molecular formulae of  $C_{55}H_{72}MgN_4O_5$  and  $C_{55}H_{70}MgN_4O_6$ , respectively. Chlorophyll in plants is useful for many purposes. It is the pigment that enables plants to manufacture their own food during the process of photosynthesis. Chlorophyll enables plants to release oxygen into

the environment during photosynthesis. Amount of chlorophyll in plants can be used to determine nutritional status [6]. The use of chlorophyll from plants as alternative therapies is common in many communities in developing countries; chlorophyll has been proven to have health benefits. Chlorophyll and its derivatives have been shown to increase red blood cells, have antioxidant properties and reduce the effect of deleterious carcinogens [7]. Moreover, the economic value, marketability, and the overall degree of acceptability of many plants particularly leafy vegetables are dependent on the greenness of stems and leaves.

Malabar spinach (*Basellae alba*) and waterleaf (*Talinum triangulare*) are two green leafy vegetables grown in the southern part of Nigeria and some tropical regions of the world. Malabar spinach and waterleaf are locally called *amunututu* and *gbure*, respectively. There are two main species of Malabar spinach, a specie with green stems and a specie with red-purple colour. Although a native of India, Malabar spinach is grown in other tropical regions of the world [7,8]. Waterleaf, a short life-span plant that can adapt to varying climatic conditions, is quite resistant to pests and diseases [8,9,10,11]. The two leafy vegetables are rich in vitamins A, B and C, as well as iron and calcium. They are low in calorie but high in antioxidants and rich sources of fiber [1,12,13,14]. The leaves have mucilaginous texture and are often used as soup thickeners in soups and stews in African and Asian cuisines [13,15,16,17]. Moreover, waterleaf has been used as remediation of soil exposed to heavy metals [18].

Blanching of vegetables inactivates enzymes responsible for senescence and induces structural and chemical changes in the tissue which initiate chlorophyll degradation. Understanding degradation kinetics, including reaction order, rate constant and activation energy, is essential to predict loss of quality during thermal processing and storage [19]. There is dearth of information on kinetic degradation of chlorophyll in Malabar spinach and waterleaf. Hence, the study is aimed at examining the kinetic degradation of chlorophyll in Malabar spinach and waterleaf during hydrothermal processing.

## 2. MATERIALS AND METHODS

### 2.1 Materials

Freshly harvested Malabar spinach – green specie (*Basella alba*) locally called *amunututu* – Fig. 1a and waterleaf (*Talinum triangulare*) locally known as *gbure* – Fig. 1b were obtained from a local farm in Ogbomoso (8°08'N, 4°15'E) and brought to Teaching and Research Unit of Agronomy Department, Ladoke Akintola University of Technology, Ogbomoso for identification.

### 2.2 Sample Preparation and Hydrothermal Processing

The leaves were plucked, washed with clean cold water and then placed on a stainless steel perforated container to allow proper draining of the water from the leaves. The washed vegetable was blanched in a thermo regulated water bath (Model HH-W21-Cr42II, Techmel and Techmel, Texas, USA), at the temperature of 60, 65, 70, 75, and 80 °C and at 0, 5 10, 15 and 20 minutes for each temperature. Each of the processed samples was analysed for its chlorophyll content using the method described below.



Fig. 1a. Malabar spinach (*Basella alba*)

### 2.3 Determination of Chlorophyll Content

Chlorophyll content was determined immediately after blanching by weighing 0.2 g of the crushed leaf and placed in a test tube. 10 cm<sup>3</sup> of 80% acetone was added to the leaf inside the test tube and left for 30 min. About 3 cm<sup>3</sup> of the resulting green extract was poured into a 4 ml cuvette and then placed in a spectrophotometer -- UVV is -NIR V-600, JASCO spectrophotometer. The absorbance of the extract was measured at 645 nm for chlorophyll a and at 663 nm for chlorophyll b. The chlorophyll pigment concentration was calculated using Lichtenthaler's equation [20].

$$Chla = \frac{1395 \times A665 - 688 \times A649}{d \times W \times 1000} \times V \times D \left( \frac{mg}{g} \right) \quad (1)$$

$$Chlb = \frac{24.96 \times A649 - 7.32 \times A665}{d \times W \times 1000} \times V \times D \left( \frac{mg}{g} \right) \quad (2)$$

$$\text{Total } chl = \frac{6263 \times A665 + 18208 \times A649}{d \times W \times 1000} \times V \times D \left( \frac{mg}{g} \right) \quad (3)$$

where: A= absorbance at specified wavelength, V = volume of total extract (ml), W = fresh weight (g), D = coefficient of dilution, d = thickness of cuvette (mm), 1000 = conversion factor μg – mg

### 2.4 Determination of Degradation Kinetics of Chlorophyll

Kinetic model for chlorophyll degradation of Malabar spinach and waterleaf was obtained using a dynamic test approach [21]. An empirical first order kinetic model was used:

$$-\frac{dc}{dt} = kc \quad (4)$$



Fig. 1b. Water leave (*Talinum triangulare*)

where,  $c$  is the concentration of chlorophyll (normalize with respect to initial concentration). The first order rate constant ( $k$ ) was Arrhenius temperature dependent.

$$\ln \frac{c}{c_0} = -kt \quad (5)$$

where  $C$  is the concentration of the chlorophyll  $a$ ,  $b$ , at any time  $t$ ,  $C_0$  is the initial chlorophyll content,  $k$  is the first-order rate constant ( $\text{min}^{-1}$ ),  $t$ : time (min).

## 2.5 Half-Life

Half-life which is the time taken to disintegrate the chlorophyll content present in vegetable leaf to half of its original value was determined by using rate constant ( $k$ ) as

$$\text{Half-life} = 0.693/k \quad (6)$$

## 3. RESULTS AND DISCUSSION

### 3.1 Kinetics of Chlorophyll Degradation in Waterleaf during Hydrothermal Processing

The chlorophyll content of waterleaf (*Talinum triangulare*) during blanching at different temperatures for varying lengths of time is presented in Table 1. Blanching at different temperatures caused significant change ( $P = .05$ ) in the chlorophyll content of waterleaf. Increase in temperature during blanching resulted in decrease in the concentration of chlorophyll. This implied that more chlorophyll was lost by processing at high temperature. Moreover, the loss of chlorophyll during blanching was time dependent. As shown in the table, increase in processing time caused significant reduction ( $P = .05$ ) in the chlorophyll content. The raw sample (control) of waterleaf contained 3.76 mg/l of chlorophyll. Processing by blanching at 60 °C for 5 min did not cause any significant change in the chlorophyll content. Blanching of waterleaf at 60 °C for 10 min caused the chlorophyll content to decrease to 3.46 mg/l and there was further decrease to 3.30 mg/l after blanching for 15 min. The chlorophyll content of waterleaf was reduced to 3.07 mg/l after blanching for 20 minutes at 60 °C. At 65 °C, the chlorophyll content of waterleaf was 3.76, 3.50, 3.36, 3.26 and 3.10 mg/l for blanching time of 0, 5, 10, 15 and 20 min, respectively. There was significant difference ( $P = .05$ ) in the

chlorophyll content of waterleaf when it was processed at 70 °C for varying time of blanching. The chlorophyll content of the blanched samples at 70 °C was 3.76, 3.15, 3.11 3.03 and 3.01 mg/l at the blanching time of 0, 5, 10, 15 and 20 min, respectively. Concentration of chlorophyll in the raw sample decreased from 3.76 to 2.50 mg/l after blanching at 75 °C. The chlorophyll content for samples blanched at 80 °C were 3.76, 3.03, 2.74, 2.38 and 2.15 mg/l at blanching time of 0, 5, 10, 15 and 20 min, respectively. Generally, there was a reduction in the concentration of chlorophyll in the leaf with increase in temperature and processing time. The percentage reduction increased with increase in processing temperature and time. Thus, the highest percentage reduction of 42.82 % was recorded at 80 °C. A similar result was obtained on ascorbic acid and chlorophyll retention of leafy vegetables during processing [22].

### 3.2 Kinetics of Chlorophyll Degradation in Malabar Spinach during Hydrothermal Processing

Table 2 shows the concentration of chlorophyll in the leaf of Malabar spinach (*Basella alba*) during hydrothermal processing at varying lengths of time. Similar to the case of waterleaf, change in temperature during blanching caused significant change ( $P = .05$ ) in the concentration of chlorophyll in the leaf of Malabar spinach. Thus, increase in blanching temperature caused decrease in chlorophyll concentration. Also, increase in blanching time, caused more reduction in the chlorophyll content. The raw sample of *Amunututu* contained 4.15 mg/l of chlorophyll. At 60 °C, the concentrations of chlorophyll were 4.23, 4.21, 4.09 and 4.08 mg/l for blanching time of 5, 10, 15 and 20 min, respectively. At 65 °C, the chlorophyll content ranged from 3.76 mg/l for the raw sample to 3.10 mg/l for the sample blanched for 20 min. Similarly, at temperature of 70 °C, the chlorophyll content were 4.15, 4.00, 4.00, 3.78 and 3.54 at blanching time of 0, 5, 10, 15, and 20 min, respectively. The concentration of chlorophyll ranged from 4.15 mg/l to 3.03 mg/l when the leaf was processed by blanching at 75 °C; the highest percentage reduction of 45.78% was recorded when it was processed at 80 °C. Thus, at varying temperatures and times, there was significant reduction ( $P = .05$ ) in the concentration of chlorophyll. As in the case of *Talinum triangulare*, the chlorophyll content decreased with increase in processing time and temperature.

In general, there was significant difference ( $P = .05$ ) in the chlorophyll content among the samples processed at different temperatures and blanching times. The degradation rates were influenced by both time and temperature. Higher temperatures accelerated chlorophyll degradation, leading to faster color loss. Also, long processing times also contributed to increased chlorophyll degradation. Thus, there was a decrease in chlorophyll content of the samples with increase in blanching time at varying blanching temperatures. Similar finding was reported during blanching of brinjal florets [22]. Chlorophyll is sensitive to heat treatment. The loss of the green color during heat treatment

is mainly attributed to the transformation of chlorophyll -- of green color, to pheophytins -- of brown color, due to the replacement of the magnesium ion in the porphyrin ring of the chlorophyll by two ions of hydrogen, reaction which takes place in acid environment [23,24]. The results obtained in this study are comparable to earlier findings on thermal degradation of chlorophyll of pureed coriander leaves [23] and *amaranthus caudatus* [24]. Similar results were obtained on study of chlorophyll retention of ivy ground leaf [25], thermodynamic study of phenolic degradation and colour of yacon [26], and degradation kinetic of pumpkin [27].

**Table 1. Changes in chlorophyll content (mg/l) of *Talinum triangulare* (waterleaf) during hydrothermal processing at varying temperatures and times**

Time (min)	60 °C	65 °C	70 °C	75 °C	80 °C
0	3.76 ± 0.00 <sup>d</sup> {0.00}	3.76 ± 0.00 <sup>d</sup> {0.00}	3.76 ± 0.00 <sup>d</sup> {0.00}	3.76 ± 0.00 <sup>d</sup> {0.00}	3.76 ± 0.00 <sup>d</sup> {0.00}
5	3.76 ± 0.00 <sup>d</sup> {0.00}	3.50 ± 0.02 <sup>d</sup> {7.42}	3.15 ± 0.08 <sup>b</sup> {16.22}	3.07 ± 0.02 <sup>c</sup> {18.35}	3.03 ± 0.42 <sup>d</sup> {19.41}
10	3.46 ± 0.05 <sup>c</sup> {7.97}	3.36 ± 0.06 <sup>c</sup> {10.63}	3.11 ± 0.00 <sup>ab</sup> {17.28}	3.00 ± 0.00 <sup>c</sup> {20.21}	2.74 ± 0.09 <sup>c</sup> {27.12}
15	3.30 ± 0.03 <sup>b</sup> {12.23}	3.26 ± 0.06 <sup>b</sup> {13.29}	3.03 ± 0.04 <sup>a</sup> {19.41}	2.78 ± 0.11 <sup>b</sup> {26.06}	2.38 ± 0.03 <sup>b</sup> {36.70}
20	3.07 ± 0.10 <sup>a</sup> {18.35}	3.10 ± 0.02 <sup>a</sup> {17.55}	3.01 ± 0.01 <sup>a</sup> {19.95}	2.50 ± 0.11 <sup>a</sup> {33.51}	2.15 ± 0.08 <sup>a</sup> {42.82}

Values above are means of duplicate ± the standard deviation

Means with different letters along the same column are significantly different at ( $P = .05$ )

Values in parenthesis represent percentage decrease in chlorophyll content during processing

**Table 2. Changes in chlorophyll content (mg/l) of Malabar spinach (*Basellae alba*) during hydrothermal processing at varying temperatures and times**

Time (Min)	60 °C	65 °C	70 °C	75 °C	80 °C
0	4.15 ± 0.12 <sup>ab</sup> {0.00}	4.15 ± 0.12 <sup>ab</sup> {0.00}	4.15 ± 0.12 <sup>ab</sup> {0.00}	4.15 ± 0.12 <sup>ab</sup> {0.00}	4.15 ± 0.12 <sup>ab</sup> {0.00}
5	4.15 ± 0.00 <sup>ab</sup> {0.00}	4.15 ± 0.08 <sup>a</sup> {0.00}	4.00 ± 0.00 <sup>c</sup> {3.61}	3.74 ± 0.06 <sup>c</sup> {9.87}	3.43 ± 0.11 <sup>c</sup> {17.34}
10	4.08 ± 0.01 <sup>a</sup> {1.69}	4.03 ± 0.03 <sup>a</sup> {2.89}	4.00 ± 0.00 <sup>c</sup> {3.61}	3.49 ± 0.04 <sup>b</sup> {15.90}	2.74 ± 0.16 <sup>b</sup> {33.97}
15	4.09 ± 0.01 <sup>a</sup> {1.44}	3.96 ± 0.06 <sup>a</sup> {4.57}	3.78 ± 0.35 <sup>b</sup> {8.92}	3.33 ± 0.14 <sup>b</sup> {19.76}	2.56 ± 0.00 <sup>ab</sup> {38.31}
20	4.08 ± 0.02 <sup>a</sup> {1.69}	3.90 ± 0.01 <sup>a</sup> {3.61}	3.54 ± 0.08 <sup>a</sup> {14.69}	3.03 ± 0.08 <sup>a</sup> {26.98}	2.25 ± 0.17 <sup>a</sup> {45.78}

Values above are means of duplicate plus the standard deviation

Different letters along the same column are significantly different at ( $P = .05$ )


Values in parenthesis represent percentage decrease in chlorophyll content during processing

**Table 3. Kinetics of chlorophyll degradation of *Basella alba* at different blanching temperatures and times**

Time (min)	$\text{Ln}\left(\frac{C_t}{C_0}\right)$				
	60 °C	65 °C	70 °C	75 °C	80 °C
5	0.019	0	-0.037	-0.04	-0.190
10	0.0144	-0.029	-0.037	-0.173	-0.415
15	-0.0145	-0.047	-0.101	-0.220	-0.483
20	-0.0170	-0.036	-0.159	-0.315	-0.612

**Table 4. Kinetics of chlorophyll degradation of *Talinum triangulare* at different blanching temperatures and times**

Time (min)	$\text{Ln}\left(\frac{C_t}{C_0}\right)$				
	60 °C	65 °C	70 °C	75 °C	80 °C
5	0	-0.071	-0.177	-0.215	-0.215
10	-0.083	-0.112	-0.180	-0.225	-0.309
15	-0.130	-0.142	-0.215	-0.301	-0.457
20	-0.202	-0.193	-0.222	-0.408	-0.558



**Fig. 2. Plot of  $\text{Ln}\left(\frac{C_t}{C_0}\right)$  against time for (a) *Basella alba* and (b) *Talinum triangulare***

### 3.3 Determination of kinetic degradation Parameter

In order to arrive at the reaction rate constants for the degradation kinetics of Malabar spinach and waterleaf, a first order degradation was presumed. Tables 3 and 4 show degradation kinetics of Malabar spinach and waterleaf, respectively. The plot of  $\text{Ln}\left(\frac{C_t}{C_0}\right)$  against time in which rate constant K was obtained as slope of

the graph. From the graphs (Figs. 2a and 2b), correlation coefficient obtained was greater than 0.9 in all blanching temperature (60, 65, 70, 75 and 80 °C) and time (5, 10, 15 and 20 min) which indicated that the degradation followed the first order kinetics. The values of rate constants for the degradation of *Basella alba* and *Talinum triangulare* are presented in Tables 5 and Table 6, respectively.




Fig. 3. Plot of ln (K) against T<sup>-1</sup>for (a) *Basella alba* and (b) *Talinum triangulare*

Table 5. Rate constant for *Basella alba*

Temperature °C	K	Ln(K)	1/T
60	0.0185	-3.989	3.0E-3
65	0.054	-2.919	2.96E-3
70	0.0115	-4.465	2.92E-3
75	0.005	-5.298	2.87E-3
80	0.0115	-4.465	2.83E-3

Table 6. Rate constant for *Talinum triangulare*

Temperature °C	K	Ln(K)	1/T
60	0.0595	-2.8218	3.0E-3
65	0.0305	-3.490	2.96E-3
70	0.156	-1.858	2.92E-3
75	0.1235	-2.0915	2.87E-3
80	0.0905	-2.4024	2.83E-3

The rate constant increase from 0.005 – 0.054 min<sup>-1</sup> and 0.0180 – 0.1150 min<sup>-1</sup> for Malabar spinach and waterleaf, respectively. It was observed that the rate of chlorophyll degradation of Malabar spinach and waterleaf increased with increase in temperature and blanching time. This finding was in agreement with an earlier finding where the rate constant for kinetic degradation of ODAP in *Lathyrus sativus* increased with increase in temperature [28]. Activation energy  $E_a$  (calm<sup>-1</sup>) was calculated as a product of gas constant, R (1.987 calm<sup>-1</sup>K<sup>-1</sup>) and the slope of the graph obtained by plotting Ln $k$  versus '1/T'. Figs. 3 (a and b). The gradient of these graphs gave the activation energy and is given as 28.996 KJ for *Basella alba* (a) and 10.214 KJ for *Talinum triangulare* (b).

#### 4. CONCLUSION

This study has shown that chlorophyll degradation occurs rapidly during hydrothermal processing leading to the loss of characteristic green color of the leaves. The degradation of chlorophyll in Malabar spinach and waterleaf follows first-order kinetic, with the rate of degradation depending on blanching temperature and time. The highest percentage reduction of 42.82 and 45.78% were recorded for waterleaf and Malabar spinach, respectively after processing at 80 °C for twenty minutes. High temperatures and longer processing times resulted in increased chlorophyll degradation. In view of its nutritional importance, care must always be taken to minimize degradation of chlorophyll during processing. Hence,

understanding the kinetics of chlorophyll degradation in Malabar spinach and waterleaf during hydrothermal processing is expedient for optimizing processing conditions to preserve the colour and other quality characteristics of these leafy vegetables which are considered to be sources of alternative therapies.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Authors hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Agunbiade SO, Ojezele MO, Alao OO. Evaluation of the nutritional, phytochemical compositions and likely medicinal benefits of *Vernonia amygdalina*, *Talinum triangulare* and *Ocimum basilicum* leafy vegetables." *Advances in Biological Research*. 2015;9(3):151–155. Available:<https://doi.org/10.5829/idosi.abr.2015.9.3.93185>
2. Chaudhari VM, Oinam BS, Gouthami NS, Nikhil T, Rashmi S, Shweta S, Umesh T, Badri LN. Unlocking the nutritional power of vegetables: A guide to vibrant health. *European Journal of Nutrition & Food Safety*. 2024;16(8):247-61. Available:<https://doi.org/10.9734/ejnfs/2024/v16i81512>.
3. Vincente AR, Manganaris GA, Ortiz CM, Sozzi GO, Crisosto CH. Nutritional quality of fruits and vegetables. In *Postharvest handling*. 2014; 69-122. Academic Press.
4. Kumar D, Kumar S, Shekhar C. Nutritional components in green leafy vegetables: A review. *Journal of Pharmacognosy and Phytochemistry*. 2020;9(5):2498-2502.
5. Xue S, Yao P, Kang J, Zheng X, Liu C, Gao Y, Zhang D, Zhang X, Hao Y. MdSGR2 Negatively regulates chlorophyll degradation in apple. *Horticulturae*. 2024; 10(5) 439.
6. Amnah MA, Nora M, Alkehayez AHA, Nora AA. Effects of chlorophyll on body functioning and blood glucose levels. *Asian Journal of Clinical Nutrition*. 2017;9:64-70.
7. Schippers RR. Alphabetical treatment of vegetables. In *Plant Resources of Tropical Africa 2—Vegetables*. In: Grubben GJH, Denton O. A, Messiaen CM, Schippers R.R, Lemmens RHMJ, Oyen LPA. PROTA Foundation: 668. Wageningen, Netherlands; 2004.
8. Aja PM, Okaka ANC, Onu PN, Ibiam U, Urako AJ. Phytochemical composition of *Talinum triangulare* (waterleaf) leaves. *Pakistan Journal of Nutrition*. 2010; 9(6):527–530. Available:<https://doi.org/10.3923/pjn.2010.527.530>
9. Nya EJ, Eka MJ. Morphological characterization and hybridization of *Talinum triangulare* land races for desirable metric characters in South Eastern Nigeria. *The International Journal of Science and Technoledge*. 2015;3(7):192–197.
10. Liu GD, Simonne, EH, Morgan KT, Hochmuth GJ, Agehara S., Mylavarapu R, Williams PB. Chapter 2. Fertilizer Management for Vegetable Production in Florida." *EDIS 2021 (VPH)*. Available:<https://doi.org/10.32473/edis-cv296-2021>
11. Eleazu CO, Eleazu KC. Bioactive constituents and *In vitro* antioxidant capacity of waterleaf (*Talinum triangulare*) as affected by domestic cooking. *European Journal of Medicinal Plants*. 2013; 3(4):540–551. Available:<https://doi.org/10.9734/EJMP/2013/4577>
12. Swarna J, Lokeswari TS, Smita M, Ravindhran R. Characterisation and determination of *in vitro* antioxidant potential of betalains from *Talinum triangulare* (Jacq) Willd. *Food Chemistry*. 2013;141(4):4382–90. Available:<https://doi.org/10.1016/j.foodchem.2013.06.108>
13. Kristine BS, Chien-Chang B, Consolacion R. Chemical Constituents of *Talinum triangulare*. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*. 2015;6(1):167–171.
14. De Oliveira Amorim AP, de Carvalho AR, Lopes NP, Castro RN, de Oliveira MCC, de Carvalho MG. Chemical compounds



- isolated from *Talinum triangulare*. Food Chemistry. 2014;160:204–208. Available: <https://doi.org/10.1016/j.foodchem.2014.01.114>
15. Willie ES, Eze, CC. Effect of removal of inflorescence on fresh vegetable yield in waterleaf (*Talinum triangulare*). Journal of Agriculture and Veterinary Science. 2016;9(7):1–4. Available: <https://doi.org/10.9790/2380-0907010104>
  16. Iren OB, Ayito EO, Okon-inyang PO, Ofem KI, Soil Nutrient Status and Yield of Waterleaf (*Talinum triangulare Jacq*) as Influenced by Rates of Organomineral Fertilizer in a Rainforest Utisol, Nigeria. International Journal of Applied Research. 2017;3(5):581–585.
  17. Anyalogbu EAA, Anadi CC, Nwweje-Anyalowu PC, Nnoli, MC. Use of (*Talinum triangulare*) in remediation of soil exposed to heavy metals: a green technology approach. World Journal of Pharmaceutical and Life Sciences. 2017;3(9):48–53.
  18. Canjura FL, Schwartz SJ, Nuness RV. Degradation kinetics of chlorophylls and chlorophyllides. Journal of Food Science. 1991;56(6):1639-1643.
  19. Lichtenthaler HK. Chlorophylls and carotenoids: pigments of photosynthetic biomembranes. Methods in Enzymology. 1987;(148):350-382.
  20. Penicaud C, Achir N, Dhuique-Mayer C, Dornier M, Bohuon P. Degradation of  $\beta$ -carotene during fruit and vegetable processing or storage: reaction mechanisms and kinetic aspects: a review. Fruits. (2011); 66:(6)417-440. DOI:10.1051/fruits/2011058.
  21. Negi PS, Roy SK. Effect of blanching and drying methods on  $\beta$ -carotene, ascorbic acid and chlorophyll retention of leafy vegetables. LWT-Food Science and Technology. 2000;33(4):295-298.
  22. Sanjuán N, Clemente G, Bon J, Mulet A. The effect of blanching on the quality of dehydrated broccoli florets. European Food Research and Technology. 2001;213: 474-479.
  23. Rudra SG, Singh H, Basu S, Shivhare US. Enthalpy entropy compensation during thermal degradation of chlorophyll in mint and coriander puree. Journal of Food Engineering. 2008;86(3): 379-387.
  24. Nayak PK, Chandrasekar CM, Haque A, Kesavan RK. Influence of pretreatments on the degradation kinetics of chlorophylls in morisa xak (*Amaranthus caudatus*) leaves after microwave drying. Journal of Food Process Engineering. 2021;44(9): e13790.
  25. Potisate Y, Phoungchandang S. Chlorophyll retention and drying characteristics of ivy gourd leaf (*Coccinia grandis Voigt*) using tray and heat pump-assisted dehumidified air drying. Drying Technology. 2010;28(6): 786-797.
  26. Lago CC, Noreña CPZ. Thermodynamic and kinetics study of phenolics degradation and color of yacon (*Smallanthus sonchifolius*) microparticles under accelerated storage conditions. Journal of Food Science and Technology. 2017;54:4197-4204.
  27. Song J, Wang X, Li D, Liu C. Degradation kinetics of carotenoids and visual colour in pumpkin (*Cucurbita maxima L.*) slices during microwave-vacuum drying. International Journal of Food Properties. 2017;(20): S632-S643.
  28. Tarade KM, Singhal RS, Jayram RV, Pandit AB. Kinetics of degradation of ODAP in *Lathyrus sativus L.* flour during food processing. Food Chemistry. 2006;(104):643–649.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/123199>