



Potential Edible Film Biocomposite (Canna Starch, Nanochitosan and Fish Oil) for Extend Shelf Life Pangasius Sausages Based on Organoleptic Characteristics, Water Content, and pH: A Review

Al Marisa Rinaldi ^{a*}, Emma Rochima ^a,
Mochamad Untung ^a and Iis Rostini ^a

^a Department of Fisheries, Faculty of Fisheries and Marine Sciences, Padjadjaran University, Indonesia.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJFAR/2023/v22i1561

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/97857>

Review Article

Received: 25/01/2023

Accepted: 27/03/2023

Published: 05/04/2023

ABSTRACT

In general, commercial sausages can only last at room temperature for 1 day, because sausages are made from raw materials that are easily degraded in quality, especially fish sausages where fish are included in perishable foods that are easily degraded due to the high water content in fish which is suitable for microbial growth so a solution is needed to overcome these problems. Biocompote film made from fish oil, nano chitosan and canna starch has the potential to be used

*Corresponding author: Email: marisa18002@mail.unpad.ac.id;

as a packaging substitute for casings that are more environmentally friendly and can extend the shelf life by analyzing pH, water content and organoleptic to determine the effect of the application of edible film on catfish sausages.

Keywords: Sausage; catfish; edible film; fish oil; nanochitosan; canna starch.

1. INTRODUCTION

Sausages are generally made from beef or chicken, there are few or even rare sausages made from fish meat, in fact fish meat has fiber that is more easily digested and contains good fatty acids. Utami et al. [1]. One of the food packaging materials that has a plastic-like shape, is made from biopolymer material, and is safe for human consumption, namely edible film. Apart from being safe for human consumption, edible film can be a good alternative to synthetic plastics for food packaging applications because it has biodegradable properties, can increase the shelf life of food, and is safe for health. Balti et al. [2].

Edible film is a thin edible sheet made of biopolymer and functions as a food product packaging. In general, there are 3 classes of biopolymers that are often used, namely hydrocolloids, lipids, and composites Santoso et al. [3]. Putri et al. [4] Chitosan is a natural polysaccharide that is non-toxic and easily biodegradable. Chitosan has a shape similar to cellulose and the difference lies in the second C chain group. The ability to suppress bacterial growth is due to chitosan having a positively charged polycation which is able to inhibit the growth of bacteria and mold. Mekawati et al. [5] Starch is often used in the food industry as a biodegradable film to replace plastic polymers because it is economical, renewable, and provides good physical characteristics. Kusumawati and Putri et al. [6]. The high amylose content is an advantage of canna starch, because the amylose content has the ability to form gels and is suitable for producing the product desired by Sariyanti and Utami [7]. One form of utilization of fishery industry waste that can be utilized is processing, namely fish oil. It is known that the heads and bones of fish contain fat which is rich in vitamin A and omega-3 fatty acids which are good for health. Fish oil also contains essential fatty acids that the body cannot produce naturally. There are two types of omega-3 fatty acids found in fish oil, namely docosahexaenoic acid (DHA) and eicosapentaenoic acid (EPA). Fish oil also contains high amounts of vitamins A and D, two

types of fat-soluble vitamins Apituley et al. [8] because the amylose content has the ability to form gels and is suitable for producing the desired product by Sariyanti and Utami [7]. One form of utilization of fishery industry waste that can be utilized is processing, namely fish oil. It is known that the heads and bones of fish contain fat which is rich in vitamin A and omega-3 fatty acids which are good for health. Fish oil also contains essential fatty acids that the body cannot produce naturally. There are two types of omega-3 fatty acids found in fish oil, namely docosahexaenoic acid (DHA) and eicosapentaenoic acid (EPA). Fish oil also contains high amounts of vitamins A and D, two types of fat-soluble vitamins Apituley et al. [8] because the amylose content has the ability to form gels and is suitable for producing the desired product by Sariyanti and Utami [7]. One form of utilization of fishery industry waste that can be utilized is processing, namely fish oil. It is known that the heads and bones of fish contain fat which is rich in vitamin A and omega-3 fatty acids which are good for health. Fish oil also contains essential fatty acids that the body cannot produce naturally. There are two types of omega-3 fatty acids found in fish oil, namely docosahexaenoic acid (DHA) and eicosapentaenoic acid (EPA). Fish oil also contains high amounts of vitamins A and D, two types of fat-soluble vitamins Apituley et al. [8] One form of utilization of fishery industry waste that can be utilized is processing, namely fish oil. It is known that the heads and bones of fish contain fat which is rich in vitamin A and omega-3 fatty acids which are good for health. Fish oil also contains essential fatty acids that the body cannot produce naturally. There are two types of omega-3 fatty acids found in fish oil, namely docosahexaenoic acid (DHA) and eicosapentaenoic acid (EPA). Fish oil also contains high amounts of vitamins A and D, two types of fat-soluble vitamins Apituley et al. [8] One form of utilization of fishery industry waste that can be utilized is processing, namely fish oil. It is known that the heads and bones of fish contain fat which is rich in vitamin A and omega-3 fatty acids which are good for health. Fish oil also contains essential fatty acids that the body cannot produce naturally. There are two types of

omega-3 fatty acids found in fish oil, namely docosahexaenoic acid (DHA) and eicosapentaenoic acid (EPA). Fish oil also contains high amounts of vitamins A and D, two types of fat-soluble vitamins Apituley et al. [8] Fish oil also contains essential fatty acids that the body cannot produce naturally. There are two types of omega-3 fatty acids found in fish oil, namely docosahexaenoic acid (DHA) and eicosapentaenoic acid (EPA). Fish oil also contains high amounts of vitamins A and D, two types of fat-soluble vitamins Apituley et al. [8] Fish oil also contains essential fatty acids that the body cannot produce naturally. There are two types of omega-3 fatty acids found in fish oil, namely docosahexaenoic acid (DHA) and eicosapentaenoic acid (EPA). Fish oil also contains high amounts of vitamins A and D, two types of fat-soluble vitamins (Apituley et al. [8]. According to research Leger et al. [9]. EPA has potential as an antioxidant, and cardioprotective. The content of EPA and DHA also has benefits as anti-inflammatory and anti-arrhythmic substances which are beneficial for heart function Endo et al. [10].

2. BIOCOSMPOSITE EDIBLE FILM

One of the food packaging materials that has a plastic-like shape, is made from biopolymer material, and is safe for human consumption, namely edible film. Apart from being safe for human consumption, edible film can be a good alternative to synthetic plastics for food packaging applications because it has biodegradable properties, can increase the shelf life of food, and is safe for health Balti et al. [2] in Suryono et al. [11]. Quoting from Utami et al. [1] Based on its function, Edible packaging is divided into two groups, namely as a coating (edible coating) and in the form of sheets (edible film). One food packaging that has a plastic-like shape, is made from biopolymer material, and is


safe for human consumption, namely edible film Balti et al. [2] Based on the components, edible films are divided into three categories, namely, hydrocolloids (containing polysaccharides, proteins, or alginates), lipids or fats (fatty acids, acylglycerols or waxes), and composites (made of hydrocolloids and lipids). Rosida et al. [12] Fats (lipids), carbohydrates (polysaccharides), and proteins (polypeptides) are components of the raw material for making edible films which have thermoplastic properties, making them easy to print as edible films. This polymer has the advantage of being biodegradable and renewable Widodo et al. [13]. According to Nahwi 2016 in Febianti et al. [14] Food products such as fresh vegetables, fruits and even sausages can be packaged using edible film which functions as a barrier to the diffusion of oxygen gas, water vapor, carbon dioxide and flavor components because it creates internal conditions that match the needs of the packaged product, so it can inhibit decline in quality Mustapa et al. [15].

3. CANNA STARCH


According to Dewi et al [16], starch is a type of polysaccharide that can be found in nature in abundant quantities so it is easy to find, starch also has biodegradable properties, and is economical in price. Another advantage possessed by starch is that it has properties that are suitable for making edible films or edible coatings because it can form a film structure that is strong enough Widodo et al. [13]. Starch is included in the hydrocolloid group which is commonly used as a basic material for making edible films. There are several advantages of edible films made from hydrocolloids, some of which include having the desired mechanical properties, as well as having the ability to protect food products from oxygen, lipids and carbon dioxide Nofiandi et al. [19].

Table 1. Ingredients for edible film

Edible Films	Characteristics of edible films	Source
Hydrocolloid (Protein, pectin, alginate, carbohydrates)	<ul style="list-style-type: none"> advantages in the form of high adhesion to packaged food ingredients, selective to oxygen and carbon dioxide The weakness is that it is stiff and easily cracked 	Dewi et al. [16]
Lipids (fatty acids, fatty waxes)	<ul style="list-style-type: none"> The advantages are good moisture barrier properties (moisture). brittle weakness 	Abhishaben et al. [17]
Composite (Combination of hydrocolloids and lipids)	increasing the advantages and overcoming the disadvantages of hydrocolloid and lipid films	Murdinah et al. [18]



(a)



(b)

Fig. 1 Chemical structure of Amylose (a) and Amylopectin (b)

Source: Hassan et al. [20]




Fig. 2 Chitosan

Source: Rosida et al. [12]

Starch also has an important role in making edible films, both as a binder and thickener Herawati et al. [21]. Starch is composed of the components amylose and amylopectin. Due to the presence of amylose and amylopectin in starch, where amylopectin gives sticky properties and amylose gives gel properties Mustapa et al. [15]. One of the raw materials that has the

potential to be used as starch is starch from canna tubers, because canna starch has good physical and chemical properties, especially in a balanced amylose and amylopectin content, namely 24% amylose and 76% amylopectin (Santoso et al. [22] in Santoso et al. [23]). Based on research conducted by Radnaduhita et al [24,25] the use of gathot starch in chicken

sausage edible coatings obtained a pH value that was quite good even though it was not optimal, and the water content value still met the Indonesian national standard, namely a maximum of 67%.

4. NANO CHITOSAN

Chitosan or in chemistry it is called a unit (Poly-β-(1,4)-D glucosamine) is a carbohydrate polymer derived from deacetylation of chitin which is a natural biopolymer which has the second most abundant amount in nature after cellulose. Rumengan et al. [26]. The biggest chitosan producer is made from shrimp and crab shells. Chitosan is generally used in the industrial, pharmaceutical, food and cosmetic fields for several other uses, including as a stabilizer, thickener, texturizer and gelling agent. Mustapa et al. [15].

Chitosan has potential as an edible film because it has good physico-chemical properties, is non-toxic, has good antimicrobial properties (Ningrum et al. 2019). The antimicrobial activity of chitosan depends on several factors such as molecular weight (MW), degree of deacetylation (DDA), conformational structure, interaction between the positive charge (position C-2 of the glucosamine monomer) polymer, and the anionic component of chitosan Monsalve et al. [27].

Nanotechnology can extend the shelf life besides that nanotechnology can also detect the presence of pathogens/pesticides/poisons. Many chitosan studies have been carried out by modifying both chemically by increasing the degree of deacetylation, and physically by changing the size of chitosan into nanochitosan which has better absorption capacity in acting as an antibacterial and antifungal compared to the ability of chitosan which has a regular size. Komariah [28]. Based on the statement of Mohanraj et al [29], in general, nanoparticles have sizes ranging from 1-1000 nm. Synthesis of nano-chitosan can be carried out by several methods including ionic gelation and sonochemical methods Putri et al. [4]. *Edible Films* Alemán et al. 2016 in Monsalve et al. [27]. Based on other studies, it can be stated that chitosan can produce antibacterial activity in the form of higher gram-negative bacteria by decreasing the degree of deacetylation, molecular weight, and pH of chitosan Younes et al. 2014 in Rumengan et al. [26].

5. FISH OIL

Fish oil is one of the essential oils that has many important nutrients needed by the human body,

because fish oil contains about 75% unsaturated fatty acids and 25% saturated fatty acids. Yulianto et al. [30]. Fish fat or oil has special features when viewed from the composition of the fatty acid content, because fish oil has polyunsaturated fatty acids. *polyunsaturated fatty acids*(PUFA) in large quantities, which are known as omega-6, omega-3, Pandiangan et al. [31]. Palmitoleic, lauric acid, eicosapentaenoic (EPA), decosahexaenoc (DHA), arachidonic acid (ARA), alpha linolenic acid (ALA), and lauric acid(LA) are some examples of essential fatty acids needed by the body which can be found in salmon fish oil.

Apart from being beneficial for health, fish oil can also be used as a constituent of edible films, as according to Abhishaben et al. [32]. Lipid-based edible films (fish oil) exhibit good moisture barrier properties of water vapor. Based on research Santoso et al. [3] that the addition of unsaturated fatty acids to the solution had a significant effect on reducing the rate of water vapor transmission of the edible film. Generally fish oil extraction was carried out using the wet rendering method using solvent or steam (steam) and dry rendering without using a solvent Rozi et al. [33] Below is a table of omega-3 fatty acids from various sources.

Table 2. Sources of omega-3 fatty acids

Source	The amount of omega-3 content
Mackerel	2.5 gr
Herring	1.7 gr
Salmon	1.2 gr
Crustaceans/lobsters	0.2 gr
Squid	0.6 gr
salmon oil	19.9 gr
Cod liver oil	18.5 gr
Herring oil	11.4 gr

Source: Hope 2007 in Diana 2012 [34]

6. PATIN FISH SAUSAGE

Fish Sausage is a processed fishery product that uses raw materials in the form of crushed fish meat or surimi, at least 50%, which is mixed with flour and other additives, then put into sausage casings and undergoes a boiling or steaming process. Indonesian National Standards Agency [35].

Fish meat has several advantages, namely having fiber that is more easily digested, and has good fat content. However, sausages made from fish meat have deficiencies in perishable food properties or are prone to deterioration in

quality in a relatively short time Utami et al. [1]. Because fish meat has a high water content, meat with a high water content is a good substrate for bacterial growth because the bacteria's need for compounds can be fulfilled, which will later become a source of carbon, nitrogen, and other needs Pradana et al. [36].

Generally nonmicrobial spoilageThe most common occurrence in sausages is oxidation, which leads to the use of chemicals such as sulfur dioxide or synthetic antioxidants, such as butylated hydroxytoluene (BHT), propyl gallate (PG), and butylated hydroxyanisole (BHA) to prevent oxidative reactions in sausages. Conversely, ingredients added to sausages such as TVP, ISP, flour, and other vegetable proteins result in a decrease in the quality and shelf life of sausages. The decline in the quality of sausages can be seen by the presence of mucus, water, and milk Pradana et al. [36]. Therefore a way is needed to be able to maintain the quality of the sausage, namely by using packaging that can inhibit the decline in the quality of the sausage, one of which is by using edible film. Based on the research conducted Freitas et al. [37] by adding omega-3 fish oil to an edible film with a concentration of omega-3 (25%, 40%), the results were in the form of an edible film color that was acceptable to the panelists. Based on Purwamatja et al [38] the addition of chitosan solution can improve the texture of chicken meatballs.

7. PLASTICIZER

Plasticizers is one of the other components that make up edible films, in the form of low molecular weight compounds that have the benefit of softening the structure of edible films Rosida et al. [9]. Plasticizers have a major role in increasing film flexibility, reducing porosity, cracking tendency and reducing brittleness Sofia et al. [39]. Plasticizers or commonly called plasticizers are added to the components of the edible film to improve the mechanical properties of the edible film. According to Putra et al. [40] the plasticizer has a function in reducing the stiffness of the polymer so that a flexible and elastic edible film is obtained.

Sorbitol, mannitol, and polyethylene glycol are the most commonly used plasticizers in edible films, but polyols such as glycerol are the most common due to their stability and compatibility with hydrophilic compounds Monsalve et al. [27].

The glycerol plasticizer functions to increase elasticity by reducing the degree of hydrogen bonding and increasing the distance between the molecules of the polymer. Glycerol is a trihydroxy alcohol consisting of three carbon atoms. So, each carbon atom has an -OH group. Glycerol is a hydrophilic molecule that is relatively small and easily inserted between protein chains and forms hydrogen bonds with gluten groups and proteins Rosida et al. [9].

8. CHARACTERISTICS OF SAUSAGES TO KNOW THE POTENTIAL OF EDIBLE FILM IN PATIN FISH SAUSAGES

8.1 pH

The measuring instrument used in measuring pH is a pH meter, a pH meter is a measuring tool that can provide information about the degree of acidity of a solution Hadiatna and Susana [41]. The degree of acidity can experience a change in value at a certain time. This change occurs erratically due to the influence of several factors. These factors include temperature, and the decomposition process of organic matter. On the pH meter there is a probe made of a non-conducting glass cylinder that functions as a sensor. This tool is able to measure the acidity contained in water, by utilizing the HCL compound which soaks the electrode wire (Onny 2017 in Hadiatna and Susana [41].

pH analysis needs to be done because sausage is a food product that has characteristics suitable for microbial growth, so it is susceptible to damage. Sausages are made from meat raw materials which have a high water content of 68-75%, are rich in minerals, contain easily fermentable carbohydrates, and have a pH that allows for the growth of microorganisms, so one way to determine the decline in the quality of sausages is by measuring Soeparno's 2010 in Hawa et al. [42].

8.2 Water Content

Water content is the amount of water that is in an object, such as rocks, soil, agricultural materials, foodstuffs, food products, and so on Prasetyo et al. [43]. In foodstuffs, the presence of water content is usually associated with quality and is a measure of whether the ingredients are dry or solid. Other benefits of the presence of water in a material include as an index of stability during the shelf life, as well as a determinant of organoleptic

quality, especially in taste, texture, and tenderness Winarno [44].

Moisture content test is a very important test method in a food industry which functions to determine the quality and food resistance to possible damage. There are several methods that can be used to measure the water content of food, namely: distillation method (thermogravimetric), drying method, physical method and chemical method (*Karl Fisher Method*) Daud et al. [45].

8.3 Sensory

Sensory characterization is a test that is carried out by using the human senses as the main tool for measuring the acceptability of a product. Sensory test is a test that is very important to do, especially in the agricultural and food product industry. According to Zhao et al. [46] sensory tests provide very thorough assessment results, even in some assessments using sensory tests provide very thorough assessment results.

Based on Gonçalves et al. [47] sensory test is divided into three groups, namely scale test and descriptive test, difference test (*Different test*), and acceptance test (*Preference Test*). Based on Permady et al. [48] descriptive test consisting of a scoring or scaling test, scoring test or scoring is one of the tests in organoleptic characterization that uses panelists to provide an assessment of the samples tested by measuring and comparing similar products Tarjoko et al. [49]. In the scoring method, numbers are used to assess the intensity of a product, either in increasing or decreasing order.

In the assessment of food ingredients, the characteristic that determines whether a product is acceptable or not is its sensory properties. The senses used in assessing sensory properties are the senses of sight (appearance), touch (texture), smell (aroma) and taste (taste) Suryono et al. [42].

9. CONCLUSION

Catfish has the potential to be used as raw material for sausages due to its delicious taste and good fatty acid content. Deterioration in quality of catfish sausages can be inhibited by using edible film as a substitute for sausage casings. The addition of fish oil concentrations to canna starch and nanochitosan edible films can have a better effect on the organoleptic, pH, and water content of the sausages.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Utami R, Agustini T, Amalia U. Application of Edible Coating Semi Refined Carrageenan on the Shelf Life of Kurisi Fish Sausage (*Nemipterus Nematophorus*) at Cold Storage. 2017; 6(2), 24–32.
2. Balti R, et al. Development and characterization of bioactive edible films from spider crab (*Maja crispata*) chitosan incorporated with spirulina extract. *International Journal of Biological Macromolecules*. 2017.
3. Santoso B, Amilita D, Priyanto G. Development of Composite Edible Film Based on Corn Starch with the Addition of Palm Oil and Tween 20. 2018; 38(2), 119–124.
4. Putri A, Sundaryono A, Candra I. Characterization of Chitosan Nanoparticles from Sweet Potato Leaf Extract (*Ipomoea batatas* L) Using Ionic Gelation Method. *ALOTROP, Journal of Education and Chemistry*, 2018; 2(2), 203–207.
5. Mekawati, Fachriyah E, Sumardjo D. Application of Chitosan Transformation of Shrimp Waste Chitin (*Penaeus merguensis*) for Adsorption of Lead Metal Ions. *Journal of Science and Mathematics*, 2000; 8(2), 51–54.
6. Kusumawati D.H, Putri W D Physical And Chemical Characteristics Of Corn Starch Edible Film Incorporated With Black Ginger Juice. *Journal of Food and Agroindustry*. 2013;1(1):90-100.
7. Sariyanti I, Utami P. Utilization of Ganyong Starch (*Canna Edulis*) as a Raw Material for Color Barriers in Fabric. 2018;35(2); 67–74.
8. Apituley D. A, Sormin R, Nanlohy E. Characteristics and Fatty Acid Profiles of Fish Oil from Head and Bones of Tuna (*Thunnus albacares*). *AGRITEKNO: Journal of Agricultural Technology*. 2020; 9(1); 10-19.
9. Leger T, Azarnoush K, Traoré A, Cassagnes L, Rigaudière J, Jouve C, Pagès G, Bouvier D, Sapin V, Pereira B, Bonny J, Demaison L. Antioxidant and Cardioprotective Effects of EPA on Early Low-Severity Sepsis through UCP3 and SIRT3 Upholding of the Mitochondrial Redox Potential; 2019.

10. Endo, J., dan M. Arita. 2016. Cardioprotective Mechanism of Omega-3 Polyunsaturated Fatty Acids. *Journal of Cardiology*. Volume 67 (1): 22-27.
11. Suryono C, Ningrum L, Dewi TR Descriptive Likelihood and Organoleptic Tests for 5 Packaging and Seribu Islands Products. *Tourism Journal*. 2018; 5(2); 95–106.
12. Rosida D. F, Hapsari N, Dewati R. Edible Coatings and Films from Renewable Natural Biopolymer Materials. 2018.
13. Widodo L. U, Wati S. N, Vivi AP Making Edible Film from Yellow Pumpkin and Chitosan with Glycerol as a Plasticizer. *Journal of Food Technology*. 2019; 13(1); 59–65.
14. Febianti, M., Ghozali, A. A., Redjeki, S dan Iriani, I. 2020. Edible Film dari Tepung Kappa Karagenan dan Kitosan Cangkang Rajungan dengan Gliserol. *ChemPro*, 1(01): 16–21.
15. Mustapa R, Fajar R Raswen E. Utilization of Chitosan as a Basic Material for Making Edible Films from Yellow Sweet Potato Starch. 2017. 25; 2896–2896.
16. Dewi R, Rahmi R, Nasrun N. Improvement of Mechanical Properties and Water Vapor Transmission Rates of Edible Bioplastic Films Using Palm Oil and Sago Starch-Based Glycerol Plasticizers. *Unimal Chemical Technology Journal*. 2021; 8(1); 61.
17. Abhishaben M. S, Chandegara V. K, Jithender B, Pankajkumar MS Whey Protein Isolate based Biodegradable Food Packaging Film as affected by Protein to Glycerol Ratio, pH and Sonication Amplitude. *International Journal of Current Microbiology and Applied Sciences*. 2019; 8(03), 895–909.
18. Murdinah, Darmawan M, Fransiska D.Characteristics of Edible Film From Composite Alginate, Gluten And Beeswax (Beeswax). *Journal of Postharvest and Marine and Fisheries Biotechnology*. 2007; 2(1).
19. Nofiandi D, Ningsih W, Putri ASL Manufacture and Characterization of Edible Film from Breadfruit Starch Polyblend-Polyvinyl Alcohol with Propylene glycol as a Plasticizer. *Catalyst Journal*. 2016; 1(2); 1–12.
20. Hassan B, Ali S, Chatha S, Hussain A. I, Zia K. M, Akhtar N. Highlights SC. *International Journal of Biological Macromolecules*. 2017.
21. Herawati H. Potential of hydrocolloids as additives in quality food and non-food products. 2018.
22. Santoso B, Pratama F, Hamzah B, Pambayun R. Physical and chemical characteristics of canna and gadung starch modified by the crosslinking method. *Agritech Journal*. 2015; 35(03); 273.
23. Santoso A. F, Fibrianto K. The Effect of Red Dragon Fruit Peel Extract (*Hylocereus Polyrhizus*) on the Quality of Chicken Sausage. 2017; 5(4); 92–96.
24. Radnadhita A, Nuhriawangsa A, Kartikasari L. Antioxidant Activation Application of Gathot Flour (fermented cassava) in Edible Film Chicken Sausage at Room Temperature. *Livestock and Animal research*. 2021; 19(2): 227-237.
25. Radnadhita A, Nuhriawangsa A, Kartikasari L. Antioxidant Activation Application of Gathot Flour (fermented cassava) in Edible Film Chicken Sausage at Room Temperature. *Livestock and Animal research*. 2021; 19(2): 227-237.
26. Rumengan I, Suptijah P, Salindeho N, Stenly W, Luntungan A. Nanochitosan from fish scales: its application as a packaging for fishery products. 2018.
27. Monsalve B. J, Ramírez-toro, C, Bolívar G. Trends in Food Science & Technology Mechanisms of action of novel ingredients used in edible films to preserve microbial quality and oxidative stability in sausages - A review. *Trends in Food Science & Technology*. 2019; 89(May); 100–109.
28. Komariah A. Staphylococcus aureus (in vitro) Antibacterial Activity of Nano Chitosan on Staphylococcus aureus. *National Seminar on Biology Education XI FKIP UNS*. 2014; 11(1); 371–377.
29. Mohanraj VJ, Chen Y. Nanoparticles : A Review. *Tropical Journal of Pharmaceutical Research*. 2006; 5(1).
30. Yulianto A, Nugroho I, Swandari M. Formulation Of Emulsion Gourami Fish Oil (*Osphronemus gourami* L.). *Pharmaceutical Scientific Journal*. 2019; 38–43.
31. Pandiangan M, Kaban, J., Wirjosentono, B and Silalahi, J. 2019. Analysis of Omega 3 and Omega 6 Fatty Acid Content in Goldfish Oil (*Cyprinus Carpio*). *Talent Conference Series: Science and Technology (ST)*, 2(1), 37–44.
32. Abhishaben M. S, Chandegara V. K, Jithender B, Pankajkumar MS Whey Protein Isolate based Biodegradable Food

- Packaging Film as affected by Protein to Glycerol Ratio, pH and Sonication Amplitude. International Journal of Current Microbiology and Applied Sciences. 2019; 8(03); 895–909
33. Rozi A, et.al. Bintang Visitama Spirituality: The Power of Entrepreneurship. Attack. 2019.
34. Diana M Omega 3. Journal of Public Health. 2012; 6(2)
35. Indonesian National Standard Agency. 2013. SNI Fish Sausage. 11.
36. Pradana A, Hermanianto J, Profesional M, Food T and Agriculture FT Use of T VP and Pasteurization Application in Making Fried Beef Sausage at PT. X The Use of TVP and Application of Pasteurization in Production of Fried Beef Sausage in PT. X. 2019; 6(2), 99–107.
37. Freitas J, São B, De Medeiros, Cristina F and Oliveira E. Development and characterization of active film with omega-3 as a proposal for enrichment of butter Development and characterization of active film with omega-3 as a proposal for enrichment of butter. 2019.
38. Purwatmaja A, Widati Aris, Widyastuti E. Effect of Soaking Chicken Meatballs in Chitosan Solution in terms of Microbiological and Physical Quality. 2013.
39. Sofia I, Murdiningsih H, Yanti N. Manufacture and Study of Physicochemical, Mechanical, and Functional Properties of Edible Film from Chitosan Tiger Shrimp. Journal of Renewable Natural Materials. 2017; 5(2); 54–60.
40. Putra A. D, Johan V. S, Efendi R. Addition of sorbitol as a plasticizer in making breadfruit starch edible films. Come on Faculty of Agriculture. 2017; 4(2); 1–15.
41. Hadiatna F, Susana R. Design of a Smart pH Meter as a Measuring Tool for Monitoring Nutrient Solutions. 2019; 7(2); 404–414.
42. Hawa C, Ginting U. Y, Susilo B, Wiganti L. Physicochemical Study of Edible Chicken Feet Gelatin-Based Sausage Casings. Agointek. 2020; 14(2); 67–74.
43. Prasetyo T, Firas A, Sujadi H. Implementation of Moisture Detection Devices in Foods Based on the Internet of Things. 2019; 5(2); 81–96.
44. Winarno FG Food Chemistry and Nutrition. PT Gramedia Pustaka Utama. Jakarta. 2004.
45. Daud A, Suriat, Nuzulyanti. Study of the Application of Factors Affecting Determination Accuracy. LUTJANUS. 2020; 11–16.
46. Zhao G, Zhan C. F, He R. Research on Data Mining Methods for Organoleptic Determination of Amomum Villosum Product. IEEE Int. Conf. Bioinform. biomed. Work. BIBMW. 2011; 873–880.
47. Gonçalves R, Hester J, Carvalho N, Pinho P, Tentzeris M. Passive Sensors for Food Quality Monitoring and Counterfeiting. 2014; 4–7.
48. Permadi M. R, Oktafa H, Agustianto K. Design of a Food Sensory Test System with Preference Test (Hedonic and Hedonic Quality) Tests, Case Study of Bread, Using Radial Basis Function Network Algorithm. MIKROTIK: Journal of Informatics Management. 2018; 8(1); 29–42.
49. Tarjoko, Suyono, Yulia, Anjasari LN Application of Healthy Kitchens and Use of Natural Laru to Improve the Quality of Coconut Sugar. SOLMA Journal. 2019; 08(01); 39–46.

© 2023 Rinaldi et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/97857>